

# Inverse Scattering Construction of a Dipole Black Ring

Jorge V. Rocha<sup>‡</sup>, Maria J. Rodriguez<sup>#</sup>, and Amitabh Virmani<sup>◇</sup>

<sup>‡</sup> *Centro Multidisciplinar de Astrofísica - CENTRA,  
Dept. de Física, Instituto Superior Técnico, Technical University of Lisbon,  
Av. Rovisco Pais 1, 1049-001 Lisboa, Portugal  
jorge.v.rocha@ist.utl.pt*

<sup>#</sup> *Max-Planck-Institut für Gravitationsphysik,  
Albert-Einstein-Institut, 14476 Golm, Germany  
maria.rodriguez@aei.mpg.de*

<sup>◇</sup> *Physique Théorique et Mathématique,  
Université Libre de Bruxelles and International Solvay Institutes  
Campus Plaine C.P. 231, B-1050 Bruxelles, Belgium  
avirmani@ulb.ac.be*

## Abstract

Using the inverse scattering method in six dimensions we construct the dipole black ring of five dimensional Einstein-Maxwell-dilaton theory with dilaton coupling  $a = 2\sqrt{2/3}$ . The 5d theory can be thought of as the NS sector of low energy string theory in Einstein frame. It can also be obtained by dimensionally reducing six-dimensional vacuum gravity on a circle. Our new approach uses  $GL(4, \mathbb{R})$  integrability structure of the theory inherited from six-dimensional vacuum gravity. Our approach is also general enough to potentially generate dipole black objects carrying multiple rotations as well as more exotic multi-horizon configurations.

## Contents

<b>1</b>	<b>Introduction</b>	<b>2</b>
<b>2</b>	<b>Inverse Scattering Construction of a Dipole Black Ring</b>	<b>3</b>
2.1	The Set-Up . . . . .	3
2.2	Seed and Soliton Transformations . . . . .	4
2.3	Dipole Black Ring Solution . . . . .	6
<b>3</b>	<b>Discussion and Outlook</b>	<b>9</b>

## 1 Introduction

Almost a decade after the first black ring solution was written down by Emparan and Reall [1], black rings are still a continual source of excitement in higher dimensional general relativity and in microscopic description of black holes in string theory; see [2, 3, 4] for reviews. A technical problem in the black ring literature, whose solution is still elusive, is the construction of the most general asymptotically flat black ring in a simple five-dimensional supergravity theory – a five-parameter black ring solution carrying mass, two angular momenta, electric charge and magnetic dipole charge. Not only this solution is not known, but also the requisite set of techniques to carry out a systematic construction are not fully understood. The most important stumbling block in these considerations is the dipole charge. The original dipole solution [5] was constructed using educated guesswork. There also exists in the literature an algorithmic construction of a black ring solution carrying dipole charge [6]. However, it cannot be employed to generate multiple rotations. In this paper we remedy this situation for Einstein-Maxwell-dilaton theory with dilaton coupling  $a = 2\sqrt{2/3}$ . This theory is obtained by dimensionally reducing six-dimensional vacuum gravity on a circle. It can also be thought of as the NS sector of low energy string theory in Einstein frame. We exploit  $GL(4, \mathbb{R})$  integrability structure of the theory inherited from six-dimensional vacuum gravity and thus employ the inverse scattering method in six dimensions. A key point in this procedure is the identification of the seed that can be used to generate the dipole charge.

The new construction presented here below can in principle be used to construct the doubly spinning dipole black ring and/or the most general black ring in this theory. This situation is in contrast with the previous approaches, e.g., [6], which rely on a specific  $SL(2, \mathbb{R}) \times SL(2, \mathbb{R})$  structure, and cannot be extended in a natural way to construct, say, the doubly spinning dipole black ring. Our approach does not rely on  $SL(2, \mathbb{R}) \times SL(2, \mathbb{R})$  structure, but rather uses the full  $GL(4, \mathbb{R})$  symmetry of the theory. For a discussion on using integrability techniques

to construct solutions of supergravity theories we refer the reader to [7, 8]. In the next section we present our construction; in the following section we conclude with some observations and comments.

## 2 Inverse Scattering Construction of a Dipole Black Ring

This section is the core of the paper. We first present our setup and then the construction of the dipole ring solution.

### 2.1 The Set-Up

The action of the five-dimensional theory we are interested in is

$$S = \frac{1}{16\pi G} \int d^5x \sqrt{-g} \left( R - \frac{1}{2} \partial_\mu \phi \partial^\mu \phi - \frac{1}{4} e^{-a\phi} F_{\mu\nu} F^{\mu\nu} \right), \quad (2.1)$$

with

$$a = \frac{2\sqrt{2}}{\sqrt{3}}. \quad (2.2)$$

This action can also be obtained by the circle reduction of six-dimensional vacuum gravity using the ansatz

$$ds_6^2 = e^{\frac{\phi}{\sqrt{6}}} ds_5^2 + e^{-\frac{\sqrt{3}\phi}{\sqrt{2}}} (dw + A)^2. \quad (2.3)$$

The sixth dimension is parametrized by  $w$  and  $F = dA$ . The five-dimensional theory naturally supports a magnetic one-brane, an electric zero-brane (and smeared versions thereof), and dipole black rings. In the literature solutions of the ‘electric’ version of the theory under consideration are studied in greater detail [9]. This is obtained by defining

$$H = e^{-a\phi} \star_5 F, \quad H = dB, \quad (2.4)$$

then

$$S = \frac{1}{16\pi G} \int d^5x \sqrt{-g} \left( R - \frac{1}{2} \partial_\mu \phi \partial^\mu \phi - \frac{1}{12} e^{a\phi} H_{\mu\nu\rho} H^{\mu\nu\rho} \right). \quad (2.5)$$

The action (2.5) can be interpreted as the NS sector of low energy string theory in Einstein frame. The fundamental string is a solution of this action. The dilaton  $\sigma$  of string theory is  $\sigma = -\sqrt{\frac{3}{8}}\phi$  and the string metric is  $g_{\mu\nu}^{(s)} = e^{-\sqrt{\frac{2}{3}}\phi} g_{\mu\nu}$ . Via string dualities solutions of (2.5) can be related to other single brane configurations.

In the following we restrict our attention to the ‘magnetic’ version (2.1) and its uplift to six-dimensional vacuum gravity via (2.3).




Figure 1: The figure depicts sources for the seed metric  $G_0$ . The rods are located in the  $(\rho, z)$  space at the  $z$ -axis with  $\rho = 0$ . The solid rods have positive density and the dashed rod has negative density. They add up to an infinite rod with uniform density such that  $\det G_0 = -\rho^2$ .

## 2.2 Seed and Soliton Transformations

The inverse scattering method has been widely used to construct analytically many of currently available asymptotically flat black hole solutions in four and five dimensions. The construction consists in a series of soliton transformations on a seed solution. Through these ‘dressing’ transformations, or Belinsky-Zakharov (BZ) procedure [10], a more general solution can be generated exhibiting new free parameters. We use this method below. For our analysis we also make use of the results of higher-dimensional Weyl and rod diagram representations from [11, 12, 13].

The rod configuration for the seed solution of dipole black ring when lifted to six-dimensions is shown in figure 1. The thick solid lines correspond to rod sources of uniform density  $+1/2$ , and the dashed line corresponds to a rod source of uniform density  $-1/2$ . The rod in the  $t$  direction corresponds to the horizon. In figure 1 when  $a_0 = a_1$  and  $a_4 = a_2$  the negative density rod and the rod in the  $w$  direction are eliminated and the solution describes a neutral static black ring multiplied with the flat direction  $w$ . This is an unbalanced configuration. The negative density rod is included in the seed following [14] to facilitate adding the  $S^1$  angular momentum to the ring. The positive density rod  $[a_2, a_4]$  along the  $w$  direction is included to facilitate adding dipole charge to the ring.

The seed metric corresponding to the rod configuration of figure 1 is given by

$$ds_6^2 = (G_0)_{ab} dx^a dx^b + e^{2\nu_0} (d\rho^2 + dz^2), \quad (2.6)$$

where

$$G_0 = \text{diag} \left\{ -\frac{\mu_0}{\mu_2}, \frac{\rho^2 \mu_4}{\mu_1 \mu_3}, \frac{\mu_1 \mu_3}{\mu_0}, \frac{\mu_2}{\mu_4} \right\}, \quad \det G_0 = -\rho^2, \quad (2.7)$$

and the conformal factor of the seed is

$$e^{2\nu_0} = k^2 \frac{\mu_1 \mu_3}{\mu_0} \frac{(\mu_0 \mu_1 + \rho^2)(\mu_0 \mu_2 + \rho^2)(\mu_0 \mu_3 + \rho^2)(\mu_1 \mu_4 + \rho^2)(\mu_2 \mu_4 + \rho^2)(\mu_3 \mu_4 + \rho^2)}{(\mu_1 \mu_3 + \rho^2)^2 \prod_{i=0}^4 (\mu_i^2 + \rho^2)}. \quad (2.8)$$

The integration constant  $k$  will be fixed in the next subsection. Our ordering of coordinates is  $x^a = (t, \phi, \psi, w)$ , with  $t$  corresponding to the timelike coordinate,  $\phi$  describing the azimuthal angle on the  $S^2$  and  $\psi$  being the angle along the  $S^1$  component of the ring. The  $\mu_i$ 's are pole trajectories of the dressing matrices, commonly referred to as solitons,  $\mu_i = \sqrt{\rho^2 + (z - a_i)^2} - (z - a_i)$ , and  $a_i$  are the rod endpoints. In writing these expressions we have followed established conventions from the literature. We refer the reader to e.g. [3] for notational details.

We assume the ordering

$$a_0 \leq a_1 \leq a_2 \leq a_4 \leq a_3. \quad (2.9)$$

The labeling (and hence the ordering) is a little unusual, but is motivated to simplify the presentation of solution after the inverse scattering transformation. We will see that the endpoints  $a_0$  and  $a_4$  will not appear in the rod diagram of the final solution.

The solution (2.7) and (2.8) with ordering (2.9) is singular and not in itself of direct physical interest. First, with a 1-soliton transformation we add an anti-soliton which mixes  $t$  and  $\psi$  directions in such a way that the negative density rod  $[a_0, a_1]$  moves in the  $t$  direction and cancels the segment  $[a_0, a_1]$  of the positive density rod. Taking  $a_4 = a_2$  at this stage gives the singly spinning black ring of [1]; this step is the six dimensional analogue of the one used in [14]. Next, keeping  $a_2 \leq a_4$ , we apply a further 1-soliton transformation that gives the dipole black ring. This latter 1-soliton transformation mixes the  $w$  and  $\phi$  directions in such a way that the positive density rod  $[a_2, a_4]$  moves in the  $\phi$  direction and *merges* with the positive density rod  $[a_4, a_3]$ . The first transformation leaves behind naked singularities at  $z = a_0$ , but choosing the corresponding BZ vector appropriately completely eliminates this singularity. Requiring the final rod  $[a_2, a_3]$  to have uniform rod direction fixes the second BZ vector.

In more detail the steps for generating the dipole ring solution by a 2-soliton transformation are as follows:

1. Perform two 1-soliton transformations on the seed solution (2.7) followed by a rescaling. More precisely (i) remove an anti-soliton at  $z = a_0$  with trivial BZ vector  $(1, 0, 0, 0)$ , (ii) remove a soliton at  $z = a_4$  with trivial BZ vector  $(0, 0, 0, 1)$ , (iii) rescale the metric by  $\mu_0/\mu_4$ . The resulting metric is

$$G'_0 = \text{diag} \left\{ -\frac{\bar{\mu}_4}{\mu_2}, -\frac{\mu_0 \bar{\mu}_3}{\mu_1}, \frac{\mu_1 \bar{\mu}_4}{\bar{\mu}_3}, \frac{\mu_0}{\bar{\mu}_2} \right\}. \quad (2.10)$$

This metric will be the seed for our next soliton transformation.

2. The generating matrix corresponding to (2.10) is:

$$\psi'_0(\lambda, \rho, z) = \text{diag} \left\{ -\frac{\bar{\mu}_4 - \lambda}{\mu_2 - \lambda}, -\frac{(\mu_0 - \lambda)(\bar{\mu}_3 - \lambda)}{\mu_1 - \lambda}, \frac{(\mu_1 - \lambda)(\bar{\mu}_4 - \lambda)}{\bar{\mu}_3 - \lambda}, \frac{\mu_0 - \lambda}{\bar{\mu}_2 - \lambda} \right\}. \quad (2.11)$$

3. Perform now a 2-soliton transformation with  $G'_0$  as seed and undo the rescaling. More precisely (i) add an anti-soliton at  $z = a_0$  with BZ vector  $(1, 0, c_1, 0)$ , (ii) add a soliton at  $z = a_4$  with BZ vector  $(0, c_2, 0, 1)$ , and (iii) rescale by  $\mu_4/\mu_0$ . Denote the final metric by  $G$ . The final rescaling ensures that  $\det G = -\rho^2$ .
4. Construct  $e^{2\nu}$ . The result  $(e^{2\nu}, G)$  is the six-dimensional solution we want. Appropriately tuning  $c_1$  and  $c_2$  and Kaluza-Klein reducing this solution along the  $w$  direction we obtain the smooth five-dimensional dipole black ring solution of the theory under consideration.

In the next subsection we first present the resulting solution in the Weyl coordinates and analyze the rod structure and regularity of the solution. We then transform it into the more convenient  $(x, y)$  coordinates. The resulting metric is shown to be identical to the one presented in [5].

### 2.3 Dipole Black Ring Solution

The  $G_{tt}$ ,  $G_{t\psi}$  and  $G_{\psi\psi}$  components of the metric after performing the above described inverse scattering transformations takes the form:

$$G_{tt} = -\frac{\mu_0 \left[ \mu_3(\mu_0\mu_1 + \rho^2)^2(\mu_0\mu_2 + \rho^2)^2 - c_1^2\mu_1\mu_2(\mu_0 - \mu_3)^2\rho^4 \right]}{\mu_2 \left[ \mu_3(\mu_0\mu_1 + \rho^2)^2(\mu_0\mu_2 + \rho^2)^2 + c_1^2\mu_0^2\mu_1\mu_2(\mu_0 - \mu_3)^2\rho^2 \right]}, \quad (2.12)$$

$$G_{t\psi} = -\frac{c_1\mu_1\mu_3(\mu_0 - \mu_3)(\mu_0^2 + \rho^2)(\mu_0\mu_1 + \rho^2)(\mu_0\mu_2 + \rho^2)}{\left[ \mu_3(\mu_0\mu_1 + \rho^2)^2(\mu_0\mu_2 + \rho^2)^2 + c_1^2\mu_0^2\mu_1\mu_2(\mu_0 - \mu_3)^2\rho^2 \right]}, \quad (2.13)$$

$$G_{\psi\psi} = \frac{\mu_1\mu_3 \left[ \mu_3(\mu_0\mu_1 + \rho^2)^2(\mu_0\mu_2 + \rho^2)^2 - c_1^2\mu_0^4\mu_1\mu_2(\mu_0 - \mu_3)^2 \right]}{\mu_0 \left[ \mu_3(\mu_0\mu_1 + \rho^2)^2(\mu_0\mu_2 + \rho^2)^2 + c_1^2\mu_0^2\mu_1\mu_2(\mu_0 - \mu_3)^2\rho^2 \right]}. \quad (2.14)$$

Similarly, the  $G_{\phi\phi}$ ,  $G_{w\phi}$  and  $G_{ww}$  components of the metric take the form:

$$G_{\phi\phi} = \frac{\rho^2 \left[ \mu_1\mu_4^2(\mu_2\mu_4 + \rho^2)^2(\mu_3\mu_4 + \rho^2)^2 + c_2^2\mu_2\mu_3(\mu_1 - \mu_4)^2\rho^6 \right]}{\mu_1\mu_3\mu_4 \left[ \mu_1(\mu_2\mu_4 + \rho^2)^2(\mu_3\mu_4 + \rho^2)^2 - c_2^2\mu_2\mu_3(\mu_1 - \mu_4)^2\rho^4 \right]}, \quad (2.15)$$

$$G_{w\phi} = \frac{c_2\mu_2(\mu_1 - \mu_4)\rho^2(\mu_4^2 + \rho^2)(\mu_2\mu_4 + \rho^2)(\mu_3\mu_4 + \rho^2)}{\mu_4 \left[ \mu_1(\mu_2\mu_4 + \rho^2)^2(\mu_3\mu_4 + \rho^2)^2 - c_2^2\mu_2\mu_3(\mu_1 - \mu_4)^2\rho^4 \right]}, \quad (2.16)$$

$$G_{ww} = \frac{\mu_2 \left[ \mu_1(\mu_2\mu_4 + \rho^2)^2(\mu_3\mu_4 + \rho^2)^2 + c_2^2\mu_2\mu_3\mu_4^2(\mu_1 - \mu_4)^2\rho^2 \right]}{\mu_4 \left[ \mu_1(\mu_2\mu_4 + \rho^2)^2(\mu_3\mu_4 + \rho^2)^2 - c_2^2\mu_2\mu_3(\mu_1 - \mu_4)^2\rho^4 \right]}. \quad (2.17)$$

The remaining components not related to these by symmetry vanish. Note that setting  $c_1 = c_2 = 0$  we get back the static seed metric (2.7). The final conformal factor reads as:

$$e^{2\nu} = k^2 \frac{(\mu_0\mu_3 + \rho^2)(\mu_1\mu_4 + \rho^2) \left[ M_{0,0} + c_1^2 M_{c_1,0} + c_2^2 M_{0,c_2} + c_1^2 c_2^2 M_{c_1,c_2} \right]}{\mu_0(\mu_0\mu_1 + \rho^2)(\mu_0\mu_2 + \rho^2)(\mu_1\mu_3 + \rho^2)^2(\mu_2\mu_4 + \rho^2)(\mu_3\mu_4 + \rho^2) \prod_{i=0}^4 (\mu_i^2 + \rho^2)} \quad (2.18)$$




Figure 2: The figure depicts sources for the six-dimensional lift of the dipole black ring. The direction for each rod is indicated. The points  $a_0$  and  $a_4$  are not indicated in the figure, as they no longer represent real turning points.

where

$$M_{0,0} = \mu_1\mu_3(\mu_0\mu_1 + \rho^2)^2(\mu_0\mu_2 + \rho^2)^2(\mu_2\mu_4 + \rho^2)^2(\mu_3\mu_4 + \rho^2)^2, \quad (2.19)$$

$$M_{c_1,0} = \mu_0^2\mu_1^2\mu_2(\mu_0 - \mu_3)^2\rho^2(\mu_2\mu_4 + \rho^2)^2(\mu_3\mu_4 + \rho^2)^2, \quad (2.20)$$

$$M_{0,c_2} = -\mu_2\mu_3^2(\mu_1 - \mu_4)^2\rho^4(\mu_0\mu_1 + \rho^2)^2(\mu_0\mu_2 + \rho^2)^2, \quad (2.21)$$

$$M_{c_1,c_2} = -\mu_0^2\mu_1\mu_2^2\mu_3(\mu_0 - \mu_3)^2(\mu_1 - \mu_4)^2\rho^6. \quad (2.22)$$

The metric functions at this stage look somewhat complicated. However, as we will shortly see, these functions dramatically simplify when we express them in  $(x, y)$  coordinates. To verify that the metric describes the singly spinning dipole black ring, we first analyze the rod structure of the solution. This analysis is straightforward [12]. We find

- The semi-infinite rod  $z \in (-\infty, a_0]$  has direction  $(0, 1, 0, 0)$ . The  $G_{\psi\psi}$  component of the metric however diverges as  $z \rightarrow a_0$  for general values of the BZ parameter  $c_1$ .
- Similarly the rod  $z \in [a_0, a_1]$  has direction  $(0, 1, 0, 0)$ . Again the  $G_{\psi\psi}$  component of the metric diverges as  $z \rightarrow a_0$  for general values of the BZ parameter  $c_1$ .

Fortunately, the singularities in both  $(-\infty, a_0]$  and  $[a_0, a_1]$  rods are completely removed by setting

$$|c_1| = \sqrt{\frac{2(a_1 - a_0)(a_2 - a_0)}{(a_3 - a_0)}}. \quad (2.23)$$

The condition (2.23) will be imposed from now onwards. Having imposed this condition we have effectively moved the negative density rod  $z \in [a_0, a_1]$  from the  $\psi$  direction to the  $t$  direction and have canceled it against the segment  $[a_0, a_1]$  of the rod  $[a_0, a_2]$ . In the process we have successfully generated rotation in the  $\psi$  direction as is illustrated by the rod direction of the rod  $[a_1, a_2]$ .

- The finite timelike rod  $[a_1, a_2]$  has rod direction  $(1, 0, \Omega_\psi, 0)$ , where

$$\Omega_\psi = \sqrt{\frac{(a_1 - a_0)}{2(a_2 - a_0)(a_3 - a_0)}}. \quad (2.24)$$

- The finite rod  $[a_2, a_4]$  has direction  $(0, 1, 0, \frac{2(a_4 - a_2)}{c_2})$ .
- The finite rod  $[a_4, a_3]$  has direction  $(0, 1, 0, \frac{(a_4 - a_1)c_2}{a_3 - a_4})$ .

Note that directions of rods  $[a_2, a_4]$  and  $[a_4, a_3]$  are in general different. However, if we choose

$$|c_2| = \sqrt{\frac{2(a_4 - a_2)(a_3 - a_4)}{(a_4 - a_1)}}, \quad (2.25)$$

then the two rod directions are identical. The condition (2.25) will also be imposed from now onwards. Having imposed this condition we have effectively displaced the positive density rod  $z \in [a_2, a_4]$  from the  $w$  direction towards the  $\phi$  direction and have merged it to the rod  $[a_4, a_3]$ . In the process we have successfully generated a dipole charge as illustrated by the rod direction  $(0, 1, 0, \Omega_w)$  of the rod  $[a_2, a_3]$ , where

$$\Omega_w = \sqrt{\frac{2(a_4 - a_2)(a_4 - a_1)}{(a_3 - a_4)}}. \quad (2.26)$$

- Finally, the semi-infinite rod  $z \in [a_3, \infty)$  has rod direction  $(0, 0, 1, 0)$ .

All of this discussion is succinctly illustrated in the rod diagram in figure 2.

To verify that the above described solution is indeed the singly spinning dipole black ring of [5] lifted to six dimensions, we now rewrite the metric in the ring coordinates  $(x, y)$ . First recall that the five dimensional fields for the dipole ring take the form [5]

$$\begin{aligned} ds_5^2 = & -\frac{F(y)}{F(x)} \left( \frac{H(x)}{H(y)} \right)^{1/3} \left( dt + C(\nu, \lambda) R \frac{1+y}{F(y)} d\psi \right)^2 \\ & + \frac{R^2}{(x-y)^2} F(x) (H(x)H(y)^2)^{1/3} \left[ -\frac{G(y)}{F(y)H(y)} d\psi^2 - \frac{dy^2}{G(y)} + \frac{dx^2}{G(x)} + \frac{G(x)}{F(x)H(x)} d\phi^2 \right] \end{aligned} \quad (2.27)$$

where

$$F(\xi) = 1 + \lambda\xi, \quad G(\xi) = (1 - \xi^2)(1 + \nu\xi), \quad H(\xi) = 1 - \mu\xi, \quad (2.28)$$

and

$$C(\nu, \lambda) = \sqrt{\lambda(\lambda - \nu) \frac{1 + \lambda}{1 - \lambda}}. \quad (2.29)$$

The vector field  $A$  and dilaton  $\phi$  take the following expressions:

$$A = C(\nu, -\mu) R \left( \frac{1+x}{H(x)} \right) d\phi, \quad e^{-\phi} = \left( \frac{H(x)}{H(y)} \right)^{\sqrt{2/3}}. \quad (2.30)$$



The six-dimensional lift can be readily obtained using the KK ansatz (2.3). We get

$$ds_6^2 = -\frac{F(y)}{F(x)} \left( dt + C(\nu, \lambda) R \frac{1+y}{F(y)} d\psi \right)^2 + \frac{H(x)}{H(y)} \left( dw + C(\nu, -\mu) R \frac{1+x}{H(x)} d\phi \right)^2 \quad (2.31)$$

$$+ \frac{R^2}{(x-y)^2} F(x) H(y) \left[ -\frac{G(y)}{F(y)H(y)} d\psi^2 - \frac{dy^2}{G(y)} + \frac{dx^2}{G(x)} + \frac{G(x)}{F(x)H(x)} d\phi^2 \right].$$

Note in particular that in six dimensions all fractional powers of metric functions have disappeared.

Our aim now is to show that the metric in Weyl coordinates (2.12)–(2.22) is identical to the metric (2.31). To establish this we follow the procedure of [12] and appendix A.2 of [14]. The coordinate transformation from Weyl  $(\rho, z)$  to ring coordinates  $(x, y)$  is

$$\rho = \frac{R^2 \sqrt{-G(x)G(y)}}{(x-y)^2}, \quad z = \frac{R^2(1-xy)[2+\nu(x+y)]}{2(x-y)^2}. \quad (2.32)$$

The rod end points are related to parameters  $\nu, \lambda, \mu$  as

$$a_0 = \frac{R^2}{2}\alpha, \quad a_1 = -\frac{R^2}{2}\nu, \quad a_2 = \frac{R^2}{2}\nu, \quad a_3 = \frac{R^2}{2}, \quad a_4 = \frac{R^2}{2}\beta. \quad (2.33)$$

Here  $-\infty < \alpha \leq -\nu$  and  $\nu \leq \beta < 1$  are constants to be determined. After a straightforward calculation we see that with the choice

$$\alpha = \frac{\nu(1+\lambda) - 2\lambda}{(1-\lambda)}, \quad \beta = \frac{\nu(1-\mu) + 2\mu}{(1+\mu)}, \quad k^2 = \frac{(1+\mu)(1-\lambda)}{(1-\nu)^2}, \quad (2.34)$$

all metric components match. The conditions  $-\infty < \alpha \leq -\nu$  and  $\nu \leq \beta < 1$  and the ordering of the rod end points imply that

$$0 < \nu \leq \lambda < 1 \quad \text{and} \quad 0 \leq \mu < 1. \quad (2.35)$$

We therefore have also recovered the correct bounds on the parameters of the singly spinning dipole black ring. The periodicities of angular coordinates are  $\Delta\phi = \Delta\psi = 2\pi k$ . With  $k$  given in (2.34), these periodicities also agree with the results of [5]. Thus we have shown that the solution (2.12)–(2.22) is identical to (2.31).

### 3 Discussion and Outlook

Using  $\text{GL}(4, \mathbb{R})$  integrability structure of the Einstein-Maxwell-dilaton theory with dilaton coupling  $a = 2\sqrt{2/3}$ , we presented an inverse scattering construction of a singly spinning dipole black ring solution. The finite rod in the Kaluza-Klein direction  $w$  plays the key role in our construction. Recall that finite rods in Kaluza-Klein directions correspond to KK bubbles [11]. Therefore, loosely said, KK bubble is converted into magnetic dipole in our inverse scattering construction.

The novel construction presented above can in principle be generalized to construct the doubly spinning dipole black ring and/or the most general black ring in this theory. We emphasize once again that this situation is in contrast with the previous approaches, e.g., [6], which rely on a specific  $SL(2, \mathbb{R}) \times SL(2, \mathbb{R})$  structure, and cannot be extended in a natural way to construct, say, the doubly spinning dipole black ring. Our approach does not rely on a  $SL(2, \mathbb{R}) \times SL(2, \mathbb{R})$  structure, but rather uses the full  $GL(4, \mathbb{R})$  symmetry of the theory.

The approach presented above has some resemblance to the approach taken in [15], where the first static self-gravitating loop of string was described. There, starting with the product of five dimensional Euclidean Schwarzschild and time, and performing a twisted KK reduction, a static ring solution was obtained. The solution is regular, but the horizon of the ring in that case is degenerate (extremal) with zero area. The solution is balanced not by rotation but by a background field. Due to the presence of the background field, the solution is not asymptotically flat, instead it approaches a fluxbrane. In our approach, since we mix the  $w$  and  $\phi$  directions via the inverse scattering method, we generate the five-dimensional magnetic charge without the need to do a twisted KK reduction. Thus, we obtain an asymptotically flat solution in five-dimensions instead of a fluxbrane. A related technical point that is worth mentioning is that in our generation of the singly spinning dipole ring it is not necessary to perform any mixing of the Killing coordinates to obtain the desired asymptotic structure.

Finally, in the process of constructing the dipole black ring we also found its six dimensional rod diagram description. A common picture seems to be that black hole solutions are uniquely characterized by their charges and rod diagrams [16]. We expect that the rod diagram description of the dipole black ring presented above will play an essential role in the uniqueness proofs within the described dilaton theory.

The new procedure implemented in this paper sets the building blocks for the construction of more general and exotic black hole solutions with dipole charges. One could add dipole charges to e.g. the bicycling black ring of [17] or a further rotation to the dipole black ring. In this work we have not attempted to construct the doubly spinning dipole solution, or the most general black ring solution, but we hope to return to these problems elsewhere.

## Acknowledgements

We thank Joan Camps, Pau Figueras and Oscar Varela for discussions. We specially thank Roberto Emparan for comments on a draft version and for bringing reference [15] to our attention as well as the organizers of the ‘Gravity’ workshop, 17-29 July 2011, held at Centro de Ciencias Pedro Pascual, Benasque, Spain where this work was completed. JVR is supported by *Fundação para a Ciência e Tecnologia* (FCT)-Portugal through contract no. SFRH/BPD/47332/2008. AV was supported by IISN Belgium (conventions 4.4511.06 and

4.4514.08) and by the Belgian Federal Science Policy Office through the Interuniversity Attraction Pole P6/11.

## References

- [1] R. Emparan, H. S. Reall, “A Rotating black ring solution in five-dimensions,” *Phys. Rev. Lett.* **88**, 101101 (2002). [hep-th/0110260].
- [2] R. Emparan and H. S. Reall, “Black rings,” *Class. Quant. Grav.* **23**, R169 (2006) [arXiv:hep-th/0608012].
- [3] R. Emparan, H. S. Reall, “Black Holes in Higher Dimensions,” *Living Rev. Rel.* **11**, 6 (2008). [arXiv:0801.3471 [hep-th]].
- [4] M. J. Rodriguez, “On the black hole species (by means of natural selection),” Invited review talk, Proceedings of the Twelfth Marcel Grossmann Meeting on General Relativity (2011). [arXiv:1003.2411 [hep-th]].
- [5] R. Emparan, “Rotating circular strings, and infinite nonuniqueness of black rings,” *JHEP* **0403**, 064 (2004). [hep-th/0402149].
- [6] S. S. Yazadjiev, “Solution generating in 5D Einstein-Maxwell-dilaton gravity and derivation of dipole black ring solutions,” *JHEP* **0607**, 036 (2006). [hep-th/0604140].  
S. S. Yazadjiev, “Completely integrable sector in 5-D Einstein-Maxwell gravity and derivation of the dipole black ring solutions,” *Phys. Rev.* **D73**, 104007 (2006). [hep-th/0602116].
- [7] P. Figueras, E. Jamsin, J. V. Rocha and A. Virmani, “Integrability of Five Dimensional Minimal Supergravity and Charged Rotating Black Holes,” *Class. Quant. Grav.* **27**, 135011 (2010) [arXiv:0912.3199 [hep-th]].
- [8] G. A. Alekseev, “Thirty years of studies of integrable reductions of Einstein’s field equations,” arXiv:1011.3846 [gr-qc]. Talk given at 12th Marcel Grossmann Meeting on General Relativity (MG 12), Paris, France, 12-18 Jul 2009.
- [9] T. Harmark and N. A. Obers, “Thermodynamics of spinning branes and their dual field theories,” *JHEP* **0001**, 008 (2000) [arXiv:hep-th/9910036].
- [10] V. A. Belinski and V. E. Zakharov, “Integration Of The Einstein Equations By The Inverse Scattering Problem Technique And The Calculation Of The Exact Soliton Solutions,” *Sov. Phys. JETP* **48**, 985 (1978) [*Zh. Eksp. Teor. Fiz.* **75**, 1953 (1978)].

- V. A. Belinski, V. E. Sakharov, “Stationary Gravitational Solitons with Axial Symmetry,” Sov. Phys. JETP **50** 1, 1979, [Zh.Eksp.Teor.Fiz. **77** 3, 1979].
- V. Belinski and E. Verdaguer, “Gravitational solitons,” Cambridge, UK: Univ. Pr. (2001).
- A. A. Pomeransky, “Complete integrability of higher-dimensional Einstein equations with additional symmetry, and rotating black holes,” Phys. Rev. D **73**, 044004 (2006) [arXiv:hep-th/0507250].
- [11] R. Emparan, H. S. Reall, “Generalized Weyl solutions,” Phys. Rev. **D65**, 084025 (2002). [hep-th/0110258].
- [12] T. Harmark, “Stationary and axisymmetric solutions of higher-dimensional general relativity,” Phys. Rev. **D70**, 124002 (2004). [arXiv:hep-th/0408141 [hep-th]].
- [13] B. Kleihaus, J. Kunz, E. Radu, M. J. Rodriguez, “New generalized nonspherical black hole solutions,” JHEP **1102**, 058 (2011). [arXiv:1010.2898 [gr-qc]].
- [14] H. Elvang, P. Figueras, “Black Saturn,” JHEP **0705**, 050 (2007). [hep-th/0701035].
- [15] F. Dowker, J. P. Gauntlett, G. W. Gibbons and G. T. Horowitz, “Nucleation of p-branes and fundamental strings,” Phys. Rev. D **53**, 7115 (1996) [arXiv:hep-th/9512154].
- [16] S. Tomizawa, Y. Yasui and A. Ishibashi, “Uniqueness theorem for charged dipole rings in five-dimensional minimal supergravity,” Phys. Rev. D **81**, 084037 (2010) [arXiv:0911.4309 [hep-th]].
- J. Armas and T. Harmark, “Uniqueness Theorem for Black Hole Space-Times with Multiple Disconnected Horizons,” JHEP **1005**, 093 (2010) [arXiv:0911.4654 [hep-th]].
- [17] H. Elvang and M. J. Rodriguez, “Bicycling Black Rings,” JHEP **0804**, 045 (2008) [arXiv:0712.2425 [hep-th]].
- K. Izumi, Prog. Theor. Phys. **119**, 757 (2008) [arXiv:0712.0902 [hep-th]].