

# Gradient representations and affine structures in $AE_n$

Axel Kleinschmidt and Hermann Nicolai

Max-Planck-Institut für Gravitationsphysik (Albert-Einstein-Institut), Mühlenberg 1,  
D-14476 Golm, Germany

E-mail: [axel.kleinschmidt@aei.mpg.de](mailto:axel.kleinschmidt@aei.mpg.de) and [hermann.nicolai@aei.mpg.de](mailto:hermann.nicolai@aei.mpg.de)

Received 29 June 2005, in final form 30 June 2005

Published 11 October 2005

Online at [stacks.iop.org/CQG/22/4457](http://stacks.iop.org/CQG/22/4457)

## Abstract

We study the indefinite Kac–Moody algebras  $AE_n$ , arising in the reduction of Einstein’s theory from  $(n + 1)$  spacetime dimensions to one (time) dimension, and their distinguished maximal regular subalgebras  $A_{n-1} \equiv \mathfrak{sl}_n$  and  $A_{n-2}^{(1)}$ . The interplay between these two subalgebras is used, for  $n = 3$ , to determine the commutation relations of the ‘gradient generators’ within  $AE_3$ . The low-level truncation of the geodesic  $\sigma$ -model over the coset space  $AE_n/K(AE_n)$  is shown to map to a suitably truncated version of the  $SL(n)/SO(n)$  nonlinear  $\sigma$ -model resulting from the reduction Einstein’s equations in  $(n + 1)$  dimensions to  $(1 + 1)$  dimensions. A further truncation to diagonal solutions can be exploited to define a one-to-one correspondence between such solutions, and null geodesic trajectories on the infinite-dimensional coset space  $\mathfrak{H}/K(\mathfrak{H})$ , where  $\mathfrak{H}$  is the (extended) Heisenberg group, and  $K(\mathfrak{H})$  its maximal compact subgroup. We clarify the relation between  $\mathfrak{H}$  and the corresponding subgroup of the Geroch group.

PACS numbers: 02.20.Tw, 11.30.Na, 04.20.Jb

## 1. Introduction

Infinite-dimensional symmetries in gravity were first discovered by Geroch [1] in the context of  $(3 + 1)$ -dimensional general relativity after dimensional reduction to  $(1 + 1)$  dimensions (see [2, 3] for an introduction and further references). The group structure was later shown to be associated with the affine extension of Ehlers’s  $SL(2, \mathbb{R})$  symmetry [4, 5], and similar affinizations of hidden symmetries were discovered for other (super-)gravity theories [4–6]. Evidence for the emergence of an even larger symmetry corresponding to the Kac–Moody theoretic over-extension  $AE_3$  of the original  $SL(2, \mathbb{R})$  was presented in [7], following earlier suggestions of [4, 8]. For gravity in  $(n + 1)$  spacetime dimensions, the conjectured symmetry is  $AE_n$ , while for 11-dimensional supergravity it is  $E_{10}$ , which contains  $AE_{10}$  as a subalgebra governing the gravitational sector of this theory.

In [9], this conjecture was re-examined using the insights from the study of cosmological billiards (reviewed in [10]). Besides a remarkable dynamical match of a certain ‘geodesic’ one-dimensional  $\sigma$ -model with the gravity theory, the conjecture was made that the Kac–Moody algebra<sup>1</sup> allows for a re-emergence of the dependence on the coordinates along which the theory had been reduced. This would entail a ‘dimensional transmutation’, in the sense that the evolution of the geometrical data of a higher-dimensional theory, usually governed by a set of partial differential equations, can be mapped to a *one-dimensional null-geodesic motion on some infinite-dimensional coset space*. This conjecture was based on the observation that the hyperbolic algebra  $E_{10}$  contains a set of generators possessing the correct structure for higher order gradients in the suppressed directions. These, and their analogues in  $AE_n$ , will be called ‘gradient generators’ in the present paper. Further progress in the study of the one-dimensional  $\sigma$ -model is partly (but not only) hindered by the lack of known commutation relations. Although the irreducible representations appearing in level expansions of these algebras w.r.t. to their  $\mathfrak{sl}_n$  subalgebras can be determined rather efficiently on the computer [11], the commutators are much harder to obtain<sup>2</sup>. Some progress on this front was reported recently in [12] where an algorithm for computing commutation relations in a Borel subalgebra was outlined. Different aspects of the one-dimensional model were studied in [13–15].

In this paper we study the  $\sigma$ -model based on  $AE_n$ , extending previous results of [10], in order to examine aspects of the general picture explained above. Our focus is on  $AE_n$  rather than  $E_{10}$ , since the core difficulties which one encounters in matching the one-dimensional  $\sigma$ -model and the higher-dimensional field equations appear right away at levels  $\pm 1$  in a graded decomposition of  $AE_n$  under its  $\mathfrak{sl}_n$  subalgebra. In other words, the key problem of elevating the linear duality of free spin-2 theories to the nonlinear level must be faced already in the first step, whereas for  $E_{10}$ , the difficulties only become visible at  $\ell = 3$  and beyond, because the duality relating the 3- and 6-form fields is still a *linear* one, modulo metric factors, just like in Maxwell theory.

As we will show, the ‘gradient representations’ are intimately linked to the affine subalgebra  $A_{n-2}^{(1)}$  of  $AE_n$ . Exploiting the interplay of this affine subalgebra with the finite-dimensional  $A_{n-1} \equiv \mathfrak{sl}_n$  subalgebra, we are able to derive an infinite new set of structure constants for  $AE_3$ . After the restriction of the  $AE_n$   $\sigma$ -model to the affine  $A_{n-2}^{(1)}$  subsector, we exhibit a map between special solutions of the  $\sigma$ -model and solutions of the gravitational field equations with  $(n - 1)$  commuting spacelike Killing vectors, corresponding to a reduction from  $(n + 1)$  dimensions to  $(1 + 1)$  dimensions, but with a restricted space dependence. These results are analogous to previous ones in [9, 10, 13–15], but permit us to expose the remaining discrepancies in the simplest possible context. In order to focus on these difficulties, we truncate the affine  $\sigma$ -model further to a ‘Heisenberg coset model’  $\mathfrak{H}/K(\mathfrak{H})$ , which can be solved exactly, and whose gravitational counterpart corresponds to diagonal metric solutions with two commuting (spacelike) Killing vectors (known as ‘polarized Gowdy cosmologies’ [16, 17]). We will then exhibit an explicit one-to-one correspondence between these two models, by showing how the general initial data of the Heisenberg  $\sigma$ -model and the null geodesic trajectories on  $\mathfrak{H}/K(\mathfrak{H})$  which they generate can be mapped to a general space- and time-dependent diagonal metric configuration satisfying Einstein’s equation. In this way, we are able to validate the ‘gradient hypothesis’, and thereby the main conjecture of [9], at least in this simplified context.

We also elucidate the relation between the standard action of the (restricted) Geroch group on diagonal solutions, and the action of the Heisenberg group  $\mathfrak{H}$  on the null geodesics.

<sup>1</sup> We often use the acronyms ‘KMA’ for ‘Kac–Moody algebra’ and ‘CSA’ for ‘Cartan subalgebra’.

<sup>2</sup> See also remarks after (2.18) to appreciate the challenge.

More specifically, we will show that the action of these two groups coincides on the domain where the Geroch group acts non-trivially. Let us recall that only ‘half’ of the Geroch group acts non-trivially in the standard realization [2, 5], whereas the other half merely shifts the integration constants arising in the definition of the higher order ‘dual potentials’, and has no effect on the physical metric. For this reason, the standard Geroch group affects only part of the initial data (the full initial data of the off-diagonal degrees of freedom, but only the ‘initial coordinates’ of the scale factors). By contrast, the realization proposed here is such that the Heisenberg subgroup of the Geroch group acts non-trivially on *all* initial data.

Other attempts to generate spacetime dependence through algebraic constructions based on  $E_{11}$  [18] and similar ‘very-extended’ algebras [19] have been developed and discussed in [20–22]. In [20], this was achieved by including a certain irreducible representation of the relevant very-extended algebra containing translation generators and (infinitely many) other generalized central charges, whereas in [21] spacetime is thought to occur through an auxiliary parameter and gradient representation in restricted models. A key difference between the present approach and [18] is that the correspondence exhibited here works *only after the gauge degrees of freedom on both sides have been eliminated by suitable gauge choices*: on the  $\sigma$ -model side this implies the complete elimination of the degrees of freedom associated with the maximal compact subgroups, whereas on the gravitational side, it involves not only gauge fixing the vielbein (as discussed in section 5.1), but also solving the canonical constraints. As a consequence, the global symmetry relates solutions which are *physically distinct*. By contrast, the proposal of [18, 20] seeks a ‘covariant formulation’. This means that the symmetry is actually much larger than the relevant very-extended Kac–Moody group, as it must contain general coordinate transformations (for instance, via the closure with the conformal group [23]) and other gauge transformations; hence, one must disentangle transformations relating gauge equivalent configurations from those which generate physically inequivalent solutions. While it is not possible to really discriminate between the different proposals by analysing low-level degrees of freedom, the issue will be decided by whether and how the higher level fields can be fitted into the scheme. The present paper is a first step in this direction.


This paper is structured as follows. First we define and analyse the KMA  $AE_n$  and its two distinguished maximal regular subalgebras  $A_{n-2}^{(1)}$  and  $A_{n-1}$  and define the gradient representations in this context in section 2. In section 3, we compute infinitely many new structure constants involving the gradient generators in the special case of  $AE_3$  by combining these two subalgebras. The  $\sigma$ -model based on  $AE_n$  is defined in section 4 where we study its restriction to  $A_{n-2}^{(1)}$ . (The restriction to  $A_{n-1}$  is studied in appendix B where we also study gravity coupled to  $p$ -forms.) The map from (parts of) the affine model to the two-dimensional reduction of gravity is deduced in section 5. Finally, by using the Heisenberg model we examine the relation of the  $\sigma$ -model to the Geroch group in section 6. Appendix A contains a proof of the fact that the compact subalgebra of a (split) KMA is *not* a KMA.

## 2. Distinguished maximal regular subalgebras of $AE_n$

### 2.1. Definition of $AE_n$

The indefinite Kac–Moody algebras  $AE_n$ <sup>3</sup> (for  $n \geq 3$ ) are defined in the usual way via the Chevalley–Serre presentation [24] associated with the Dynkin diagram displayed in figure 1. The simple positive and negative generators are denoted by  $e_a$  and  $f_a$  respectively, and the

<sup>3</sup> We will designate by  $AE_n$  both the group and the Lie algebra, as it should be clear from the context which is meant.



**Figure 1.** The Dynkin diagram of  $AE_n$  with labelling of the simple roots shown. Solid nodes (black and grey) belong to  $A_{n-1} \equiv \mathfrak{sl}_n$  which gets enhanced to  $\mathfrak{gl}_n$ . Grey nodes mark the common ‘horizontal’  $A_{n-2} \equiv \mathfrak{sl}_{n-1}$  subalgebra.

Cartan subalgebra (CSA) generators by  $h_a$ , where  $a = 1, 2, \dots, n$ . We consider the algebra over  $\mathbb{R}$  in split real form.  $AE_n$  is hyperbolic for  $n \leq 9$ ; this means that all Dynkin subdiagrams obtained by removing one or more nodes are either affine or finite. We will also need to make use of the Chevalley involution  $\omega$ , defined on the simple generators by

$$\omega(e_a) = -f_a, \quad \omega(f_a) = -e_a, \quad \omega(h_a) = -h_a, \tag{2.1}$$

and the generalized transposition

$$x^T := -\omega(x). \tag{2.2}$$

The ‘maximal compact’ subalgebra  $K(AE_n)$  of  $AE_n$  is then defined as the invariant algebra w.r.t. the involution  $\omega$ , namely

$$K(AE_n) := \{x \in AE_n | x = \omega(x)\} \tag{2.3}$$

and consists of the ‘antisymmetric’ Lie algebra elements in view of (2.2).

In this section we shall consider two distinguished maximal regular subalgebras of  $AE_n$ , namely  $\mathfrak{sl}_n$  and the affine  $A_{n-2}^{(1)}$ , obtained by removing the nodes labelled  $n$  and  $1$ , respectively. The first generates the group of special linear transformations acting on the spatial  $n$ -bein of gravity in  $(n + 1)$  spacetime dimensions, and can be enlarged to  $\mathfrak{gl}_n$  by inclusion of the CSA generator associated with the white node in figure 1. The second subalgebra corresponds to the generalization of the Geroch group that is obtained after the reduction of pure gravity from  $(n + 1)$  dimensions to  $(1 + 1)$  dimensions.

2.2.  $A_{n-1} \equiv \mathfrak{sl}_n$  subalgebra and level decomposition of  $AE_n$

The regular subalgebra of type  $A_{n-1} \equiv \mathfrak{sl}_n$  is generated by considering only commutators of the simple generators associated with nodes 1 up to  $(n - 1)$  in figure 1. By including the Cartan subalgebra element  $h_n$  one can extend  $\mathfrak{sl}_n$  to  $\mathfrak{gl}_n$ . Its generators are denoted by  $K^a_b$  ( $a, b = 1, 2, \dots, n$ ) and obey the standard commutation relations

$$[K^a_b, K^c_d] = \delta^c_b K^a_d - \delta^a_d K^c_b. \tag{2.4}$$

For these generators, the transposition (2.2) reduces to  $(K^T)^a_b = K^b_a$ . Equation (2.4) entails the following identification of the  $\mathfrak{gl}_n$  elements with the Chevalley–Serre generators of  $AE_n$  for  $i = 1, 2, \dots, n - 1$

$$e_i = K^i_{i+1}, \quad f_i = K^{i+1}_i, \quad h_i = K^i_i - K^{i+1}_{i+1}. \tag{2.5}$$

Regularity of the  $\mathfrak{gl}_n$  subalgebra means that the standard invariant bilinear form on  $AE_n$ , which is defined by

$$\langle e_i | f_j \rangle = \delta_{ij}; \quad \langle h_i | h_j \rangle = A_{ij} \tag{2.6}$$

coincides with the usual Cartan–Killing form on  $\mathfrak{gl}_n$  when restricted to the latter subalgebra ( $A_{ij}$  is the Cartan matrix of  $AE_n$ , which is simply laced). Explicitly, the scalar product on  $\mathfrak{gl}_n$  is given by

$$\langle K_b^a | K_d^c \rangle = \delta_b^c \delta_d^a - \delta_b^a \delta_d^c. \tag{2.7}$$

It is straightforward to check that this is indeed consistent with (2.6). Let us also express the trace  $K = \sum_{a=1}^n K_a^a$  in  $\mathfrak{gl}_n$  in terms of the CSA generators  $h_a$ ; we have

$$K = -(n - 1)h_1 - (2n - 2)h_2 - (2n - 3)h_3 - \dots - nh_n. \tag{2.8}$$

Solving for  $h_n$  gives

$$h_n = -K^1_1 - K^2_2 + K^n_n = -K + K^3_3 + K^4_4 + \dots + K^{n-1}_{n-1} + 2K^n_n. \tag{2.9}$$

The adjoint representation of  $AE_n$  can be decomposed under the (adjoint) action of  $\mathfrak{gl}_n$  into an infinite tower of  $\mathfrak{sl}_n$  representations. The  $\mathfrak{sl}_n$  level  $\ell$ , or simply the *level*, of a given representation counts the number of occurrences of the simple root  $\alpha_n$  in the corresponding  $AE_n$  root  $\alpha$ , i.e.

$$\alpha = \ell \alpha_n + \sum_{j=1}^{n-1} m^j \alpha_j. \tag{2.10}$$

This level is left invariant by the action of  $\mathfrak{sl}_n$  and provides an elliptic slicing of the forward lightcone in the  $AE_n$  root lattice. Level  $\ell = 0$  contains the adjoint of  $\mathfrak{gl}_n$ . At level  $\ell = 1$  we have the representation

$$[0, 1, 0, 0, \dots, 0, 1] \longleftrightarrow E^{a_1 \dots a_{n-2}, a_{n-1}} \tag{2.11}$$

associated with the Young tableau

$a_1$	$a_{n-1}$
$a_2$	
$\vdots$	
$a_{n-2}$	

This tensor is therefore antisymmetric in its first  $(n - 2)$  indices  $[a_1 \dots a_{n-2}]$ , and obeys

$$E^{[a_1 \dots a_{n-2}, a_{n-1}]} = 0. \tag{2.12}$$

Under Chevalley transposition we obtain

$$F_{a_1 \dots a_{n-2}, a_{n-1}} \equiv -\omega(E^{a_1 \dots a_{n-2}, a_{n-1}}). \tag{2.13}$$

Under  $\mathfrak{gl}_n$  the tensors  $E^{a_1 \dots a_{n-2}, a_{n-1}}$  and  $F_{a_1 \dots a_{n-2}, a_{n-1}}$  transform contravariantly or covariantly, as indicated by the position of indices, for instance

$$[K_b^a, E^{c_1 \dots c_{n-2}, c_{n-1}}] = \delta_b^{c_1} E^{ac_2 \dots c_{n-2}, c_{n-1}} + \dots + \delta_b^{c_{n-1}} E^{c_1 \dots c_{n-2}, a}. \tag{2.14}$$

Hence the level is counted by the operator  $\frac{1}{n-1} K$ .

The identification of the  $AE_n$  Chevalley–Serre basis is completed by relating the generators  $e_n$  and  $f_n$  to the level  $\ell = \pm 1$  generators via

$$e_n = E^{3^4 \dots n, n}, \quad f_n = F_{3^4 \dots n, n}. \tag{2.15}$$

The commutator  $[(\ell = 1), (\ell = -1)]$  is best written using an auxiliary (dummy) tensor  $X_{a_1, \dots, a_{n-2}, a_{n-1}}$  (with the same Young symmetries as  $F$ ) in the form

$$\begin{aligned} [X_{a_1, \dots, a_{n-2}, a_{n-1}} E^{a_1 \dots a_{n-2}, a_{n-1}}, F_{b_1, \dots, b_{n-2}, b_{n-1}}] = & -(n-2)! (X_{b_1, \dots, b_{n-2}, b_{n-1}} K \\ & - X_{b_1, \dots, b_{n-2}, e} K^e_{b_{n-1}} - (n-2) K^e_{[b_{n-2}} X_{b_1, \dots, b_{n-3}]e, b_{n-1}}). \end{aligned} \tag{2.16}$$

This is consistent with the normalization<sup>4</sup>

$$\langle E^{a_1 \dots a_{n-2}, n-1} | F_{b_1, \dots, b_{n-2}, b_{n-1}} \rangle = -\frac{n-2}{n-1} (n-2)! (\delta_{b_1 \dots b_{n-2}}^{a_1 \dots a_{n-2}} \delta_{b_{n-1}}^{a_{n-1}} + \delta_{b_{n-1}}^{[a_1} \delta_{[b_2 \dots b_{n-2}}^{a_2 \dots a_{n-2}}] \delta_{b_1]}^{a_{n-1}}), \tag{2.17}$$

corresponding to the standard normalization for the Chevalley generators

$$\langle e_n | f_n \rangle = 1. \tag{2.18}$$

All higher levels in the  $\mathfrak{sl}_n$  decomposition can be obtained by taking multiple commutators of the level  $\ell = 1$  generator  $E^{a_1 \dots a_{n-2}, a_{n-1}}$ . For  $AE_3$ , the decomposition into irreducible representations of  $\mathfrak{sl}_3$  is known up to level  $\ell = 56$  [11]. Counting outer multiplicities, the total number of  $\mathfrak{sl}_3$  representations up to that level is [25]

$$\#(\text{representations for } \ell \leq 56) = 20\,994\,472\,770\,550\,672\,476\,591\,949\,725\,720. \tag{2.19}$$

Consequently, the complete table of structure constants up to that level will already contain more than  $10^{62}$  entries! Hence, the ‘gradient representations’ that we will consider below constitute only a tiny subsector (but not a subalgebra) of the full Lie algebra.

### 2.3. Affine $A_{n-2}^{(1)}$ subalgebra

Figure 1 also shows that  $AE_n$  has a regular affine subalgebra  $A_{n-2}^{(1)} \equiv \widehat{\mathfrak{sl}_{n-1}} \oplus \mathbb{R}\hat{c} \oplus \mathbb{R}\hat{d}$  generated by nodes 2 up to  $n$  (the circular subdiagram in figure 1)<sup>5</sup>. We write the corresponding traceless generators as  $\bar{K}_m^\alpha$  with  $m \in \mathbb{Z}$ , and Greek indices  $\alpha, \beta, \dots \in \{2, \dots, n\}$ . The generators which belong to both this affine subalgebra and the  $\mathfrak{sl}_{n-1}$  constitute what we call the horizontal algebra  $\mathfrak{sl}_{n-1}$  corresponding to the nodes 2 up to  $n-1$ , with generators

$$\bar{K}_{0\beta}^\alpha = K^\alpha_\beta - \frac{1}{n-1} \delta^\alpha_\beta K^\gamma_\gamma \tag{2.20}$$

in the  $\mathfrak{gl}_n$  from above. The relevant nodes in figure 1 are marked in grey.

The affine commutation relations are

$$[\bar{K}_{m\beta}^\alpha, \bar{K}_{n\delta}^\gamma] = \delta^\gamma_\beta \bar{K}_{m+n\delta}^\alpha - \delta^\alpha_\delta \bar{K}_{m+n\beta}^\gamma + m\delta_{m,-n} \left( \delta^\alpha_\delta \delta^\gamma_\beta - \frac{1}{n-1} \delta^\alpha_\beta \delta^\gamma_\delta \right) \hat{c}. \tag{2.21}$$

The central element  $\hat{c}$  and the derivation  $\hat{d}$  required for the  $n$ -dimensional CSA of  $A_{n-2}^{(1)}$  can be identified in the CSA of the rank  $n$  KMA  $AE_n$ . Requiring the affine level counting operator  $\hat{d}$  to obey

$$[\hat{d}, e_n] = -e_n, \quad [\hat{d}, e_i] = 0 \quad (i = 2, \dots, n-1), \quad \langle \hat{d} | \hat{d} \rangle = 0, \tag{2.22}$$

<sup>4</sup> Symmetrization and antisymmetrization are defined with ‘strength one’ throughout this paper, such as for instance in

$$\delta_{b_1 b_2}^{a_1 a_2} := \frac{1}{2} \delta_{b_1}^{a_1} \delta_{b_2}^{a_2} - \frac{1}{2} \delta_{b_2}^{a_1} \delta_{b_1}^{a_2}, \quad \bar{\delta}_{b_1 b_2}^{a_1 a_2} := \frac{1}{2} \delta_{b_1}^{a_1} \delta_{b_2}^{a_2} + \frac{1}{2} \delta_{b_2}^{a_1} \delta_{b_1}^{a_2}.$$

where a bar over  $\delta$  denotes symmetrization, and no bar means antisymmetrization.

<sup>5</sup> We follow the convention that  $\widehat{\mathfrak{sl}_{n-1}}$  denotes the loop algebra based on  $A_{n-2} \equiv \mathfrak{sl}_{n-1}$ . The non-twisted affine algebra  $A_{n-2}^{(1)}$  is obtained by adjoining the central element  $\hat{c}$  and the derivation  $\hat{d}$  to the Cartan subalgebra, which then has dimension  $n$  [24]. (The rank of  $A_{n-2}^{(1)}$  is  $n-1$ .)

results in the following expressions for  $\hat{c}$  and  $\hat{d}$  in terms of the  $\mathfrak{gl}_n$  elements:

$$\hat{c} = \sum_{i=2}^n h_i = -K^1_1, \tag{2.23}$$

$$\hat{d} = \frac{n-2}{2(n-1)}K^1_1 - \frac{1}{n-1}K^\alpha_\alpha, \tag{2.24}$$

such that

$$\begin{aligned} K = K^a_a = K^1_1 + K^\alpha_\alpha &= -(n-1)\hat{d} - \frac{n}{2}\hat{c}, \\ \implies K^\alpha_\alpha &= -(n-1)\hat{d} - \frac{n-2}{2}\hat{c}. \end{aligned} \tag{2.25}$$

The CSA elements  $\hat{c}$  and  $\hat{d}$  are normalized according to

$$\langle \hat{c} | \hat{d} \rangle = -1, \quad \langle \hat{d} | \hat{d} \rangle = \langle \hat{c} | \hat{c} \rangle = 0, \tag{2.26}$$

and the other non-vanishing inner product is

$$\langle \bar{K}^\alpha_{m\beta} | \bar{K}^\gamma_{n\delta} \rangle = \delta_{m,-n} \left( \delta^\alpha_\delta \delta^\gamma_\beta - \frac{1}{n-1} \delta^\alpha_\beta \delta^\gamma_\delta \right), \tag{2.27}$$

which is compatible with (2.6).

Alternatively, the adjoint of  $AE_n$  can be decomposed under the action of  $A_{n-2}^{(1)}$ . In this case, the grading is labelled by the *affine level*, which is equal to the coefficient of the root  $\alpha_1$  (now providing a parabolic slicing of the forward lightcone in the root lattice). This decomposition was introduced and studied in [26] for  $AE_3$ , but we will not require these results here<sup>6</sup>.

#### 2.4. Horizontal $\mathfrak{sl}_{n-1}$ and gradient representations

Writing the Kac–Moody algebra  $AE_n$  as a graded representation of its  $\mathfrak{sl}_n$  subalgebra is not independent of making use of the affine  $A_{n-2}^{(1)}$  subalgebra. This is due to the identification of common generators in the horizontal algebra  $\mathfrak{sl}_{n-1} \equiv A_{n-2} = A_{n-1} \cap A_{n-2}^{(1)}$ . We repeat that this is the  $A_{n-2}$  subalgebra of  $AE_n$  generated by the  $n-2$  (grey) nodes 2 up to  $n-1$  in diagram 1, with the generators given in equation (2.20). In particular, the affine generator  $\bar{K}^\alpha_{m\beta}$  on affine level  $m$ , transforming in the adjoint of the horizontal  $\mathfrak{sl}_{n-1}$ , also transforms under  $\mathfrak{sl}_n \supset \mathfrak{sl}_{n-1}$ , and it is natural to ask to which  $\mathfrak{sl}_n$  representation it belongs. This is most easily determined by considering the lowest weight vector and its Dynkin labels with respect to  $\mathfrak{sl}_n$ . The lowest weight vector lies in the root space of the level- $m$   $AE_n$  root (assuming  $m \geq 1$  from now on)

$$\alpha = m\alpha_n + (m-1) \sum_{j=2}^{n-1} \alpha_j \tag{2.28}$$

corresponding to  $\mathfrak{sl}_n$  Dynkin labels  $[m-1, 1, 0, \dots, 0, 1]$ . We see that this is consistent with the common horizontal  $\mathfrak{sl}_{n-1}$ : in terms of the Dynkin labels restricting to the horizontal  $A_{n-2}$  of  $A_{n-2}^{(1)}$  means dropping the first entry of  $[m-1, 1, 0, \dots, 0, 1]$  and the resulting representation is indeed simply the adjoint  $[1, 0, \dots, 0, 1]$  of the horizontal  $\mathfrak{sl}_{n-1}$  as anticipated. The generator

<sup>6</sup> Beyond the so-called basic representation on level one there are already infinitely many  $A_{n-2}^{(1)}$  representations on affine level two, which can be calculated from modularity and an appropriate implementation of the Serre relations [26]. This is true not only for  $AE_3$  but also for  $AE_n$ .

corresponding to the  $[m - 1, 1, 0, \dots, 0, 1]$  representation of  $\mathfrak{sl}_n$  is therefore associated with the following Young tableau:

	...		$c_1$	$c_{n-1}$
	...		$c_2$	
⋮		⋮	⋮	
	...		$c_{n-2}$	
	...			

This level- $m$  element of  $AE_n$  thus has  $(m - 1)$  symmetric sets of  $(n - 1)$  antisymmetric indices and then  $(n - 1)$  indices with the structure of the representation on  $\mathfrak{sl}_n$  level  $\ell = 1$ , cf (2.11)–(2.12). Because this is rather cumbersome to write, it is convenient to treat this tensor as a representation of  $\mathfrak{sl}_n$ , and to dualize the sets of antisymmetric indices by means of the  $\mathfrak{sl}_n$   $\varepsilon$ -symbol. The result is an  $\mathfrak{sl}_n$  tensor with  $(m - 1)$  lower indices

$$E_{b_1 \dots b_{m-1}}^{c_1 \dots c_{n-2}, c_{n-1}} \tag{2.29}$$

which is *symmetric* in these indices, while the upper indices belong to the level-one representation (2.11). There is a similar expression for the transposed  $F$  generator. We stress that (2.4) and (2.29) are equivalent *only* as representations of  $\mathfrak{sl}_n$ , but not  $\mathfrak{gl}_n$ , as they carry different  $\mathfrak{gl}_1$  charges. The commutation relations must therefore be amended by a correction term for the transformation under the trace  $K$  in order to account for this charge. We will come back to this point below for  $AE_3$ .

As noted in [9], the infinite tower of representations (2.29) has precisely the right structure corresponding to the *spatial gradients* of the low-level fields, such that each index  $b_i$  becomes associated with a gradient operator  $\partial/\partial x^{b_i}$ , and the irreducibility condition of the above Young tableau translates to vanishing divergence. More specifically, for  $AE_n$ , the relevant degree of freedom is the *dual graviton*, corresponding to the above tableau with only the right two columns as in (2.11), and appears already at level  $\ell = 1$  (for  $E_{10}$  it appears only at level  $\ell = 3$ ).<sup>7</sup> For this reason, we will refer to these representations as *gradient representations*. Let us emphasize that, except for the lowest levels, this is so far only a kinematical correspondence<sup>8</sup>.


The other main feature of the gradient representations is that they contain the affine generators. More specifically, the latter are obtained by performing a ‘dimensional reduction’, restricting the gradient indices (i.e. the lower indices in (2.29)) to the single value  $b_1 = \dots = b_{m-1} = 1$ , and the upper indices to the remaining values  $\alpha, \beta, \dots = 2, \dots, n$ . This truncation corresponds precisely to a dimensional reduction of Einstein’s theory from  $(n + 1)$  to  $(1 + 1)$  spacetime dimensions, with spacelike Killing vectors  $\partial/\partial x^2, \dots, \partial/\partial x^n$ , and  $x \equiv x^1$  as the surviving spatial coordinate. The precise identification of  $\bar{K}_{m\beta}^\alpha$  (for  $m > 0$ ) as part of the  $\mathfrak{sl}_n$  tensor (2.29) is

$$\bar{K}_{m\beta}^\alpha = \frac{1}{(n - 2)!} \varepsilon_{\beta\gamma_1 \dots \gamma_{n-2}} E_{\underbrace{1 \dots 1}_{(m-1)}}^{\gamma_1 \dots \gamma_{n-2}, \alpha}, \tag{2.30}$$

<sup>7</sup> The dual graviton representation has been discussed previously in [9, 27–33].

<sup>8</sup> We note that there are generators resembling a  $k$ th spatial derivative (on level  $\ell + k$ ) of any given generator (on level  $\ell$ ). These will generally occur with exponentially increasing outer multiplicity as they belong to a standard lowest weight representation of the affine algebra. The gradient representations (2.29) are distinguished by being generated from the adjoint representation of the affine algebra and hence have constant outer multiplicity, equal to 1.





**Figure 2.** The Dynkin diagram of  $AE_3$  with labelling of the simple roots. Solid nodes belong to  $\mathfrak{sl}_3$  which gets enhanced to  $\mathfrak{gl}_3$ . The grey node marks the horizontal  $A_1 \equiv \mathfrak{sl}_2$ .

where the indices  $\alpha, \beta, \dots = 2, \dots, n$  have been restricted to  $\mathfrak{sl}_{n-1}$ . For affine level  $m < 0$  the identification is

$$\bar{K}_{m\beta}^\alpha = \frac{1}{(n-2)!} \varepsilon^{\alpha\gamma_1 \dots \gamma_{n-2}} F^{\overbrace{1 \dots 1}^{(m-1)}_{\gamma_1 \dots \gamma_{n-2}, \beta}} \tag{2.31}$$

The correctness of these identifications is easy to check at levels  $|m| \leq 1$ .

One can now exploit the interplay between  $\mathfrak{sl}_n$  and  $A_{n-2}^{(1)}$  to determine the complete commutators of the ‘gradient generators’ modulo non-gradient representations. This is achieved by writing the general ansatz for a commutator in  $\mathfrak{sl}_n$  covariant form, and then restricting to the horizontal  $\mathfrak{sl}_{n-1}$  and using the affine relations (2.21). In the following section, we exemplify this procedure for  $AE_3$ .

### 3. $AE_3$ and its gradient commutators via $\mathfrak{sl}_3$ and $A_1^{(1)}$

#### 3.1. $AE_3$ in $\mathfrak{sl}_3$ form

The Dynkin diagram of  $AE_3$  is displayed in figure 2. Before computing the contribution of the gradient generators to commutators for  $AE_3$  we give the explicit formulæ for the  $\mathfrak{sl}_3$  generators and the appropriate low  $\mathfrak{sl}_3$  level commutators of  $AE_3$ .

In this case, the general identifications (2.8) and (2.9) read

$$h_3 = -K + 2K^3, \quad K = -2h_1 - 4h_2 - 3h_3. \tag{3.1}$$

The  $\mathfrak{gl}_3$  commutation relations and inner products are identical in form to (2.4) but with index ranges now restricted to  $a = 1, 2, 3$ . The decomposition of  $AE_3$  under its  $\mathfrak{sl}_3$  subalgebra results in table 1, containing levels  $\ell = 1, \dots, 5$ .

Generators belonging to the class of gradient generators have been marked with an asterisk. Our notation for representations of  $\mathfrak{gl}_3$  is the one inherited from  $\mathfrak{sl}_3$ . The general  $\ell = 1$  generator (2.11) reduces to a generator of the type

$$E^{b_1 b_2}, \tag{3.2}$$

where we have dropped the separating comma. The Young irreducibility constraint (2.12) now implies that this tensor is symmetric,

$$E^{b_1 b_2} = E^{b_2 b_1}. \tag{3.3}$$

For  $\mathfrak{sl}_3$  one can write any representation with Dynkin labels  $[p, q]$  as a tensor with  $p$  lower and  $q$  upper indices, after lowering the  $p$  pairs of antisymmetric indices with  $\varepsilon$  of  $\mathfrak{sl}_3$ . The irreducibility constraints are then symmetric in lower and upper indices separately and vanishing of any contraction. The totally antisymmetric tensor  $\varepsilon_{abc}$  used for lowering the indices, however, is *not* invariant under  $\mathfrak{gl}_3$  as can be checked easily. This means that while the  $\ell = 2$  tensor with natural indices  $E^{c_1 c_2, b_1 b_2}$  transforms under  $\mathfrak{gl}_3$  as the indices suggest, i.e.

$$[K^a_b, E^{c_1 c_2, b_1 b_2}] = \delta_b^{c_1} E^{a c_2, b_1 b_2} + \delta_b^{c_2} E^{c_1 a, b_1 b_2} + \delta_b^{b_1} E^{c_1 c_2, a b_2} + \delta_b^{b_2} E^{c_1 c_2, b_1 a}, \tag{3.4}$$

the corresponding tensor with lowered indices does *not* transform in the obvious way. Rather one has to add a compensating term for the transformation under the trace  $K = K^a_a$ . Generally,

$$[K^c_d, E_{a_1 \dots a_k}^{b_1 b_2}] = -k \delta_{(a_1}^c E_{a_2 \dots a_k) d}^{b_1 b_2} + 2 \delta_d^{(b_1} E_{a_1 \dots a_k}^{b_2) c} + k \delta_d^c E_{a_1 \dots a_k}^{b_1 b_2}. \tag{3.5}$$

**Table 1.** Adjoint of  $AE_3$  decomposed under regular  $\mathfrak{sl}_3$  subalgebra. The charge of an element  $X$  on level  $\ell$  under the trace  $K = K^a_a$  of  $\mathfrak{gl}_3$  is  $[K, X] = 2\ell X$  as explained in the text.

$\ell$	$[p_1, p_2]$	$\alpha$	$\alpha^2$	mult $\alpha$	$\mu$	Gradient
1	[0, 2]	(0, 0, 1)	2	1	1	*
2	[1, 2]	(0, 1, 2)	2	1	1	*
3	[2, 2]	(0, 2, 3)	2	1	1	*
	[1, 1]	(1, 3, 3)	-4	3	1	
4	[3, 2]	(0, 3, 4)	2	1	1	*
	[2, 1]	(1, 4, 4)	-6	5	2	
	[1, 0]	(2, 5, 4)	-10	11	1	
	[0, 2]	(2, 4, 4)	-8	7	1	
	[1, 3]	(1, 3, 4)	-2	2	1	
5	[4, 2]	(0, 4, 5)	2	1	1	*
	[3, 1]	(1, 5, 5)	-8	7	3	
	[2, 0]	(2, 6, 5)	-14	22	3	
	[0, 1]	(3, 6, 5)	-16	30	2	
	[0, 4]	(2, 4, 5)	-6	5	2	
	[1, 2]	(2, 5, 5)	-12	15	4	
	[2, 3]	(1, 4, 5)	-4	3	2	

The last term is the necessary correction: the charge of the original tensor on level  $k + 1$  with  $2k + 2$  upper indices under the trace  $K$  is  $2k + 2$ , and this charge has to be maintained. Similar remarks apply to the Chevalley transposed generators

$$F^{a_1 \dots a_p}_{b_1 \dots b_q} = (E_{a_1 \dots a_p}{}^{b_1 \dots b_q})^T. \tag{3.6}$$

We now present the commutation of the  $AE_3$  generators for levels  $|\ell| \leq 2$  in  $\mathfrak{sl}_3$  form. These can be deduced from table 1 after the generators have been normalized in the standard bilinear form<sup>9</sup>. For  $\ell = 1, 2$  we demand

$$\langle E^{b_1 b_2} | F_{d_1 d_2} \rangle = \bar{\delta}_{d_1 d_2}^{b_1 b_2} := \delta_{d_1}^{(b_1} \delta_{d_2}^{b_2)}, \tag{3.7}$$

$$\langle E_a{}^{b_1 b_2} | F^c{}_{d_1 d_2} \rangle = P_a{}^{b_1 b_2}{}^c{}_{d_1 d_2} := \delta_a^c \bar{\delta}_{d_1 d_2}^{b_1 b_2} - \frac{1}{2} \delta_a^{(b_1} \delta_{(d_1}^{b_2)} \delta_{d_2)}^c. \tag{3.8}$$

The commutation relations are then

$$[E^{b_1 b_2}, F_{d_1 d_2}] = -\bar{\delta}_{d_1 d_2}^{b_1 b_2} K + 2\delta_{(d_1}^{(b_1} K_{d_2)}^{b_2)}, \tag{3.9}$$

$$[E^{b_1 b_2}, E^{c_1 c_2}] = 2\epsilon^{b_1 c_1 a} E_a{}^{b_2 c_2}, \tag{3.10}$$

$$[F_{d_1 d_2}, F_{c_1 c_2}] = -2\epsilon_{d_1 c_1 a} F^a{}_{d_2 c_2}, \tag{3.11}$$

$$[E_a{}^{b_1 b_2}, F_{d_1 d_2}] = 2\epsilon_{ae(d_1} \delta_{d_2)}^{(b_1} E^{b_2)e}, \tag{3.12}$$

$$[F^c{}_{d_1 d_2}, E^{b_1 b_2}] = 2\epsilon^{ce(b_1} \delta_{(d_1}^{b_2)} F_{d_2)e}, \tag{3.13}$$

$$[E_a{}^{b_1 b_2}, F^c{}_{d_1 d_2}] = -P_a{}^{b_1 b_2}{}^c{}_{d_1 d_2} K + 2\delta_a^c \delta_{(d_2}^{(b_1} K_{d_2)}^{b_2)} - \bar{\delta}_{d_1 d_2}^{b_1 b_2} K_a^c - \frac{1}{2} \delta_a^{(b_1} K_{(d_1}^{b_2)}{}_{d_2)}^c. \tag{3.14}$$

Here, symmetrization over lower or upper non-contracted indices is implicit.

<sup>9</sup> In general, there remain signs  $\pm 1$  which need to be fixed consistently. Up to  $|\ell| \leq 2$  our choice is consistent.

### 3.2. $A_1^{(1)}$ and gradient commutators

The identification of the affine  $A_1^{(1)} \subset AE_3$  subalgebra is achieved through (cf (2.23))

$$\begin{aligned} \hat{c} = h_2 + h_3 = -K^1_1, \quad \hat{d} = h_1 + \frac{5}{4}h_2 + \frac{3}{4}h_3 = \frac{1}{4}K^1_1 - \frac{1}{2}(K^2_2 + K^3_3) \\ K = K^a_a = K^1_1 + K^\gamma_\gamma = -2\hat{d} - \frac{3}{2}\hat{c}, \quad \implies K^\gamma_\gamma = -2\hat{d} - \frac{1}{2}\hat{c}. \end{aligned} \tag{3.15}$$

The conventional choice for  $AE_3$  would be to take node 3 to be the horizontal node instead of 2 as we did, since this is more natural from the extension point of view. This choice would correspond to the Ehlers  $SL(2, \mathbb{R})$ , and it is also the choice taken by [26]. However, it does not reduce to the desired horizontal  $\mathfrak{sl}_2 \subset \mathfrak{sl}_3$  subalgebra, and for this reason we here make another choice, corresponding to the ‘Matzner–Misner’  $SL(2, \mathbb{R})$  in the physical interpretation. The gradient generators (2.29) on level  $\ell$  have Dynkin labels  $[\ell - 1, 2]$  and after lowering the sets of antisymmetric indices take the form

$$E_{a_1 \dots a_{\ell-1}}{}^{b_1 b_2}. \tag{3.16}$$

We now determine the contribution of the gradient generators to  $AE_3$  commutators by writing for each commutator the most general  $\mathfrak{sl}_3$  covariant ansatz compatible with the level decomposition, and then matching this ansatz with the affine commutation relations (2.21). Let us first note the projector onto a representation with Dynkin labels  $[\ell, 2]$ :

$$\begin{aligned} P_{a_1 \dots a_\ell}{}^{b_1 b_2} |^{c_1 \dots c_\ell}{}_{d_1 d_2} = \bar{\delta}_{a_1 \dots a_\ell}{}^{b_1 b_2} \delta_{d_1 d_2} - \frac{2\ell}{3 + \ell} \bar{\delta}_{a_1 \dots a_{\ell-1}}{}^{c_1 \dots c_{\ell-1}} \delta_{a_\ell}{}^{(b_1} \delta_{d_1}^{b_2)} \delta_{d_2}{}^{c_\ell} \\ + \frac{\ell(\ell - 1)}{(3 + \ell)(2 + \ell)} \bar{\delta}_{a_1 \dots a_{\ell-2}}{}^{c_1 \dots c_{\ell-2}} \bar{\delta}_{a_{\ell-1} a_\ell}{}^{b_1 b_2} \bar{\delta}_{d_1 d_2}{}^{c_{\ell-1} c_\ell}, \end{aligned} \tag{3.17}$$

where the right-hand side also has to be symmetrized over the  $a$  and  $c$  indices. Here,  $\bar{\delta}$  denotes the strength one symmetrizing projector. The second and third terms are only present when  $\ell \geq 1$  or  $\ell \geq 2$ , respectively. The projector has the unusual property that it also serves as a projector on the representation  $[2, \ell]$  when acting from the right<sup>10</sup>.

The restriction to those elements in  $E_{a_1 \dots a_{\ell-1}}{}^{b_1 b_2}$  belonging strictly to the affine subalgebra gives an identification of  $\bar{K}_{\ell\beta}^\alpha$  as explained above. Here, the identification of the affine subalgebra within  $AE_3$  for all levels  $\ell > 0$  reads

$$\bar{K}_{\ell\beta}^\alpha = \varepsilon_{\beta\gamma} \underbrace{E_{1 \dots 1}}_{(\ell-1)}{}^{\gamma\alpha} \tag{3.18}$$

and for  $-\ell < 0$

$$\bar{K}_{-\ell\beta}^\alpha = \varepsilon^{\alpha\gamma} \underbrace{F^{1 \dots 1}}_{(\ell-1)}{}_{\gamma\beta}. \tag{3.19}$$

These formulae specialize (2.30) and (2.31) to the case  $n = 3$ . Note that we use Euclidean conventions resulting in  $\varepsilon^{23} = \varepsilon_{23} = 1$ . It is easy to check that indeed ( $\ell \geq 1$ )

$$\left[ \hat{d}, \underbrace{E_{1 \dots 1}}_{(\ell-1)}{}^{\alpha\beta} \right] = -\ell \underbrace{E_{1 \dots 1}}_{(\ell-1)}{}^{\alpha\beta} \tag{3.20}$$

and

$$\left[ \hat{c}, \underbrace{E_{1 \dots 1}}_{(\ell-1)}{}^{\alpha\beta} \right] = 0. \tag{3.21}$$

Here the correction term in (3.5) is essential in order to obtain a vanishing result.

<sup>10</sup> That this is not naturally so can be seen, for instance, from considering the projector on the Riemann tensor symmetries.

Upon restriction to the affine subalgebra the projectors (3.17) simplify to

$$P_{\underbrace{1 \dots 1}_\ell}^{\beta_1 \beta_2} |^{c_1 \dots c_{\ell-1}}_{d_1 d_2} = \bar{\delta}_{1 \dots 1}^{c_1 \dots c_\ell} \bar{\delta}_{d_1 d_2}^{\beta_1 \beta_2} \tag{3.22}$$

for positive  $\ell$  and for negative  $\ell$  accordingly. The identification (3.18) together with the affine relations (2.21) then determines the commutators of the gradient representations. Note that the identification also fixes the normalization of the gradient tensor in the  $AE_3$  invariant form to be

$$\langle E_{a_1 \dots a_\ell}{}^{b_1 b_2} | F^{c_1 \dots c_k}{}_{d_1 d_2} \rangle = \begin{cases} P_{a_1 \dots a_\ell}{}^{b_1 b_2} |^{c_1 \dots c_\ell}{}_{d_1 d_2} & \text{for } \ell = k \\ 0 & \text{for } \ell \neq k \end{cases}, \tag{3.23}$$

by using equation (2.27).

The commutator of two  $E$  generators on levels  $\ell$  and  $k$  is then

$$[E_{a_1 \dots a_{\ell-1}}{}^{b_1 b_2}, E_{c_1 \dots c_{k-1}}{}^{d_1 d_2}] = -E_{a_1 \dots a_{\ell-1} c_1 \dots c_{k-1} e}{}^{d_1 (b_1 \varepsilon^{b_2) d_2 e}} - E_{a_1 \dots a_{\ell-1} c_1 \dots c_{k-1} e}{}^{d_2 (b_1 \varepsilon^{b_2) d_1 e}} + \dots \tag{3.24}$$

The dots on the right-hand side indicate other possible  $AE_3$  generators appearing in this commutators which will, however, vanish upon restriction to the affine subalgebra  $A_1^{(1)}$ . The term on the right is fixed up to normalization from the tensor symmetries and the normalization is fixed from the affine subalgebra. A similar expression holds for  $[F, F]$ .

For  $\ell > k$  we obtain

$$[E_{a_1 \dots a_\ell}{}^{b_1 b_2}, F^{c_1 \dots c_k}{}_{d_1 d_2}] = -2P_{a_1 \dots a_\ell}{}^{b_1 b_2} |^{e_1 \dots e_\ell}{}_{f_1 f_2} P_{g_1 \dots g_k}{}^{h_1 h_2} |^{c_1 \dots c_k}{}_{d_1 d_2} \cdot \varepsilon_{e_\ell h_1 i} \delta_{h_2}^{f_1} \bar{\delta}_{e_1 \dots e_k}^{g_1 \dots g_k} E_{e_{k+1} \dots e_{\ell-1}}{}^{f_2 i} + \dots \tag{3.25}$$

Finally, level  $\ell + 1 > 0$  commutes with level  $-\ell - 1$  into the adjoint of  $\mathfrak{gl}_3$

$$[E_{a_1 \dots a_\ell}{}^{b_1 b_2}, F^{c_1 \dots c_\ell}{}_{d_1 d_2}] = P_{a_1 \dots a_\ell}{}^{b_1 b_2} |^{e_1 \dots e_\ell}{}_{f_1 f_2} P_{g_1 \dots g_\ell}{}^{h_1 h_2} |^{c_1 \dots c_\ell}{}_{d_1 d_2} \cdot (\delta_{e_1}^{g_1} \dots \delta_{e_\ell}^{g_\ell} \delta_{h_1}^{f_1} \delta_{h_2}^{f_2} K - 2\delta_{e_1}^{g_1} \dots \delta_{e_\ell}^{g_\ell} \delta_{h_1}^{f_1} K^{f_2}{}_{h_2} + \ell \delta_{h_1}^{f_1} \delta_{h_2}^{f_2} \delta_{e_1}^{g_1} \dots \delta_{e_{\ell-1}}^{g_{\ell-1}} K^{g_\ell}{}_{e_\ell}). \tag{3.26}$$

This commutator is exact since there are no other Lie algebra elements besides those of  $\mathfrak{gl}_3$  at level  $\ell = 0$ . The omitted terms (= dots) in the other  $AE_3$  commutators are necessary for the Jacobi identities. This phenomenon first occurs at level  $\ell = 3$ , since this is the first time a non-gradient representation is present (cf table 1). Checking the Jacobi identities using solely the gradient generators shows a violation which is resolved when one also takes into account the  $\ell = 3$  representation with Dynkin labels [1, 1].

#### 4. Affine truncation of the $AE_n/K(AE_n)$ $\sigma$ -model

Having clarified the embeddings of the subgroups and having determined the gradient commutators, we next turn to the one-dimensional ‘geodesic’  $\sigma$ -model over the coset space  $AE_n/K(AE_n)$ . This model is governed by the Lagrangian [10]

$$L = \frac{1}{2n} \langle \mathcal{P} | \mathcal{P} \rangle, \tag{4.1}$$

where all quantities depend only on the affine parameter (‘time’)  $t$ , and  $n(t)$  is the lapse function required for reparametrization invariance  $t \rightarrow \tilde{t}(t)$ . The quantity  $\mathcal{P}(t)$  and the  $K(AE_n)$  gauge connection  $\mathcal{Q}(t)$  are determined from the Cartan form

$$\partial_t \mathcal{V} \mathcal{V}^{-1} = \mathcal{Q} + \mathcal{P} \tag{4.2}$$

with  $\mathcal{V}(t) \in AE_n/K(AE_n)$ . By construction,  $\mathcal{Q}$  belongs to the compact subalgebra  $K(AE_n)$  (fixed by the Chevalley involution, cf equation (2.3)), and  $\mathcal{P}$  to its complement in  $AE_n$ . With these definitions and notation, the equations of motion assume the simple looking form [14]

$$n\partial_t(n^{-1}\mathcal{P}) = [\mathcal{Q}, \mathcal{P}]. \tag{4.3}$$

To get these equations into the standard second-order form, we would have to choose coordinates on the  $AE_n/K(AE_n)$  coset manifold and to substitute them into (4.2), but we will skip this step here. Although these objects are highly formal constructs at this point, readers need not worry about our lack of knowledge of what the ‘group’  $AE_3$  really is, because the truncated level expansion provides an algorithm which is such that all operations take place in the Lie algebra, and involve only a finite number of steps if one truncates at finite level. As we will show these steps remain well defined for the *affine truncation* where one retains an infinite number of Lie algebra elements.

Varying the lapse  $n$ , we get the (Hamiltonian) constraint

$$\langle \mathcal{P}|\mathcal{P} \rangle = 0. \tag{4.4}$$

Consequently, the ‘trajectory’ described by (4.3) on the coset manifold is a *null geodesic*.

The system described by this model is formally integrable [9]. For every solution  $\mathcal{V}$  giving rise to  $\mathcal{Q}$  and  $\mathcal{P}$  satisfying (4.3) we can define a conserved (Noether) charge

$$\mathcal{J} = n^{-1}\mathcal{V}^{-1}\mathcal{P}\mathcal{V}, \quad \partial_t\mathcal{J} = 0 \tag{4.5}$$

taking values in the Lie algebra and obeying a null condition

$$\langle \mathcal{J}|\mathcal{J} \rangle = 0. \tag{4.6}$$

Similarly, we can define a (symmetric) ‘metric’  $\mathcal{M}$  associated with the  $\infty$ -bein  $\mathcal{V}$  by  $\mathcal{M} = \mathcal{V}^T\mathcal{V}$ , which is related to  $\mathcal{J}$  via

$$\mathcal{J} = \frac{1}{2}n^{-1}\mathcal{M}^{-1}\partial_t\mathcal{M}. \tag{4.7}$$

Equation (4.5) implies the following constraint on  $\mathcal{J}$ , and hence on the initial data,

$$\mathcal{J}^T = \mathcal{M}\mathcal{J}\mathcal{M}^{-1}. \tag{4.8}$$

The general solution for  $\mathcal{M}$  can be (formally) written as

$$\mathcal{M}(t) = \exp(v(t)\mathcal{J}^T)\mathcal{M}(0)\exp(v(t)\mathcal{J}). \tag{4.9}$$

where

$$v(t) := \int_{t_0}^t n(t') dt', \tag{4.10}$$

with  $t_0$  being some arbitrary initial time. Translated back to  $\mathcal{V}(t)$ , the general solution is

$$\mathcal{V}(t) = k(t)\mathcal{V}(0)\exp(v(t)\mathcal{J}). \tag{4.11}$$

Here,  $k(t)$  belongs to the compact subgroup  $K(AE_n)$  and is not determined by the equations of motion. This indeterminacy corresponds to the freedom of choosing a gauge for  $\mathcal{V}(t)$ . We can now work out the above equations of motion (4.3) level by level, following the low-level results of [10].

4.1. Restriction to  $A_{n-2}^{(1)}$

We will now restrict the one-dimensional geodesic  $\sigma$ -model (4.1) to its *affine subsector*; this is a consistent truncation. As we explained, the affine sector originates from the zeroth level in the  $A_{n-2}^{(1)}$  analysis. To this end we define

$$\begin{aligned} S_m^{\alpha\beta} &= \frac{1}{2}(\bar{K}_{m\beta}^\alpha + \bar{K}_{-m\alpha}^\beta), \\ J_m^{\alpha\beta} &= \frac{1}{2}(\bar{K}_{m\beta}^\alpha - \bar{K}_{-m\alpha}^\beta), \end{aligned} \tag{4.12}$$

which obey the symmetry relations

$$S_m^{\alpha\beta} = S_{-m}^{\beta\alpha}, \quad J_m^{\alpha\beta} = -J_{-m}^{\beta\alpha}. \tag{4.13}$$

The generators  $S_m^{\alpha\beta}, J_m^{\alpha\beta}$  for  $m \geq 0$ , together with  $\hat{c}$  and  $\hat{d}$ , define a basis of  $A_{n-2}^{(1)}$ . The commutation relations of  $A_{n-2}^{(1)}$  in this basis are

$$[\hat{d}, S_m^{\alpha\beta}] = -m J_m^{\alpha\beta}, \quad [\hat{d}, J_m^{\alpha\beta}] = -m S_m^{\alpha\beta} \tag{4.14}$$

and, for  $m, n \geq 0$ ,

$$[J_m^{\alpha\beta}, S_n^{\gamma\delta}] = \frac{1}{2}\delta^{\beta\gamma} S_{m+n}^{\alpha\delta} + \frac{1}{2}\delta^{\beta\delta} S_{m-n}^{\alpha\gamma} - \frac{1}{2}\delta^{\alpha\gamma} S_{m-n}^{\delta\beta} - \frac{1}{2}\delta^{\alpha\delta} S_{m+n}^{\gamma\beta} + \frac{1}{2}m\hat{c}\delta_{m,n}\delta^{\alpha\gamma}\delta^{\beta\gamma}, \tag{4.15}$$

$$[J_m^{\alpha\beta}, J_n^{\gamma\delta}] = \frac{1}{2}\delta^{\beta\gamma} J_{m+n}^{\alpha\delta} - \frac{1}{2}\delta^{\beta\delta} J_{m-n}^{\alpha\gamma} + \frac{1}{2}\delta^{\alpha\gamma} J_{m-n}^{\delta\beta} - \frac{1}{2}\delta^{\alpha\delta} J_{m+n}^{\gamma\beta}, \tag{4.16}$$

$$[S_m^{\alpha\beta}, S_n^{\gamma\delta}] = \frac{1}{2}\delta^{\beta\gamma} J_{m+n}^{\alpha\delta} + \frac{1}{2}\delta^{\beta\delta} J_{m-n}^{\alpha\gamma} - \frac{1}{2}\delta^{\alpha\gamma} J_{m-n}^{\delta\beta} - \frac{1}{2}\delta^{\alpha\delta} J_{m+n}^{\gamma\beta}. \tag{4.17}$$

Observe that the central charge drops out in the  $[J, J]$  and  $[S, S]$  commutators, but is present in the ‘mixed’ commutators  $[J, S]$ . The elements  $J_m$  (with  $m \geq 0$ ) generate the (centreless) ‘maximal compact’ subalgebra  $K(A_{n-2}^{(1)}) \equiv K(\widehat{\mathfrak{sl}_{n-1}})$ .<sup>11</sup>

Next, we expand the Cartan form in this basis as in (4.2), with

$$\begin{aligned} \mathcal{Q}(t) &= Q_{\alpha\beta}^{(0)} J_0^{\alpha\beta} + \sum_{m \geq 1} Q_{\alpha\beta}^{(m)} J_m^{\alpha\beta}, \\ \mathcal{P}(t) &= P_{\alpha\beta}^{(0)} S_0^{\alpha\beta} + \sum_{m \geq 1} P_{\alpha\beta}^{(m)} S_m^{\alpha\beta} + \partial_t \hat{\rho} \hat{\rho}^{-1} \hat{d} + \partial_t \sigma \hat{c}. \end{aligned} \tag{4.18}$$

The quantity  $\hat{\rho}$  will be seen to be directly related to the ‘dilaton’ field (= volume density for the internal manifold), while  $e^\sigma$  is related to the conformal factor in the dimensional reduction of Einstein’s theory to two dimensions. Observe that the position of indices no longer matters, as the tensors appearing on the rhs are to be regarded as  $SO(n - 1)$  tensors only. Also,

$$Q_{\alpha\beta}^{(0)} = -Q_{\beta\alpha}^{(0)}, \quad P_{\alpha\beta}^{(0)} = +P_{\beta\alpha}^{(0)} \tag{4.19}$$

whereas the higher modes are not subject to any such symmetry restrictions.

In this general, non-gauge-fixed form, the equations of motion (4.3) read

$$\begin{aligned} n\partial_t(n^{-1}\hat{\rho}^{-1}\partial_t\hat{\rho}) &= 0, \quad n\partial_t(n^{-1}\partial_t\sigma) = \frac{1}{2}\sum_{m=1}^{\infty} m Q_{\alpha\beta}^{(m)} P_{\alpha\beta}^{(m)}, \\ n\partial_t(n^{-1}P_{\alpha\beta}^{(0)}) &= 2Q_{(\alpha|\gamma}^{(0)} P_{\gamma|\beta)}^{(0)} + \frac{1}{2}\sum_{m=1}^{\infty} (Q_{(\alpha|\gamma}^{(m)} P_{\beta)\gamma}^{(m)} - Q_{\gamma(\beta}^{(m)} P_{|\gamma|\alpha)}^{(m)}) \end{aligned} \tag{4.20}$$

<sup>11</sup> That the latter is *not* a Kac–Moody algebra is demonstrated in appendix A. Also note that the additional CSA elements  $\hat{c}$  and  $\hat{d}$  do not survive the projection to the compact subalgebra and hence  $K(A_{n-2}^{(1)})$  equals the compact subalgebra of the loop algebra  $\widehat{\mathfrak{sl}_{n-1}}$ .

for the level-zero degrees of freedom, where the vertical bars on the rhs of the third equation indicate that symmetrization should be performed only over the indices  $\alpha$  and  $\beta$ . At levels  $\ell \geq 1$ , we obtain (recall that there is no symmetry under the exchange of  $\alpha$  and  $\beta$ )

$$n \partial_t (n^{-1} P_{\alpha\beta}^{(\ell)}) = \ell \hat{\rho}^{-1} \partial_t \hat{\rho} Q_{\alpha\beta}^{(\ell)} + \frac{1}{2} \sum_{m=0}^{\ell} (Q_{\alpha\gamma}^{(m)} P_{\gamma\beta}^{(\ell-m)} - Q_{\gamma\beta}^{(m)} P_{\alpha\gamma}^{(\ell-m)}) + \frac{1}{2} \sum_{m=0}^{\infty} (Q_{\alpha\gamma}^{(\ell+m)} P_{\beta\gamma}^{(m)} - Q_{\gamma\beta}^{(\ell+m)} P_{\gamma\alpha}^{(m)} + Q_{\beta\gamma}^{(m)} P_{\alpha\gamma}^{(\ell+m)} - Q_{\gamma\alpha}^{(m)} P_{\gamma\beta}^{(\ell+m)}). \tag{4.21}$$

Finally, the Hamiltonian constraint in a general gauge is

$$\langle \mathcal{P} | \mathcal{P} \rangle = -2 \hat{\rho}^{-1} \partial_t \hat{\rho} \partial_t \sigma + P_{\alpha\beta}^{(0)} P_{\alpha\beta}^{(0)} + \frac{1}{2} \sum_{m \geq 1} P_{\alpha\beta}^{(m)} P_{\alpha\beta}^{(m)} = 0. \tag{4.22}$$

The first equation in (4.20) can be integrated straightforwardly, with the result

$$n(t) = \partial_t \ln \hat{\rho}(t) \quad \Rightarrow \quad v(t) = \ln \hat{\rho}(t), \tag{4.23}$$

where integration constants have been chosen conveniently. Hence, the choice of the function  $\hat{\rho}(t)$  can be viewed as a choice of gauge for the lapse  $n(t)$ ; this function is only subject to the requirement that it be a monotonic function of the affine parameter  $t$ . This result can be plugged into the general solution (4.11) to give

$$\mathcal{V}(t) = k(t) \mathcal{V}(0) \exp(\ln \hat{\rho}(t) \mathcal{J}). \tag{4.24}$$

Although the above equations of motion constitute a consistent truncation of the full  $AE_n/K(AE_n)$   $\sigma$ -model, one may ask how they embed into the latter. Within  $AE_n/K(AE_n)$ , the above equations of motion may receive new contributions from those ‘non-gradient fields’ whose associated Lie algebra elements ‘commute back’ into the affine subalgebra. The role and significance of these extra contributions is not clear.

We emphasize that expression (4.18) is the most general, because we have not chosen a gauge for the local subgroup  $K(A_{n-2}^{(1)})$ . In other words, *no matter which gauge is chosen, the equations of motion can always be cast into the forms (4.20), (4.21) and (4.22).*

#### 4.2. Triangular gauge

The equations of motion derived above can now be considered in various gauges, thereby fixing the factor  $k(t)$  in (4.11). A convenient choice, and one that has been used almost exclusively in the previous work, is the *triangular gauge*, for which the  $\infty$ -bein  $\mathcal{V}(t)$  depends only on the level  $\ell \geq 0$  fields. In this gauge, one obtains

$$\partial_t \mathcal{V} \mathcal{V}^{-1} = Q_{\alpha\beta}^{(0)} J_0^{\alpha\beta} + P_{\alpha\beta}^{(0)} S_0^{\alpha\beta} + \sum_{m \geq 1} P_{\alpha\beta}^{(m)} \bar{K}_{m\beta}^{\alpha} + \partial_t \hat{\rho} \hat{\rho}^{-1} \hat{d} + \partial_t \sigma \hat{c}. \tag{4.25}$$

Consequently,

$$Q_{\alpha\beta}^{(m)} = P_{\alpha\beta}^{(m)} \quad \text{for } m \geq 1. \tag{4.26}$$

Note that the second term on the rhs of (4.21) vanishes for this choice of gauge, in agreement with previous calculations [14]. While  $P_{\alpha\beta}^{(0)}$  is symmetric,  $P_{\alpha\beta}^{(m)}$  contains both symmetric and antisymmetric parts for  $m > 0$ , and we can therefore decompose it into irreducible  $\mathfrak{so}_{n-1}$  representations:

$$P_{\alpha\beta}^{(m)} = \bar{P}_{\alpha\beta}^{(m)} + \bar{Q}_{\alpha\beta}^{(m)} \tag{4.27}$$

with

$$\bar{P}_{\alpha\beta}^{(m)} = \bar{P}_{\beta\alpha}^{(m)}, \quad \bar{Q}_{\alpha\beta}^{(m)} = -\bar{Q}_{\beta\alpha}^{(m)}, \quad (4.28)$$

remembering that  $\bar{P}_{\alpha\beta}^{(m)}$  is traceless. With these definitions the dynamical equations (4.20) and (4.21) become

$$n\partial_t(n^{-1}\hat{\rho}^{-1}\partial_t\hat{\rho}) = 0, \quad (4.29)$$

$$n\partial_t(n^{-1}\partial_t\sigma) = \frac{1}{2} \sum_{m \geq 1} m(\bar{Q}_{\alpha\beta}^{(m)}\bar{Q}_{\alpha\beta}^{(m)} + \bar{P}_{\alpha\beta}^{(m)}\bar{P}_{\alpha\beta}^{(m)}), \quad (4.30)$$

$$n\partial_t(n^{-1}\bar{P}_{\alpha\beta}^{(0)}) = 2 \sum_{m \geq 0} \bar{Q}_{(\alpha|\gamma}^{(m)}\bar{P}_{\gamma|\beta)}^{(m)}, \quad (4.31)$$

$$n\hat{\rho}^\ell\partial_t(n^{-1}\hat{\rho}^{-\ell}\bar{P}_{\alpha\beta}^{(\ell)}) = 2 \sum_{m \geq 0} (\bar{Q}_{(\alpha|\gamma}^{(\ell+m)}\bar{P}_{\gamma|\beta)}^{(m)} + \bar{Q}_{(\alpha|\gamma}^{(m)}\bar{P}_{\gamma|\beta)}^{(\ell+m)}), \quad (4.32)$$

$$n\hat{\rho}^\ell\partial_t(n^{-1}\hat{\rho}^{-\ell}\bar{Q}_{\alpha\beta}^{(\ell)}) = 2 \sum_{m \geq 0} (-\bar{Q}_{\gamma[\alpha}^{(\ell+m)}\bar{Q}_{\beta]\gamma}^{(m)} + \bar{P}_{\gamma[\alpha}^{(\ell+m)}\bar{P}_{\beta]\gamma}^{(m)}), \quad (4.33)$$

where the last two equations hold for  $\ell \geq 1$ . The Hamiltonian constraint (4.22) now reads

$$\langle \mathcal{P} | \mathcal{P} \rangle = -2\hat{\rho}^{-1}\partial_t\hat{\rho}\partial_t\sigma + \bar{P}_{\alpha\beta}^{(0)}\bar{P}_{\alpha\beta}^{(0)} + \frac{1}{2} \sum_{m \geq 1} (\bar{Q}_{\alpha\beta}^{(m)}\bar{Q}_{\alpha\beta}^{(m)} + \bar{P}_{\alpha\beta}^{(m)}\bar{P}_{\alpha\beta}^{(m)}) = 0. \quad (4.34)$$

Are there other viable gauge choices? While the triangular or Borel parametrization is by no means the only possibility for finite-dimensional matrices, the situation is more subtle for infinite-dimensional groups. In fact, it appears that so far the triangular parametrization is the only manageable one for indefinite KMAs. Nevertheless, indications were found recently [34] that the triangular parametrization must be relaxed if one is to include M theoretic corrections, and to extend the ‘dictionary’ beyond the very first few levels. Here we simply emphasize again that independently of the gauge, the equations of motion can always be written in the forms (4.20), (4.21) and (4.22). The only difference is that the equality (4.26) will fail to hold in a non-triangular gauge.

## 5. Comparison with two-dimensional reduction

### 5.1. Relation to higher-dimensional vielbein

The infinite ‘matrix’  $\mathcal{V}(t) \in AE_n/K(AE_n)$  can be thought of as an  $\infty$ -bein analogous to the vielbein of general relativity. Indeed, it contains the (spatial) vielbein as a finite-dimensional ‘submatrix’, if one restricts attention to the level zero sector, i.e. upon restriction of the full coset to the finite-dimensional coset space  $GL(n)/SO(n)$  (note that we keep the  $GL(1)$  factor here as a remnant of the full algebra). As is well known, pure gravity in  $(n+1)$  spacetime dimensions exhibits a hidden  $SL(n-1)$  symmetry after reduction to three dimensions. This symmetry is obtained through an enlargement of the manifest  $SL(n-2)$  symmetry upon dualization of  $(n-2)$  Kaluza–Klein vectors to  $(n-2)$  scalar fields. After further reduction to two dimensions the system becomes integrable with a Lax pair, and admits an  $A_{n-2}^{(1)}$  symmetry as a ‘solution generating group’ [4–6]; for  $n=3$  this group is known as the Geroch group [1, 2]. Together with the  $GL(n)$  acting on the spatial  $n$ -bein, we thus recover precisely the subgroups of  $AE_n$  discussed in the foregoing section.



Let us therefore spell out this correspondence in a little more detail, in order to facilitate the comparison between the one-dimensional geodesic  $\sigma$ -model introduced in the foregoing section, and the  $D = 2$  theory to be discussed in the following subsection. For this purpose, we need to fix gauges such that

$$E_M^A = \left( \begin{array}{c|c} N & 0 \\ \hline 0 & e_m^a \end{array} \right). \tag{5.1}$$

This equation displays the decomposition of the  $(n + 1)$ -bein into a lapse factor  $N$ , and a spatial  $n$ -bein  $e_m^a$ ; the shift variables (which enforce the spatial diffeomorphism constraint) have been set to zero, as required for the comparison with the Kac–Moody  $\sigma$ -model. As we said already, the spatial  $n$ -bein can be viewed as the level-0 sector of the  $\infty$ -bein  $\mathcal{V}$ , and thereby of the coset  $AE_3/K(AE_3)$ . As is well known [10], the restriction of the  $\sigma$ -model Lagrangian (4.1) to the  $\ell = 0$  sector coincides precisely with the dimensional reduction of Einstein’s theory to one time dimension.

In section 5.2, we will compare the one-dimensional  $\sigma$ -model equations of motion with those obtained in the reduction of Einstein’s theory to  $(1 + 1)$  dimension (which depend on time and one extra spatial coordinate). For this comparison, one further step is required, namely the split of the spatial  $n$ -bein in (5.1) according to

$$E_M^A = \left( \begin{array}{c|cc} \lambda & 0 & 0 \\ \hline 0 & \lambda & 0 \\ 0 & 0 & \rho^{1/(n-1)} \bar{e}_{\bar{m}}^{\bar{a}} \end{array} \right). \tag{5.2}$$

Here we have singled out one (the first) spatial direction, and decomposed the remaining  $(n - 1)$ -bein into a unimodular part, and its determinant. So  $\det(\bar{e}_{\bar{m}}^{\bar{a}}) = 1$ , and  $\rho$  measures the volume of the  $(n - 1)$ -dimensional internal space (‘internal’ from the  $(1 + 1)$ -dimensional perspective, of course). In addition, we have adopted the conformal gauge for the zweibein, with the conformal factor  $\lambda$ , thus tying the lapse to the  $(11)$  component of the spatial metric. Finally, setting the remaining elements of the first column and the first row of  $e_m^a$  to zero is related to the hypersurface orthogonality of the Killing vectors usually assumed in the reduction to  $(1 + 1)$  dimensions [2, 5].

The special parametrization (5.2) results in the following non-vanishing components of the coefficients of anholonomicity  $\Omega_{ABC} \equiv 2E_A^M E_B^N \partial_{[M} E_{N]C}$  (with  $\partial_x \equiv \partial/\partial x^1$ )

$$\Omega_{0\bar{b}\bar{c}} = \frac{1}{n-1} \lambda^{-1} \delta_{\bar{b}\bar{c}} \partial_t \rho \rho^{-1} + \lambda^{-1} \bar{e}_{\bar{b}}^{\bar{m}} \partial_t \bar{e}_{\bar{m}\bar{c}}, \tag{5.3}$$

$$\Omega_{1\bar{b}\bar{c}} = \frac{1}{n-1} \lambda^{-1} \delta_{\bar{b}\bar{c}} \partial_x \rho \rho^{-1} + \lambda^{-1} \bar{e}_{\bar{b}}^{\bar{m}} \partial_x \bar{e}_{\bar{m}\bar{c}}, \tag{5.4}$$

$$\Omega_{011} = \lambda^{-2} \partial_t \lambda, \tag{5.5}$$

$$\Omega_{010} = -\lambda^{-2} \partial_x \lambda. \tag{5.6}$$

The spatial components  $\Omega_{abc}$  of the anholonomicity are related by duality to the first gradient representation (2.11), which we called dual graviton in 2.4. The irreducibility constraint (2.12) on this representation is equivalent to the tracelessness condition  $\Omega_{abb} = 0$  [9, 14]. In the present two-dimensional context this condition reduces to

$$\Omega_{1bb} = \Omega_{1\bar{b}\bar{b}} = \lambda^{-1} \partial_x \rho \rho^{-1} = 0. \tag{5.7}$$

Therefore, the irreducibility condition  $\Omega_{abb} = 0$  holds if and only if  $\rho = \rho(t)$ . In the following section, we will arrive at the same conclusion by a somewhat different route.

The parametrization (5.2) also yields the determinant of the original  $(n + 1)$ -bein to be

$$\det(E) = N\sqrt{g} = \lambda^2 \rho \quad (5.8)$$

where  $\sqrt{g} \equiv \det e_m^a$ . Using the identification between the original lapse  $N$  and the  $\sigma$ -model lapse  $n$  derived in [13, 14], we can relate the  $\sigma$ -model lapse to the internal volume density  $\rho$  via

$$n = Ng^{-1/2} = \rho^{-1}. \quad (5.9)$$

This result will be useful below.

To deduce the relation between the field  $\sigma$  and the conformal factor  $\lambda$ , we compute the contribution from (5.2) to the diagonal part of  $\partial_t e_a^m e_m^b$ , namely

$$\partial_t e_a^m e_m^b = -\lambda^{-1} \partial_t \lambda K^1_1 - \frac{1}{n-1} \rho^{-1} \partial_t \rho K^\alpha_\alpha + \dots \quad (5.10)$$

where the dots stand for the contributions from the off-diagonal degrees of freedom. Rewriting this in terms of the CSA elements  $\hat{c}$  and  $\hat{d}$ , cf (2.23), we obtain

$$\partial_t [\rho^{(n-2)/2(n-1)} \lambda] [\rho^{(n-2)/2(n-1)} \lambda]^{-1} \hat{c} + \partial_t \rho \rho^{-1} \hat{d}. \quad (5.11)$$

Comparing (5.11) and (4.18) we conclude that the conformal factor should be identified as<sup>12</sup>

$$e^\sigma \equiv \lambda \rho^{\frac{n-2}{2(n-1)}} \quad (5.12)$$

while  $\hat{\rho}$  indeed agrees with the internal volume density  $\rho$ . This is consistent with the result from the irreducibility constraint (5.7) that  $\rho$  is a function of time only.

### 5.2. $\sigma$ -model equations of motion in $(1 + 1)$ dimensions

Next we write the equations of motion for the  $D = 2$  theory as obtained from an  $SL(n - 1)/SO(n - 1)$   $\sigma$ -model coupled to gravity in two dimensions, and then compare them to the equations of motion of the one-dimensional  $\sigma$ -model derived in the foregoing section. This will not only allow us to extend previous results on the matching, but also, more importantly, to exhibit those terms which do *not* match, and where the dictionary needs to be modified if one is to include higher order spatial gradients. The main advantage of the  $(1 + 1)$  theory is that the mismatch assumes the simplest possible form, and can therefore be scrutinized in full detail.

The basic object of the  $(1 + 1)$ -dimensional theory is a matrix  $V(t, x)$ , which is an element of the coset space  $SL(n - 1)/SO(n - 1)$ , and which is the analogue of  $\mathcal{V}(t)$  of the previous section. The corresponding sector of the theory is governed by the standard  $\sigma$ -model Lagrangian on the worldsheet in a gravitational background. The corresponding Cartan form belongs to the horizontal  $\mathfrak{sl}_{n-1}$

$$\partial_\mu V V^{-1} = Q_\mu + P_\mu \equiv Q_{\mu\alpha\beta} J_0^{\alpha\beta} + P_{\mu\alpha\beta} S_0^{\alpha\beta} \quad (5.13)$$

with the gauge field  $Q_\mu \in \mathfrak{so}_{n-1}$  and  $P_\mu \in \mathfrak{sl}_{n-1} \ominus \mathfrak{so}_{n-1}$ , and  $\mu, \nu, \dots = t, x$ . The equations of motion read

$$D_\mu (\rho \sqrt{-\gamma} \gamma^{\mu\nu} P_\nu) = 0, \quad (5.14)$$

where  $D_\mu P_\nu \equiv \partial_\mu P_\nu - [Q_\mu, P_\nu]$  is the  $\mathfrak{so}_{n-1}$  covariant derivative, and  $\gamma_{\mu\nu}$  is the worldsheet metric, with inverse  $\gamma^{\mu\nu}$  and determinant  $\gamma$ . The additional dependence on the ‘dilaton’  $\rho$  in

<sup>12</sup> The same correction to the conformal factor is also obtained (for  $n = 3$ ) in the Matzner–Misner coset formulation of gravity reduced to two spacetime dimensions [5], see also [35].

this equation is a remnant of the reduction from higher dimensions. In the conformal gauge  $\gamma_{\mu\nu} = \lambda^2 \eta_{\mu\nu}$  (cf (5.2)), this equation simplifies to

$$D^\mu(\rho P_\mu) = 0, \tag{5.15}$$

where indices are now to be raised and lowered with the Minkowski metric  $\eta_{\mu\nu}$ . Writing these equations in terms of  $t$  and  $x$  components, we get

$$\rho^{-1} \partial_t(\rho P_t) - \rho^{-1} \partial_x(\rho P_x) = [Q_t, P_t] - [Q_x, P_x]. \tag{5.16}$$

In addition, from (5.13), we have the integrability conditions

$$\partial_t P_x - \partial_x P_t = [Q_t, P_x] - [Q_x, P_t], \quad \partial_t Q_x - \partial_x Q_t = [Q_t, Q_x] + [P_t, P_x]. \tag{5.17}$$

with the gauge connection  $Q_t, Q_x \in \mathfrak{so}_{n-1}$ . Here we have written explicitly all covariantizations in order to facilitate the comparison with the equations of motion (4.29) following from the one-dimensional  $\sigma$ -model.

In conformal gauge, the ‘dilaton’  $\rho$  obeys a free field equation, namely

$$(\partial_t^2 - \partial_x^2)\rho(t, x) = 0. \tag{5.18}$$

This equation is satisfied in particular if  $\rho$  equals one of the two-dimensional coordinates or a linear combination thereof (Weyl canonical coordinates). Since we are interested in cosmological applications, for which  $\rho$  is a timelike coordinate, we choose<sup>13</sup>

$$\rho(t, x) = t. \tag{5.19}$$

To see that this choice matches with the one-dimensional  $\sigma$ -model, we substitute (5.9) into the first equation of (4.29), which gives

$$\rho^{-1} \partial_t^2 \rho = 0 \tag{5.20}$$

and indeed agrees with (5.18) if  $\rho$  is independent of  $x$ . Furthermore, independence of  $\rho$  of  $x$  is consistent with the Young irreducibility constraint in the  $\sigma$ -model as explained above in (5.7).

The first-order equations for the conformal factor  $\lambda$  are just the Hamiltonian and diffeomorphism constraints, respectively, and read

$$\partial_t \rho \lambda^{-1} \partial_t \lambda + \partial_x \rho \lambda^{-1} \partial_x \lambda = \frac{1}{2} \rho \operatorname{Tr}(P_t^2 + P_x^2), \tag{5.21}$$

$$\partial_t \rho \lambda^{-1} \partial_x \lambda + \partial_x \rho \lambda^{-1} \partial_t \lambda = \rho \operatorname{Tr} P_t P_x. \tag{5.22}$$

With (5.19) they simplify to

$$\lambda^{-1} \partial_t \lambda = \frac{t}{2} \operatorname{Tr}(P_t^2 + P_x^2), \quad \lambda^{-1} \partial_x \lambda = t \operatorname{Tr} P_t P_x. \tag{5.23}$$

There is also a second-order equation for the conformal factor, which reads, for  $\rho = t$ ,

$$t^{-1} \partial_t(t \lambda^{-1} \partial_t \lambda) - \partial_x(\lambda^{-1} \partial_x \lambda) = \operatorname{Tr} P_x P_x. \tag{5.24}$$

In order to compare the equations (5.16)–(5.24) with the equations of the one-dimensional affine  $\sigma$ -model (4.29)–(4.33), we must truncate both models appropriately. On the side of the one-dimensional model we do this by restricting the expansion to levels  $|\ell| \leq 1$ . Making use of the equality (5.9) and the triangular gauge (4.28), the equations of motion (4.29)–(4.33) are thus truncated to

$$\begin{aligned} \rho^{-1} \partial_t(\rho P_{\alpha\beta}^{(0)}) &= 2\bar{Q}_{\gamma[\alpha}^{(0)} P_{\beta]\gamma}^{(0)} + 2\bar{Q}_{\gamma[\alpha}^{(1)} \bar{P}_{\beta]\gamma}^{(1)} \\ \partial_t \bar{P}_{\alpha\beta}^{(1)} &= 2\bar{Q}_{\gamma[\alpha}^{(0)} \bar{P}_{\beta]\gamma}^{(1)} + 2\bar{Q}_{\gamma[\alpha}^{(1)} \bar{P}_{\beta]\gamma}^{(0)} \\ \partial_t \bar{Q}_{\alpha\beta}^{(1)} &= -2\bar{Q}_{\gamma[\alpha}^{(1)} \bar{Q}_{\beta]\gamma}^{(0)} + 2\bar{P}_{\gamma[\alpha}^{(1)} \bar{P}_{\beta]\gamma}^{(0)} \end{aligned} \tag{5.25}$$

<sup>13</sup> For spacelike  $\rho$ , we would obtain a variant of the so-called Einstein–Rosen waves, see e.g. [3].

$$\rho^{-1} \partial_t (\rho \partial_t \sigma) = \frac{1}{2} (\bar{Q}_{\alpha\beta}^{(1)} \bar{Q}_{\alpha\beta}^{(1)} + \bar{P}_{\alpha\beta}^{(1)} \bar{P}_{\alpha\beta}^{(1)}) \quad (5.26)$$

which must now be matched to (5.16) and (5.17). On the side of the (1 + 1) theory, we must restrict the  $x$ -dependence of the two-dimensional quantities such that  $P_t$  and  $P_x$  become independent of  $x$  (which is tantamount to keeping only first-order spatial gradients). Then the  $\sigma$ -model equations of motion match upon the identification

$$\bar{P}^{(0)}(t) \equiv P_t(t, x_0), \quad \bar{P}^{(1)}(t) \equiv P_x(t, x_0), \quad (5.27)$$

$$\bar{Q}^{(0)}(t) \equiv Q_t(t, x_0), \quad \bar{Q}^{(1)}(t) \equiv -Q_x(t, x_0), \quad (5.28)$$

where  $x_0$  is some fixed, but arbitrarily chosen spatial point. The ‘dictionary’ (5.27) must be supplemented by the relations (5.9) and (5.12), already derived before.

While many terms thus do match, there remain several discrepancies between these equations and those of the geodesic  $\sigma$ -model. First of all, there is a mismatch in the equation of motion for the conformal factor, which is similar to that encountered in the previous work, and which in particular involves a contribution  $\propto Q^{(1)} Q^{(1)}$ , which has no analogue involving the *gauge-variant* expression  $Q_x Q_x$  in (5.23). We interpret this mismatch as another indication of the impossibility to reconcile the higher-dimensional gauge invariance with the desired correspondence. Likewise, (5.24) has a term  $\partial_x^2 \sigma$  which has no analogue in (4.30). Furthermore, the spatial gradients  $\partial_x P_t$ ,  $\partial_x P_x$  and  $\partial_x Q_t$  in (5.16) are absent in the one-dimensional model. In view of these mismatches, we will now simplify the affine model yet further to an exactly solvable model.

## 6. Restriction to the ‘Heisenberg coset’ $\mathfrak{H}/K(\mathfrak{H})$

### 6.1. The $\mathfrak{H}/K(\mathfrak{H})$ $\sigma$ -model

We consider the (prototype) affine coset  $A_1^{(1)}/K(A_1^{(1)})$  restricted to its ‘Heisenberg subspace’  $\mathfrak{H}/K(\mathfrak{H})$ . In this way, we are able to define a one-to-one correspondence between a very limited, albeit non-trivial, class of solutions of Einstein’s equations (diagonal metrics with two commuting Killing vectors), and the null geodesic motion on the infinite-dimensional manifold  $\mathfrak{H}/K(\mathfrak{H})$ . To this end we restrict the affine algebra  $A_1^{(1)}$  to its Heisenberg subalgebra

$$\text{Lie}(\mathfrak{H}) := \widehat{\mathfrak{gl}}_1 \oplus \mathbb{R}\hat{c} \oplus \mathbb{R}\hat{d}. \quad (6.1)$$

Note that our terminology is slightly unusual in that the last summand is usually not considered to be part of the Heisenberg algebra, but required here in order to obtain a non-degenerate (indefinite) metric on  $\text{Lie}(\mathfrak{H})$ . Define

$$H_m = \frac{1}{\sqrt{2}} (\bar{K}_{m2}^2 - \bar{K}_{m3}^3) \quad (6.2)$$

for all  $m \in \mathbb{Z}$ . Then the non-vanishing commutators of the Heisenberg algebra are

$$[H_m, H_n] = m\delta_{m,-n}\hat{c}, \quad [\hat{d}, H_m] = -mH_m. \quad (6.3)$$

The symmetric and antisymmetric combinations are

$$\begin{aligned} S_m &:= \frac{1}{2}(H_m + H_{-m}) & \text{for } m \geq 0, \\ J_m &:= \frac{1}{2}(H_m - H_{-m}) & \text{for } m \geq 1. \end{aligned} \quad (6.4)$$

The  $J_m$  generate the maximal compact subgroup  $K(\mathfrak{H})$  of the extended Heisenberg group  $\mathfrak{H}$ . The commutation relations (6.3) in this basis are (recall that  $m, n \geq 0$ )

$$\begin{aligned} [S_m, S_n] &= 0, & [J_m, J_n] &= 0, \\ [J_m, S_n] &= \frac{1}{2}m\delta_{m,n}\hat{c}, & & \\ [\hat{d}, S_m] &= -mJ_m, & [\hat{d}, J_m] &= -mS_m. \end{aligned} \quad (6.5)$$

Parametrizing the coset space  $\mathfrak{H}/K(\mathfrak{H})$  in triangular gauge by

$$\mathcal{V}(t) = \exp[\ln \hat{\rho}(t)\hat{d}] \cdot \exp[\sigma(t)\hat{c}] \cdot \exp\left(\sum_{m \geq 0} \phi_m(t)H_m\right), \tag{6.6}$$

we find

$$\partial_t \mathcal{V} \mathcal{V}^{-1} = \sum_{m \geq 0} \hat{\rho}^{-m} \partial_t \phi_m H_m + \hat{\rho}^{-1} \partial_t \hat{\rho} \hat{d} + \partial_t \sigma \hat{c}. \tag{6.7}$$

Defining  $P_m(t) \equiv \hat{\rho}^{-m} \partial_t \phi_m$ , we have

$$\begin{aligned} \mathcal{P}(t) &= \sum_{m \geq 0} P_m(t) S_m + \hat{\rho}^{-1} \partial_t \hat{\rho}(t) \hat{d} + \partial_t \sigma(t) \hat{c} \\ \mathcal{Q}(t) &= \sum_{m \geq 1} P_m(t) J_m. \end{aligned} \tag{6.8}$$

Using the commutation relations (6.5) we get

$$[\mathcal{Q}, \mathcal{P}] = \sum_{m \geq 0} m P_m \hat{\rho}^{-1} \partial_t \hat{\rho} S_m + \frac{1}{2} \sum_{m \geq 1} m P_m^2 \hat{c}. \tag{6.9}$$

Therefore, the equations of motion of our model read

$$\begin{aligned} n \partial_t (n^{-1} \hat{\rho}^{-1} \partial_t \hat{\rho}) &= 0, \\ n \partial_t (n^{-1} \partial_t \sigma) &= \frac{1}{2} \sum_{m \geq 1} m \hat{\rho}^{-2m} (\partial_t \phi_m)^2, \\ n \partial_t (n^{-1} \hat{\rho}^{-m} \partial_t \phi_m) &= m \hat{\rho}^{-(m+1)} \partial_t \hat{\rho} \partial_t \phi_m. \end{aligned} \tag{6.10}$$

The Hamiltonian constraint takes the form

$$-2 \hat{\rho}^{-1} \partial_t \hat{\rho} \partial_t \sigma + P_0^2 + \frac{1}{2} \sum_{m \geq 1} P_m^2 = 0. \tag{6.11}$$

Using the insights from the preceding section we solve the first equation of (6.10) by setting  $n^{-1} = \hat{\rho} = t$ . Then the null geodesic trajectory in  $\mathfrak{H}/K(\mathfrak{H})$  is explicitly parametrized by

$$\left. \begin{aligned} \phi_0(t) &= p_0 \ln t + q_0 \\ \phi_m(t) &= \frac{1}{2m} p_m t^{2m} + q_m \quad (m > 0) \end{aligned} \right\} \Rightarrow P_m(t) = p_m t^{m-1} \quad (m \geq 0), \tag{6.12}$$

$$\sigma(t) = \frac{1}{2} p_0^2 \ln t + \sum_{m \geq 1} \frac{1}{8m} p_m^2 t^{2m} + \sigma_0,$$

where the coefficient of  $\ln t$  in the last line is determined by imposing the Hamiltonian constraint (6.11), which, however, does not fix  $\sigma_0$ . Because the model is explicitly solvable, we see in particular how the solution  $\mathcal{V}(t)$  depends on the most general initial data, i.e. the initial ‘coordinates’  $q_m$  and ‘momenta’  $p_m$  for  $m \geq 0$ .

The conserved current is, from (4.5),

$$\begin{aligned} \mathcal{J} = n^{-1} \mathcal{V}^{-1} \mathcal{P} \mathcal{V} &= n^{-1} \left[ \left( \partial_t \sigma - \frac{1}{2} \sum_{m \geq 1} m \hat{\rho}^{-m} \phi_m P_m \right) \hat{c} + \hat{\rho}^{-1} \partial_t \hat{\rho} \hat{d} \right. \\ &\quad \left. + P_0 H_0 + \frac{1}{2} \sum_{m \geq 1} (-2m \hat{\rho}^{-1} \partial_t \hat{\rho} \phi_m + \hat{\rho}^m P_m) H_m + \frac{1}{2} \sum_{m \geq 1} \hat{\rho}^{-m} P_m H_{-m} \right]. \end{aligned} \tag{6.13}$$

Plugging the solution (6.12), as well as  $n^{-1}(t) = t$ , into this expression yields

$$\mathcal{J} = \left( \frac{1}{2} p_0^2 - \frac{1}{2} \sum_{m \geq 1} m p_m q_m \right) \hat{c} + \hat{d} + \frac{1}{2} \sum_{m \geq 1} p_m H_{-m} + p_0 H_0 - \sum_{m \geq 1} m q_m H_m, \tag{6.14}$$

which is evidently conserved since it depends only on the initial data. Observe that neither  $q_0$  nor  $\sigma_0$  appear in  $\mathcal{J}$ . In agreement with the general analysis of [10], the initial momenta appear in the *lower*, and the initial coordinates in the *upper* triangular half of Lie ( $\mathfrak{h}$ ).

Under global  $\mathfrak{h}$  transformations  $g$ ,  $\mathcal{J}$  changes as  $\mathcal{J} \rightarrow g \mathcal{J} g^{-1}$ . Let

$$g(\alpha, \beta, \omega) := \exp(\ln \alpha \hat{c}) \cdot \exp(\ln \beta \hat{d}) \cdot \exp\left(\sum_{n \in \mathbb{Z}} \omega_n H_n\right), \tag{6.15}$$

then

$$g \mathcal{J} g^{-1} = \left( \frac{1}{2} p_0^2 - \frac{1}{2} \sum_{m \geq 1} m (p_m - 2m\omega_{-m})(q_m - \omega_m) \right) \hat{c} + \hat{d} - \sum_{m \geq 1} m \beta^{-m} (q_m - \omega_m) H_m + p_0 H_0 + \frac{1}{2} \sum_{m \geq 1} \beta^m (p_m - 2m\omega_{-m}) H_{-m}. \tag{6.16}$$

From this we immediately read the transformation of the initial coordinates  $q_m$  and momenta  $p_m$  (for  $m \geq 1$ ) under the action of  $\mathfrak{h}$ :

$$\left. \begin{aligned} q_m &\rightarrow \beta^{-m} (q_m - \omega_m) \\ p_m &\rightarrow \beta^m (p_m - 2m\omega_{-m}) \end{aligned} \right\} \quad (m \geq 1). \tag{6.17}$$

On  $\mathcal{V}$  the corresponding transformation is  $\mathcal{V} \rightarrow k \mathcal{V} g^{-1}$  where  $k(t)$  is the (local) compensator required to restore the triangular gauge, which does not contribute to the transformation of  $\mathcal{J}$ . By considering  $\mathcal{V}$  transformations we find constant shifts (which drop out in  $g \mathcal{J} g^{-1}$ )

$$\sigma_0 \rightarrow \sigma_0 - \alpha, \quad q_0 \rightarrow q_0 - \omega_0. \tag{6.18}$$

Hence, all the constants are shifted except the (Kasner) coefficient  $p_0$ . All fields except  $\sigma$  are inert under a transformation associated with the central term. This is no longer the case for the full  $AE_n$  model where  $\hat{c}$  ceases to be central.

### 6.2. Relation to polarized Gowdy cosmologies

Remarkably, there is a one-to-one correspondence between our model, and the ‘polarized’ Gowdy type cosmological model with diagonal metrics depending on two coordinates  $(t, x)$ .<sup>14</sup> Apart from possible reparametrizations of the time parameter  $t$ , *this correspondence works only after complete elimination of the gauge degrees of freedom on both sides*. In particular, the conformal factor must be treated as a *dependent* degree of freedom via the constraints (6.21). The relevant line elements can be written in the form (see e.g. [3])

$$ds^2 = \lambda^2 e^{-Z} (-dt^2 + dx^2) + t^2 e^{-Z} dy^2 + e^Z dz^2, \tag{6.19}$$

where the function  $Z = Z(t, x)$  is subject to the two-dimensional wave equation

$$\partial_t [t \partial_t Z(t, x)] = t \partial_x^2 Z(t, x). \tag{6.20}$$

The conformal factor can be determined by straightforward integration from the first-order equations

$$\lambda^{-1} \partial_t \lambda = \frac{t}{4} [(\partial_t Z)^2 + (\partial_x Z)^2], \quad \lambda^{-1} \partial_x \lambda = \frac{t}{2} \partial_t Z \partial_x Z \tag{6.21}$$

<sup>14</sup> See also [36] for a discussion of such solutions in the framework of gravitational solitons.

which are compatible if (6.20) is satisfied. They give rise to the second-order evolution equation for  $\lambda$

$$\partial_t(t\lambda^{-1}\partial_t\lambda) - t\partial_x(\lambda^{-1}\partial_x\lambda) = \frac{t}{2}(\partial_x Z)^2. \tag{6.22}$$

The general solution of (6.20) can be written as<sup>15</sup>

$$Z(t, x) = \int_{-\infty}^{\infty} G(t, x - w)\tilde{q}(w) dw + \int_{-\infty}^{\infty} H(t, x - w)p(w) dw \tag{6.23}$$

with the Green's functions

$$G(t, x) := \frac{1}{\pi} \int_0^{\infty} \cos(kx)J_0(kt) dk, \quad H(t, x) := \frac{1}{\pi} \int_0^{\infty} \cos(kx)Y_0(kt) dk, \tag{6.24}$$

where  $J_0(z)$  and  $Y_0(z) = J_0(z) \ln(z) + \dots$  are the standard Bessel functions (see e.g. [38]). Note that the second integral in (6.24) is well defined (as a distribution) for all  $t > 0$  despite the logarithmic singularity of the integrand. Near the singular hypersurface  $t = 0$ , the general solution admits the expansion [16, 17]

$$Z(t, x) = q(x) + p(x) \ln t + F(t, x) \tag{6.25}$$

where  $q(x) \neq \tilde{q}(x)$  unless  $p(x) = 0$ , and  $\lim_{t \rightarrow 0} F(t, x) = 0$ . The conformal factor expands as

$$\ln \lambda(t, x) = \frac{1}{4}p^2(x) \ln t + \dots \tag{6.26}$$

The functions

$$q(x) = q_0 + q_1x + q_2x^2 + \dots, \quad p(x) = p_0 + p_1x + p_2x^2 + \dots \tag{6.27}$$

can be viewed as the *initial values of the coordinates and momenta on the 'big bang' hypersurface  $t = 0$* .  $p(x)$  correspond to the canonical momenta (rather than the velocities) since we know from section 5.2 that the Lagrange density contains the kinetic term  $\mathcal{L} = \rho P_t P_t$ . With  $\rho(t) = t$  the conjugate momenta are therefore

$$\Pi(t, x) = t\partial_t Z(t, x) \xrightarrow{t \rightarrow 0} p(x). \tag{6.28}$$

We emphasize that we are here working locally in a fixed coordinate chart—the Taylor expansion could be equivalently performed about any other spatial point  $x_0$ .

Evidently, the *Kasner solution* corresponds to constant  $q(x) = q_0$  and  $p(x) = p_0$  in (6.25) and (6.26). In this case we have a direct correspondence between this particular solution of Einstein's equations, and the corresponding solution (6.12) of the  $\mathfrak{H}/K(\mathfrak{H})$  model with  $p_m = q_m = 0$  for  $m \geq 1$ . In view of (6.17) and (6.18), different orbits under  $\mathfrak{H}$  are thus labelled by  $p_0$ . If  $p(x) \equiv 0$ , the solution is no longer of Kasner type; instead of a singularity, it may exhibit other features such as Cauchy horizons [16]. In the following we will not consider this case.

When the higher modes (and the higher order terms in (6.25)) are switched on, the relation is less transparent. *We now define an explicit one-to-one map between the null geodesic trajectories  $\mathcal{V}(t)$  on  $\mathfrak{H}/K(\mathfrak{H})$  characterized by (6.12), and the solutions of (6.20) characterized by the initial values (6.27), by associating the solutions with the same initial*

<sup>15</sup> For *regular* initial data, i.e.  $p(x) = 0$ , this formula can be written in the completely explicit (and manifestly causal) form [37]

$$Z(t, x) = \frac{1}{\pi} \int_{x-t}^{x+t} \frac{q(w) dw}{\sqrt{t^2 - (x-w)^2}}$$

which also follows directly from (6.24) (we thank V Moncrief for a discussion on this point).

data  $q_0, q_1, \dots$  and  $p_0, p_1, \dots$  and  $\sigma_0$ . We emphasize again that such an association makes sense *only after* gauges have been fixed; in particular, all possible coordinate transformations in  $(t, x)$  are ‘used up’ when the four-dimensional line element is cast into the form (6.19), and the diffeomorphism constraint is imposed on the initial data by solving (6.21). The action of the ‘solution generating group’  $\mathfrak{H}$  on the solutions of (6.20) is likewise *defined* by the corresponding action on the solution  $\mathcal{V}(t)$  in (6.16). We next examine how this action compares to the standard realization of the Geroch group on such solutions.

### 6.3. Geroch versus Heisenberg

To elucidate the relation between the transformations (6.17) on the initial data (6.27) on the one hand, and the action of the corresponding subgroup of the Geroch group on solutions of (6.20) on the other, we briefly recall how the latter is usually realized. The integrability of general relativity reduced to  $(1+1)$  dimensions [39, 40] is usually expressed by means of a linear system (or Lax pair) whose integrability condition is equivalent to the reduced Einstein equations [5]. In the standard formulation this is achieved by promoting the matrix  $V(t, x)$  appearing in (5.13) of section 5 to an element of the associated *loop group* by introducing an extra dependence on a spectral parameter  $\gamma$ , namely

$$V(t, x) \rightarrow \hat{V}(t, x; \gamma). \quad (6.29)$$

This matrix satisfies the linear system (Lax pair) equations [5, 6]

$$\partial_\mu \hat{V}^{-1} = Q_\mu + \frac{1 + \gamma^2}{1 - \gamma^2} P_\mu + \frac{2\gamma}{1 - \gamma^2} \epsilon_{\mu\nu} P^\nu. \quad (6.30)$$

The parameter  $\gamma$  is to be interpreted as the spectral parameter of the loop algebra over  $\mathfrak{sl}_{n-2}$ . The integrability of (6.30) implies the equations of motion (5.16) if

$$\gamma = \gamma(t, x; w) = \frac{(x - w) \pm \sqrt{(x - w)^2 - t^2}}{t} \quad \Leftrightarrow \quad w = -\frac{t}{2} \left( \gamma + \frac{1}{\gamma} \right) + x, \quad (6.31)$$

where  $w$  is sometimes called the ‘constant spectral parameter’. The appearance of *two* spectral parameters  $\gamma$  and  $w$ , one of which depends on the spacetime coordinates, is a consequence of the coupling of the  $\sigma$ -model to gravity, and the characteristic feature which distinguishes this model from flat space  $\sigma$ -models. It is precisely the coordinate dependence of the spectral parameter  $\gamma$  which allows the Geroch group to generate space-dependent solutions, as we will now explain for the polarized Gowdy cosmologies.

Diagonal solutions can be obtained by starting from the following  $w$ -dependent matrix<sup>16</sup> considered as an element of the loop group  $\widehat{GL}(1) \subset A_1^{(1)}$

$$\hat{V}(w) = \begin{pmatrix} \exp(G(w)) & 0 \\ 0 & \exp(-G(w)) \end{pmatrix}. \quad (6.32)$$

To generate spacetime-dependent solutions from this matrix, one follows the general procedure of [5] by first expressing  $w$  as a function of  $t, x$  and  $\gamma$  by virtue of (6.31), and then removing the singularity at  $\gamma = 0$  by a compensating transformation. The corresponding solution of the field equations is then obtained from  $\hat{V}$  by setting  $\gamma = 0$ , as  $\hat{V}$  is holomorphic at  $\gamma = 0$  after removal of the poles. In infinitesimal form, these steps are summarized in the combined transformation

$$\delta \hat{V}(t, x; \gamma) = \delta h(t, x; \gamma) \hat{V}(t, x; \gamma) - \hat{V}(t, x; \gamma) \delta g(w), \quad (6.33)$$

<sup>16</sup> The following considerations are based on unpublished work with Damour [41].



where  $\delta g(w) \in \widehat{\mathfrak{gl}}_1$ , and  $\delta h$  is the compensating transformation in  $K(\widehat{\mathfrak{gl}}_1)$  which removes the poles at  $\gamma = 0$ . Equation (6.33) shows that *only half of*  $\widehat{\mathfrak{gl}}_1$  has an actual effect on the solution  $V(t, x)$ , namely those  $\delta g(w) \propto w^n$  with  $n > 0$ . The other half ( $n < 0$ ) yields expressions in  $\delta \hat{V}$  for which  $\delta V = \lim_{\gamma \rightarrow 0} \delta \hat{V}(\gamma) = 0$ . Hence, the latter transformations merely shift the integration constants arising in the definition of the higher order dual potentials, and have no effect on the physical solution.

All solutions of the type (6.25) can be generated from [41]

$$G(w) = f(w) + g(w) \ln w \tag{6.34}$$

with regular functions  $f(w)$  and  $g(w)$  (which are directly related to, but not identical with, the functions  $q(w)$  and  $p(w)$  in (6.27)). The presence of a  $\ln w$  term in (6.34), which is necessary to obtain a non-trivial Kasner coefficient  $p(x)$  in (6.25), signals that we are not dealing with the standard loop group in  $w$ . The necessity of the  $\ln w$  term is another manifestation of the fact that the standard realization of the Geroch group affects only one-half of the initial data, and must therefore be ‘amended’.

From (6.31) and (6.34), we can deduce the leading contributions to  $\delta Z(t, x)$  for  $t \sim 0$  from a given  $\delta g(w)$ , following the steps described above, with the result

$$\begin{aligned} \delta g(w) = w^n &\Rightarrow \delta Z(t, x) \propto x^n, \\ \delta g(w) = w^n \ln w &\Rightarrow \delta Z(t, x) \propto x^n \ln t, \end{aligned} \tag{6.35}$$

for  $n \geq 0$ . This shows that a regular (non-logarithmic) affine level  $n$  element of the Geroch group induces a spatial dependence  $\propto x^n$  in the regular initial data in a Taylor expansion around  $x = 0$ , but that a logarithmic dependence on the loop parameter  $w$  is required to change the singular initial data (the Kasner coefficient function  $p(x)$ ). The action of the Witt–Virasoro algebra, and hence that of infinitesimal variations along  $\hat{d}$ , were discussed in [42]. The central charge acts on the conformal factor  $\lambda$  by scaling transformations [4–6], in agreement with the results derived at the end of section 6.1.

Equations (6.35) illustrate the main advantage of the affine coset model and our new realization of (a subgroup of) the Geroch group in comparison with the linear system approach: in the Heisenberg model, the *full* algebra generates non-trivial transformations on the initial data—the positive half shifts the initial coordinates  $q_m$ , and the negative half shifts the initial momenta  $p_m$ , see (6.17). By contrast, for the standard realization of the Geroch group only *half* the transformations  $\propto w^n$  for  $n > 0$  act non-trivially and change the initial coordinates encoded in  $q(x)$ ; in this sector, the action of the Geroch group agrees with the action of the upper half of  $\mathfrak{H}$ . On the other hand, in the standard approach, we must extend the Geroch transformations by allowing  $w^n \ln w$  terms in order to shift the initial momenta  $p(x)$ . Moreover, this rather *ad hoc* extension of the loop group fails at the nonlinear level, i.e. for non-diagonal metrics, because it would force us to admit *arbitrary* positive powers of  $\ln w$  [41]. This difficulty is avoided altogether in our new coset approach, where the ‘solution generating group’ acts on *all* initial data, except  $p_0$ , and which furthermore does not require modifying the loop group by logarithmic terms to generate the most general solution.

The difference between the two realizations of the affine symmetry is also evident from the way the (dynamical) fields are encoded in  $\hat{V}$  and  $\mathcal{V}$ , respectively. Since the Chevalley involution acts by [5]

$$\omega(\hat{V}(t, x; \gamma)) = (\hat{V}^T)^{-1} \left( t, x; \frac{1}{\gamma} \right) \tag{6.36}$$

the rhs of (6.30) is invariant, whence

$$\partial_\mu \hat{V}^{-1} \in K(A_{n-2}^{(1)}). \quad (6.37)$$

By contrast, for the geodesic  $\sigma$ -model of section 4, all the dynamics is contained in the coset components, see (4.18). Hence, we conclude again that the linear system (6.30) is not immediately suitable for the comparison with the results of the foregoing sections.

## 7. Discussion

There is an alternative formulation of the Lax pair for two-dimensional gravity [43], which is somewhat more similar to (4.25), in that its rhs belongs to a coset subalgebra, and involves the CSA generators  $\hat{c}$  and  $\hat{d}$  explicitly. In that formulation the Cartan form reads (using light-cone coordinates)

$$\partial_\pm \mathcal{V}^{-1}(t, x) = \pm \rho^{-1} \partial_\pm \rho (L_0 - L_{\pm 1}) + \frac{1}{2} Q_{\pm\alpha\beta} J^{\alpha\beta} + \frac{1}{2} P_{\pm\alpha\beta} K_{\pm 1\beta}^\alpha \mp \partial_\pm \sigma \hat{c}. \quad (7.1)$$

Here,  $L_0$  and  $L_{\pm 1}$  belong to the Möbius subalgebra of the Virasoro algebra (which can be embedded in the enveloping algebra of the affine algebra with  $L_0 = \hat{d}$ ). In contradistinction to the linear system (6.30), the expansion of (7.1) is truncated at affine level one, but involves the Möbius generators. (This has an effect on the set of allowed dressing transformations [43].)

In the above linear system, the generator  $L_{-1}$  acts like a derivative operator on a vertex representation, and belongs to a Witt–Virasoro algebra acting on the affine algebra via a semi-direct product. This suggests that there might exist a similar, but larger, subalgebra inside the enveloping algebra of  $AE_n$ , which would contain derivative operators in all spatial directions, and contain the standard Witt–Virasoro algebra as a subalgebra. In addition, the affine algebra might be embedded in a *toric algebra* (see e.g. [44] and references therein). Such an algebra is spanned by generators  $T_{\vec{m}}^A$  (where  $\vec{m}$  designates a vector in some vector space), with commutation relations of the form

$$[T_{\vec{m}}^A, T_{\vec{n}}^B] = f^{AB}{}_C T_{\vec{m}+\vec{n}}^C + \text{central terms}. \quad (7.2)$$

However, looking at the relation for the gradient fields derived for  $AE_3$  in section 3, we see that the naive identification of  $\vec{m}$  with the 3-vector of the multiplicities of the indices 1, 2, 3 among the ‘gradient indices’  $a_1, \dots, a_{\ell-1}$  of the generator  $E_{a_1, \dots, a_{\ell-1}}^{b_1 b_2}$  will fail since the commutator (3.24) will always *add* one index and so destroy the vector space structure on such  $\vec{m}$ . Moreover, we have seen in section 3 that the gradient representations by themselves do not close into a subalgebra of  $AE_3$ . We thus conclude that there is no toric algebra inside  $AE_n$  or its enveloping algebra.

The interpretation of the affine coset proposed in section 6 provides a more favourable realization of the Geroch group since it can incorporate transformations of both the initial coordinates and the initial velocities in the general case without extending the set of allowed transformations. We anticipate that similar results hold for the  $\sigma$ -model realization of the affine group  $A_1^{(1)}$  in comparison with the standard realization of the full Geroch group. However, the more important challenge at this point is now to find out how the hyperbolic  $AE_3$  coset model can generate the most general dependence on four-dimensional space and time coordinates, and to understand the significance of the fact that, unlike the affine elements, the gradient generators no longer form a closed subalgebra of  $AE_n$ .

It is also interesting to consider higher levels of the affine coset model by including as a next step the basic representation [26] of the affine algebra. This will show new aspects of the correspondence characteristic for the KMA  $\sigma$ -model crucial for the programme of [9].

**Acknowledgments**

We are grateful to T Damour, F Englert, V Moncrief and A Rendall for enlightening discussions, to D Bernard and T Fischbacher for correspondence and to T Damour for permission to include some unpublished results of [41].

**Appendix A. The compact subalgebra is not a KMA**

In this appendix we prove that the ‘maximal compact’ invariant subalgebra of an infinite-dimensional KMA is *not* of Kac–Moody type. This applies in particular to the algebras  $AE_n$  and  $E_{10}$ , and their invariant subalgebras.

**Proposition 1.** *Let  $\mathfrak{g}$  be any infinite-dimensional split real Kac–Moody algebra  $\mathfrak{g}$ , which is not the direct sum of (infinitely many) finite-dimensional algebras. Then the infinite-dimensional subalgebra  $\mathfrak{k} \subset \mathfrak{g}$  fixed by the Chevalley involution  $\omega$  is not a Kac–Moody algebra.*

**Proof.** The main observation is that the contravariant Hermitian form on the Kac–Moody algebra

$$(x|y) := -\langle x|\omega(y) \rangle \quad \text{for } x, y, \in \mathfrak{g} \tag{A.1}$$

is not positive definite everywhere on the split real KMA if the latter is infinite dimensional [24]: in the CSA, and only in the CSA, there exist elements of both positive and negative norm squared, because the Cartan matrix is not positive definite. However, since the CSA is not invariant under  $\omega$ , the compact subalgebra  $\mathfrak{k}$  does not contain any such elements, and the contravariant form on  $\mathfrak{k}$  (inherited from  $\mathfrak{g}$ ) is therefore positive definite on  $\mathfrak{k}$  [45]. Because  $\mathfrak{k}$  is infinite dimensional, it can thus be a KMA if and only if it is the (orthogonal) sum of infinitely many finite-dimensional algebras of KM type.

We will prove that this cannot happen by exhibiting a contradiction if the assumption were true. Therefore, assume that  $\mathfrak{k}$  is the infinite direct sum of some finite-dimensional algebras. Now consider some (regular) level decomposition  $\mathfrak{g} = \bigoplus_{\ell \in \mathbb{Z}} \mathfrak{g}_\ell$  of  $\mathfrak{g}$  such that  $0 < \dim \mathfrak{g}_\ell < \infty$  for all  $\ell \in \mathbb{Z}$ . Then  $\mathfrak{g}$  is generated by taking multiple commutators of  $\mathfrak{g}_{-1}$ ,  $\mathfrak{g}_0$  and  $\mathfrak{g}_1$ . Since the Chevalley involution  $\omega$  acts according to  $\omega(\mathfrak{g}_\ell) = \mathfrak{g}_{-\ell}$ , the invariant subalgebra admits the decomposition (not a grading!)

$$\mathfrak{k} = \bigoplus_{\ell \geq 0} \mathfrak{k}_\ell \quad \text{with } \mathfrak{k}_0 \subset \mathfrak{g}_0, \quad \mathfrak{k}_\ell \subset \mathfrak{g}_{-\ell} \oplus \mathfrak{g}_\ell \quad \Rightarrow \quad [\mathfrak{k}_0, \mathfrak{k}_\ell] \subset \mathfrak{k}_\ell. \tag{A.2}$$

Like  $\mathfrak{g}$ , the infinite-dimensional invariant subalgebra  $\mathfrak{k}$  is generated by taking multiple commutators of  $\mathfrak{k}_0$  and  $\mathfrak{k}_1$ . Since all  $\mathfrak{g}_\ell$  are of finite dimension, it follows in particular that  $\mathfrak{k}_0$  and  $\mathfrak{k}_1$  are also finite dimensional. According to our assumption they must therefore belong to a *finite* sum of finite-dimensional subalgebras of  $\mathfrak{k}$ . Hence, the algebra they generate is also finite dimensional, in contradiction with the fact that  $\mathfrak{k}$  is infinite dimensional.  $\square$

The proposition unfortunately does not give an answer to the important question what kind of Lie algebra  $\mathfrak{k}$  is, and, more pressingly, what its representation theory is. It is expected that the representation theory of  $\mathfrak{k}$  is associated with the supersymmetric extension of the one-dimensional  $\sigma$ -model [13, 46].

## Appendix B. $AE_n$ $\sigma$ -model in $\mathfrak{sl}_n$ decomposition and $p$ -forms

In this appendix, we analyse the  $\sigma$ -model based on  $AE_n$  of section 4 from its  $\mathfrak{sl}_n$  subalgebra. We will also add  $p$ -forms to the Einstein equation of the model to make contact with other oxidized theories.

### B.1. Pure gravity

We first study the  $\sigma$ -model and then compare it to the reduction of the Einstein equation.

The Cartan form in triangular gauge in  $\mathfrak{sl}_n$  form now is

$$\partial_t \mathcal{V} \mathcal{V}^{-1} = P_{ab} S^{ab} + Q_{ab} J^{ab} + \frac{1}{(n-2)!} P_{a_1 \dots a_{n-2}, a_{n-1}} E^{a_1 \dots a_{n-2}, a_{n-1}} + \dots \quad (\text{B.1})$$

with indices  $a, b$  taking values  $1, \dots, n$ . The resulting  $\sigma$ -model equation on level  $\ell = 0$ , in the truncation to  $|\ell| \leq 1$ , can be deduced from the results of section 2.2, with the result

$$\begin{aligned} n \partial_t (n^{-1} P_{ab}) &= 2 P_{c(a} Q_{b)c} + \frac{1}{2(n-2)!} (\delta_{ab} P_{c_1 \dots c_{n-2}, c_{n-1}} P_{c_1 \dots c_{n-2}, c_{n-1}} \\ &\quad - P_{c_1 \dots c_{n-2}, a} P_{c_1 \dots c_{n-2}, b} - (n-2) P_{ac_1 \dots c_{n-3}, c_{n-2}} P_{bc_1 \dots c_{n-3}, c_{n-2}}). \end{aligned} \quad (\text{B.2})$$

In the corresponding gravity theory, the space–space components of reduced Einstein equations in flat components are

$$R_{ab} = R_{ab}^{\text{temp}} + R_{ab}^{\text{spat}} = 0, \quad (\text{B.3})$$

where we defined, following [14],

$$R_{ab}^{\text{temp}} = \partial_0 \omega_{ab0} + \omega_{c0} \omega_{ab0} - 2 \omega_{0c(a} \omega_{b)c0}, \quad (\text{B.4})$$

$$R_{ab}^{\text{spat}} = \frac{1}{4} \tilde{\Omega}_{cda} \tilde{\Omega}_{cdb} - \frac{1}{2} \tilde{\Omega}_{acd} \tilde{\Omega}_{bcd} - \frac{1}{2} \tilde{\Omega}_{acd} \tilde{\Omega}_{bdc} - \frac{1}{2} \partial_c \tilde{\Omega}_{c(ab)}. \quad (\text{B.5})$$

Here,  $\tilde{\Omega}_{abc}$  is the tracefree part of the anholonomy  $\Omega_{abc} = 2e_{[a}^m e_b]^n \partial_m e_{nc}$  and  $\omega_{abc} = (\Omega_{abc} + \Omega_{cab} - \Omega_{bca})/2$  is the spin connection. We use the same decomposition (5.1) of the vielbein as in section 5.1.

By comparing equation (B.2) with equation (B.3), we see that almost all terms match upon identifying

$$n(t) \equiv N e^{-1}(t, x_0), \quad (\text{B.6})$$

$$P_{ab}(t) \equiv \omega_{abt}(t, x_0), \quad (\text{B.7})$$

$$Q_{ab}(t) \equiv -\omega_{tab}(t, x_0), \quad (\text{B.8})$$

$$P_{a_1 \dots a_{n-2}, a_{n-1}}(t) \equiv \frac{N}{2} \varepsilon_{a_1 \dots a_{n-2} bc} \tilde{\Omega}_{bc a_{n-1}}(t, x_0) \quad (\text{B.9})$$

for some fixed spatial point  $x_0$ . The terms which do not match are the second spatial derivatives and the cross-term  $\tilde{\Omega}_{acd} \tilde{\Omega}_{bdc}$  in (B.5). However, the second spatial derivatives are not expected to match in this approximation and the cross-term disappears upon using a conformal gauge for the zweibein.

## B.2. Inclusion of $p$ -forms

We now add a single  $p$ -form field to the Einstein action via their kinetic terms<sup>17</sup>

$$S = \int d^{n+1}x \det(E) \left( R - \frac{1}{2(p+1)!} F_{M_1 \dots M_{p+1}} F^{M_1 \dots M_{p+1}} \right). \quad (\text{B.10})$$

The generalization to several  $p$ -forms is straightforward and we do not consider dilaton terms here for simplicity.

The Einstein equation resulting from (B.10) can be written as

$$R_{MN} = \frac{1}{2p!} F_{M P_1 \dots P_p} F_N{}^{P_1 \dots P_p} - \frac{p}{2(p+1)!(n-1)} G_{MN} F_{P_1 \dots P_{p+1}} F^{P_1 \dots P_{p+1}}. \quad (\text{B.11})$$

Converting into flat indices and restricting to the spatial directions, we write

$$R_{ab}^{\text{temp}} + R_{ab}^{\text{spat}} = -N^{-2} T_{ab}^{\text{el}} + T_{ab}^{\text{magn}}, \quad (\text{B.12})$$

with the energy–momentum tensor split into ‘electric’ and ‘magnetic’ contributions according to

$$T_{ab}^{\text{el}} = \frac{p}{2p!} F_{t a c_1 \dots c_{p-1}} F_{t b c_1 \dots c_{p-1}} - \frac{p}{2p!(n-1)} \delta_{ab} F_{t c_1 \dots c_p} F_{t c_1 \dots c_p}, \quad (\text{B.13})$$

$$T_{ab}^{\text{magn}} = \frac{1}{2p!} F_{a c_1 \dots c_p} F_{b c_1 \dots c_p} - \frac{p}{2(p+1)!(n-1)} \delta_{ab} F_{c_1 \dots c_{p+1}} F_{c_1 \dots c_{p+1}}. \quad (\text{B.14})$$

The factor  $-N^{-2}$  in (B.12) stems from lowering the  $t$  index. Similarly, we split the Ricci tensor into temporal and spatial derivatives as before.

On the algebra side, we need to include additional generators to account for the electric and magnetic contributions to the Einstein equation. Consider the expansion of the Cartan form to be

$$\begin{aligned} \partial_t \mathcal{V} \mathcal{V}^{-1} &= P_{ab} S^{ab} + Q_{ab} J^{ab} + \frac{1}{(n-2)!} P_{a_1 \dots a_{n-2}, a_{n-1}} E^{a_1 \dots a_{n-2}, a_{n-1}} \\ &+ \frac{1}{p!} P_{a_1 \dots a_p}^{\text{el}} E^{a_1 \dots a_p} + \frac{1}{(n-p-1)!} P_{a_1 \dots a_{n-p-1}}^{\text{magn}} E^{a_1 \dots a_{n-p-1}} + \dots \end{aligned} \quad (\text{B.15})$$

In the first line of (B.15), we have the contribution of the  $AE_n$  fields as above. The second line now belongs to additional generators we have introduced into the algebra. That such generators are present in the relevant Kac–Moody algebra for the oxidized theory follows from the results of [47]. Demanding the normalizations

$$\langle E^{a_1 \dots a_p} | F_{b_1 \dots b_p} \rangle = p! \delta_{b_1 \dots b_p}^{a_1 \dots a_p}, \quad (\text{B.16})$$

$$\langle E^{a_1 \dots a_{n-p-1}} | F_{b_1 \dots b_{n-p-1}} \rangle = (n-p-1)! \delta_{b_1 \dots b_{n-p-1}}^{a_1 \dots a_{n-p-1}}, \quad (\text{B.17})$$

leads to the commutators of the new generators

$$[K^a{}_b, E^{c_1 \dots c_p}] = (-1)^p p \delta_b^{[c_1} E^{c_2 \dots c_p] a}, \quad (\text{B.18})$$

$$[E^{a_1 \dots a_p}, F_{b_1 \dots b_p}] = -\frac{p}{n-1} p! \delta_{b_1 \dots b_p}^{a_1 \dots a_p} K + p \cdot p! \delta_{[b_1 \dots b_{p-1}}^{[a_1 \dots a_{p-1}} K^{a_p] b_p]}, \quad (\text{B.19})$$

<sup>17</sup> We need not concern ourselves with possible Chern–Simons-like terms since these do not affect the Einstein equation. However, they are crucial for the gauge field equation of motions and Bianchi identities.

and similar expressions for the magnetic generator with  $p$  replaced by  $(n - p - 1)$ . From this, together with the commutator (2.16), we can deduce the following  $\ell = 0$  equation of motion for the  $\sigma$ -model in this approximation

$$\begin{aligned}
 n\partial_t(n^{-1}P_{ab}) = & 2P_{c(a}Q_{b)c} + \frac{1}{2(n-2)!}(\delta_{ab}P_{c_1\dots c_{n-2},c_{n-1}}P_{c_1\dots c_{n-2},c_{n-1}} \\
 & - P_{c_1\dots c_{n-2},a}P_{c_1\dots c_{n-2},b} - (n-2)P_{ac_1\dots c_{n-3},c_{n-2}}P_{bc_1\dots c_{n-3},c_{n-2}}) \\
 & + \frac{p}{2p!}P_{ac_1\dots c_{p-1}}^{\text{el}}P_{bc_1\dots c_{p-1}}^{\text{el}} - \frac{p}{2p!(n-1)}\delta_{ab}P_{c_1\dots c_p}^{\text{el}}P_{c_1\dots c_p}^{\text{el}} \\
 & + \frac{n-p-1}{2(n-p-1)!}P_{ac_1\dots c_{n-p-2}}^{\text{magn}}P_{bc_1\dots c_{n-p-2}}^{\text{magn}} - \frac{p}{2p!(n-1)}\delta_{ab}P_{c_1\dots c_{n-p-1}}^{\text{magn}}P_{c_1\dots c_{n-p-1}}^{\text{magn}}.
 \end{aligned}
 \tag{B.20}$$

By comparing (B.12) and (B.20) we find agreement if we identify the new terms by

$$P_{a_1\dots a_p}^{\text{el}}(t) \equiv F_{ta_1\dots a_p}(t, x_0), \tag{B.21}$$

$$P_{a_1\dots a_{n-p-1}}^{\text{magn}}(t) \equiv \frac{N}{(p+1)!}\varepsilon_{a_1\dots a_{n-p-1}b_1\dots b_{p+1}}F_{b_1\dots b_{p+1}}(t, x_0). \tag{B.22}$$

in addition to (B.6). There are sign ambiguities here, since the fields appear quadratically in (B.20). These will be fixed from the equations of motion and Bianchi identities for the  $p$ -form fields.

## References

- [1] Geroch R 1971 A method for generating solutions of Einstein's equations *J. Math. Phys.* **12** 918–24
- [2] Geroch R 1972 A method for generating solutions of Einstein's equations. II *J. Math. Phys.* **13** 394–404
- [3] Hoenselaers C and Dietz W (ed) 1984 *Solutions of Einstein's Equations: Techniques and Results* (Berlin: Springer)
- [4] Stephani H, Kramer D, MacCallum M, Honselaers C and Herlt E 2003 *Exact Solutions of Einstein's Field Equations* (Cambridge: Cambridge University Press)
- [5] Julia B 1981 Group disintegrations *Superspace and Supergravity: Proc. Nuffield Workshop (Cambridge, UK, 22 June–12 July 1980)* ed S W Hawking and M Roček (Cambridge: Cambridge University Press) pp 331–50 (LPTENS 80/16)
- [6] Julia B 1985 Kac–Moody symmetry of gravitation and supergravity theories *Applications of Group Theory in Physics and Mathematical Physics (Lectures in Applied Mathematics vol 21)* ed M Flato, P Sally and G Zuckerman (Providence, RI: American Mathematical Society) pp 355–74 (LPTENS 82/22)
- [7] Breitenlohner P and Maison D 1987 On the Geroch group *Ann. Poincaré Phys. Theor.* **46** 215–46
- [8] Nicolai H 1991 Two-dimensional gravities and supergravities as integrable system *Lectures presented at 30th Schladming Winter School (Schladming 27 Feb–5 Mar 1991)* (DESY-91-038)
- [9] Damour T, Henneaux M, Julia B and Nicolai H 2001 Hyperbolic Kac–Moody algebras and chaos in Kaluza–Klein models *Phys. Lett. B* **509** 323–30 (Preprint hep-th/0103094)
- [10] Nicolai H 1992 A hyperbolic Lie algebra from supergravity *Phys. Lett. B* **276** 333
- [11] Damour T, Henneaux M and Nicolai H 2002  $E_{10}$  and a 'small tension expansion' of M-theory *Phys. Rev. Lett.* **89** 221601 (Preprint hep-th/0207267)
- [12] Damour T, Henneaux M and Nicolai H 2003 Cosmological billiards *Class. Quantum Grav.* **20** R145–200 (Preprint hep-th/0212256)
- [13] Nicolai H and Fischbacher T 2003 Low level representations of  $E_{10}$  and  $E_{11}$  *Contribution to the Proc. Ramanujan Int. Symposium on Kac–Moody Algebras and Applications, ISKMAA-2002 (Chennai 28–31 Jan.)* (Preprint hep-th/0301017)
- [14] Fischbacher T 2005 The structure of  $E_{10}$  at higher  $A_9$  levels: a first algorithmic approach *J. High Energy Phys.* JHEP08(2005)012 (Preprint hep-th/0504230)
- [15] Kleinschmidt A and Nicolai H 2004  $E_{10}$  and  $SO(9, 9)$  invariant supergravity *J. High Energy Phys.* JHEP07(2004)041 (Preprint hep-th/0407101)

- [14] Damour T and Nicolai H 2004 Eleven dimensional supergravity and the  $E_{10}/K(E_{10})\sigma$ - model at low  $A_9$  levels *Preprint* hep-th/0410245
- [15] Kleinschmidt A and Nicolai H 2005 IIB supergravity and  $E_{10}$  *Phys. Lett. B* **606** 391–402 (*Preprint* hep-th/0411225)
- [16] Chrusciel P, Isenberg J and Moncrief V 1990 Strong cosmic censorship in polarised Gowdy spacetimes *Class. Quantum Grav.* **7** 1671–80
- [17] Kichenassamy S and Rendall A 1998 Analytic description of singularities in Gowdy spacetimes *Class. Quantum Grav.* **15** 1339–55
- [18] West P C 2001  $E_{11}$  and M theory *Class. Quantum Grav.* **18** 4443–60 (*Preprint* hep-th/0104081)
- [19] Englert F, Houart L, Taormina A and West P 2003 The symmetry of M theories *J. High Energy Phys.* JHEP09(2003)020 (*Preprint* hep-th/0304206)
- [20] West P 2003  $E_{11}$ ,  $SL(32)$  and central charges *Phys. Lett. B* **575** 333–42 (*Preprint* hep-th/0307098)
- [21] Englert F and Houart L 2004  $G^{+++}$  invariant formulation of gravity and M-theories: exact BPS solutions *J. High Energy Phys.* JHEP01(2004)002 (*Preprint* hep-th/0311255)
- [22] Kleinschmidt A and West P 2004 Representations of  $G^{+++}$  and the role of space-time *J. High Energy Phys.* JHEP02(2004)033 (*Preprint* hep-th/0312247)
- [23] West P C 2000 Hidden superconformal symmetry in M theory *J. High Energy Phys.* JHEP08(2000)007 (*Preprint* hep-th/0005270)
- [24] Kac V G 1990 *Infinite Dimensional Lie Algebras* 3rd edn (Cambridge: Cambridge University Press)
- [25] Fischbacher T private communication
- [26] Feingold A J and Frenkel I B 1983 A hyperbolic Kac–Moody algebra and the theory of Siegel modular forms of genus 2 *Math. Ann.* **263** 87–144
- [27] Curtright T 1985 Generalized gauge fields *Phys. Lett. B* **165** 304
- [28] Obers N A and Pioline B 1999 U-duality and M-theory *Phys. Rep.* **318** 113 (*Preprint* hep-th/9809039)
- [29] Hull C M 2001 Duality in gravity and higher spin gauge fields *J. High Energy Phys.* JHEP09(2001)027 (*Preprint* hep-th/0107149)
- [30] Bekaert X, Boulanger N and Henneaux M 2003 Consistent deformations of dual formulations of linearized gravity: A no-go result *Phys. Rev. D* **67** 044010 (*Preprint* hep-th/0210278)
- [31] West P 2003 Very extended  $E_8$  and  $A_8$  at low levels, gravity and supergravity *Class. Quantum Grav.* **20** 2393 (*Preprint* hep-th/0212291)
- [32] Nieto J A 1999 S-duality for linearized gravity *Phys. Lett. A* **262** 274 (*Preprint* hep-th/9910049)
- [33] Henneaux M and Teitelboim C 2005 Duality in linearized gravity *Phys. Rev. D* **71** 024018 (*Preprint* gr-qc/0408101)
- [34] Damour T and Nicolai H 2005 Higher order M theory corrections and the Kac–Moody algebra  $E_{10}$  *Class. Quantum Grav.* **22** 2849–79 (*Preprint* hep-th/0504153)
- [35] Nagar A and Nicolai H 2001 Infinite-dimensional symmetries in gravity *Gravitational Waves* ed I Ciufolini *et al* (Bristol: Institute of Physics Publishing)
- [36] Belinski V and Verdager E 2001 *Gravitational Solitons* (Cambridge: Cambridge University Press)
- [37] Courant R and Hilbert D 1989 *Methods of Mathematical Physics* vol II (New York: Wiley)
- [38] Temme N M 1996 *Special Functions: an Introduction to the Classical Functions of Mathematical Physics* (New York: Wiley)
- Jeffrey A 1995 *Handbook of Mathematical Formulas and Integrals* (New York: Academic)
- [39] Maison D 1978 Are the stationary, axially symmetric Einstein equations completely integrable? *Phys. Rev. Lett.* **41** 521
- [40] Belinski V and Zakharov V 1978 Integration of Einstein’s equations by means of inverse scattering problem technique and construction of exact soliton solutions *Sov. Phys.—JETP* **48** 985
- [41] Damour T and Nicolai H unpublished notes
- [42] Julia B and Nicolai H 1996 Conformal internal symmetry of 2d sigma-models coupled to gravity and a dilaton *Nucl. Phys. B* **482** 431 (*Preprint* hep-th/9608082)
- [43] Bernard D and Julia B 1999 Twisted self-duality of dimensionally reduced gravity and vertex operators *Nucl. Phys. B* **547** 427 (*Preprint* hep-th/9712254)
- [44] Eswara Rao S 1995 Iterated loop modules and a filtration for vertex representations of toroidal Lie algebras *Pac. J. Math.* **171** 511–28
- [45] Kac V G and Peterson D H 1984 Unitary structures in representations of infinite-dimensional groups and a convexity theorem *Invent. Math.* **76** 1
- [46] Nicolai H and Samtleben H 2005 On  $K(E_9)$  *Q. J. Pure Appl. Math.* **1** 180 (*Preprint* hep-th/0407055)
- [47] Kleinschmidt A, Schnakenburg I and West P 2004 Very extended Kac–Moody algebras and their interpretation at low levels *Class. Quantum Grav.* **21** 2493–525 (*Preprint* hep-th/0309198)