

The fields of uniformly accelerated charges in de Sitter spacetime

Jiří Bičák* and Pavel Krtouš†

*Institute of Theoretical Physics, Faculty of Mathematics and Physics, Charles University,
V Holešovičkách 2, 180 00 Prague 8, Czech Republic*

(Dated: July 1, 2002)

The scalar and electromagnetic fields of charges uniformly accelerated in de Sitter spacetime are constructed. They represent the generalization of the Born solutions describing fields of two particles with hyperbolic motion in flat spacetime. In the limit $\Lambda \rightarrow 0$, the Born solutions are retrieved. Since in the de Sitter universe the infinities \mathcal{I}^\pm are spacelike, the radiative properties of the fields depend on the way in which a given point of \mathcal{I}^\pm is approached. The fields must involve both retarded and advanced effects: Purely retarded fields do not satisfy the constraints at the past infinity \mathcal{I}^- .

PACS numbers: 04.20.-q, 04.40.-b, 98.80.Hw, 03.50.-z

The question of the electromagnetic field and associated radiation from uniformly accelerated charges has been one of the best known “perpetual problems” in classical physics from the beginning of the past century. In the pioneering work in 1909, Born gave the time-symmetric solution for the field of two point particles with opposite charges, uniformly accelerated in opposite directions in Minkowski space. In the 1920s Sommerfeld, von Laue, Pauli, Schott and others discussed the properties of the field. The controversial point that the field exhibits radiative features but that the radiation reaction force vanishes for the hyperbolic motion, and related questions, was discussed in many articles from 1960s onward. Even the December 2000 issue of *Annals of Physics* contains three papers [1] with numerous references on “electrodynamics of hyperbolically accelerated charges”.

In general relativity, solutions of Einstein’s equations, representing “uniformly accelerated particles or black holes”, are the *only* explicitly known exact *radiative* spacetimes describing *finite* sources. They are asymptotically flat at null infinity [2] (except for some special points) and have been used in gravitational radiation theory, quantum gravity and numerical relativity (cf. review [3]). One of the best known examples is the *C*-metric, describing uniformly accelerated black holes. There exists also the *C*-metric for a nonvanishing cosmological constant Λ . However, no general framework is available to analyze these spacetimes for $\Lambda \neq 0$ as that given in Ref. [2] for $\Lambda = 0$.

In this Letter, we present the generalization of the Born solutions for scalar and electromagnetic fields to the case of two charges uniformly accelerated in de Sitter universe, and explicitly show how in the limit $\Lambda \rightarrow 0$ the Born solutions are retrieved. We also study the asymptotic expansions of the fields in the neighborhood of future infinity \mathcal{I}^+ . In de Sitter spacetime, conformal infinities, \mathcal{I}^\pm , are *spacelike*, which implies the presence of particle and event horizons. It is known [4] that the radiation field is “less invariantly” defined when \mathcal{I}^+ is spacelike (it depends on the direction in which \mathcal{I}^+ is approached), but no explicit model appears to be available

so far. Our solutions can serve as prototypes for studying these issues.

In recent work [5], we analyzed fields of accelerated sources to show the *insufficiency of purely retarded fields in de Sitter spacetime*. Consider a point P near \mathcal{I}^- whose past null cone will not cross the particles’ worldlines (Fig. 1). The field at P should vanish if an incoming field is absent. However, the “Coulomb-type” field of particles cannot vanish there because of Gauss law [6]. The requirement that the field be purely retarded leads, in general, to a bad behavior of the field along the “creation light cone” of the “point” at which a source enters the universe (see Ref. [5] for detailed discussion).

It is natural to use de Sitter space for studying radiating sources in spacetimes which are not asymptotically flat and possess spacelike infinities: It is the space of constant curvature, conformal to Minkowski space, and with Huygens principle (no tails of radiation) satisfied for conformally invariant fields. The de Sitter universe also plays an important role in cosmology — not only in the context of inflationary theories but also as the “asymptotic state” of standard cosmological models with $\Lambda > 0$, which has been indeed suggested by recent observations. In addition, the Born fields generalized to de Sitter space should be relevant from quantum perspectives: for example, for studying particle production in strong fields, or accelerating detectors in the presence of a cosmological horizon.

The de Sitter universe has topology $S^3 \times \mathbb{R}$. The metric in standard “spherical” coordinates¹ $\{\tau, \chi, \vartheta, \varphi\}$ is

$$g_{\text{as}} = -d\tau^2 + \alpha^2 \cosh^2(\tau/\alpha) (d\chi^2 + \sin^2 \chi d\omega^2), \quad (1)$$

¹ It is convenient to allow the angular coordinates χ, ϑ, φ on S^3 to attain values in \mathbb{R} and use the identifications $\chi \cong \chi + 2\pi$, $\vartheta \cong \vartheta + 2\pi$, $\varphi \cong \varphi + 2\pi$, $\{\chi, \vartheta, \varphi\} \cong \{-\chi, \pi - \vartheta, \varphi + \pi\}$, and $\{\chi, \vartheta, \varphi\} \cong \{\chi, -\vartheta, \varphi + \pi\}$. Thus, the “radial” coordinate χ can be negative, the points with $\chi < 0$ being identical to those with $|\chi| > 0$, located “symmetrically” with respect to the origin $\chi = 0$. The same convention is used for \tilde{r}, r, \tilde{r} , and R (see Appendix in [5] for details).




FIG. 1: The conformal diagram of de Sitter spacetime. The lines of coordinates $\{\tilde{t}, \tilde{r}\}$ and $\{t, r\}$ are shown. Uniformly accelerated particles move along worldlines 1 and 1'. The shaded region is the domain of influence of 1, its boundary \mathcal{H} is the “creation light cone” of this particle “born” at $\tilde{t} = 0$ at “point” O . Retarded fields of 1 and 1’ cannot affect point P ; a Coulomb-type field, however, cannot vanish there.

where $d\omega^2 = d\vartheta^2 + \sin^2\vartheta d\varphi^2$, $\tau \in \mathbb{R}$, and $\alpha^2 = 3/\Lambda$. Putting

$$\chi = \tilde{r}, \quad \tau = \alpha \log\left(\tan \frac{\tilde{t}}{2}\right), \quad \tilde{t} \in \langle 0, \pi \rangle$$

in Eq. (1), the de Sitter metric can be written in the form

$$g_{\text{ds}} = \frac{\alpha^2}{\sin^2 \tilde{t}} (-d\tilde{t}^2 + d\tilde{r}^2 + \sin^2 \tilde{r} d\omega^2). \quad (2)$$

The corresponding coordinate lines are drawn in the Penrose diagram of de Sitter space in Fig. 1. The lines $\tilde{r} = \pi$ and $\tilde{r} = -\pi$ are identified, the spacelike hypersurfaces $\tilde{t} = 0$ and $\tilde{t} = \pi$ represent past (\mathcal{I}^-) and future (\mathcal{I}^+) infinities of de Sitter spacetime.

By employing conformal techniques, we recently studied [5] two particles moving with uniform acceleration² in de Sitter space. The worldlines of both uniformly accelerated particles are given by

$$\tan \tilde{t} = -\frac{\cosh \beta_o}{\sinh\left(\frac{\lambda_{\text{ds}}}{\alpha} \cosh \beta_o\right)},$$

$$\tan \tilde{r} = \pm \frac{\sinh \beta_o}{\cosh\left(\frac{\lambda_{\text{ds}}}{\alpha} \cosh \beta_o\right)};$$

they are plotted as 1 and 1’ in Fig. 1. Both particles start at antipodes of the spatial section of de Sitter space at \mathcal{I}^-




FIG. 2: The worldlines 2, 2’ of uniformly accelerated charges symmetrically located with respect to the origins of both de Sitter and conformally related Minkowski spacetimes.

and move one towards the other until $\tilde{t} = \pi/2$, the moment of the maximal contraction of de Sitter space. Then they move, in a time-symmetric manner, apart from each other until they reach future infinity at the antipodes from which they started. Their physical velocities, as measured in the “comoving” coordinates $\{\tau, \chi, \vartheta, \varphi\}$ of Eq. (1), have simple forms

$$v_\chi = \sqrt{g_{\chi\chi}} \frac{d\chi}{d\tau} = \mp \frac{a_o \alpha \tanh(\tau/\alpha)}{\sqrt{1 + a_o^2 \alpha^2 \tanh^2(\tau/\alpha)}},$$

where $|a_o|$ is the magnitude of their acceleration. In contrast to the flat space case, the particles do not approach the velocity of light in the “natural” global coordinate system. They are causally disconnected (Fig. 1) as in the flat space case: No signal from one particle can reach the other particle.

Two charges moving along the orbits of the boost Killing vector in flat space are *at rest* in the Rindler coordinate system and have a constant distance from the spacetime origin, as measured along the slices orthogonal to the Killing vector. Similarly, the worldlines 1 and 1’ are the orbits of the “static” Killing vector $\partial/\partial T$ of de Sitter space. Introducing static coordinates $T \in \mathbb{R}$, $R \in \langle 0, \alpha \rangle$ (ϑ, φ unchanged) by

$$T = \frac{\alpha}{2} \log \frac{\cos \tilde{r} - \cos \tilde{t}}{\cos \tilde{r} + \cos \tilde{t}}, \quad R = \alpha \frac{\sin \tilde{r}}{\sin \tilde{t}},$$

in which the de Sitter metric becomes

$$g_{\text{ds}} = -\left(1 - \frac{R^2}{\alpha^2}\right) dT^2 + \left(1 - \frac{R^2}{\alpha^2}\right)^{-1} dR^2 + R^2 d\omega^2,$$

² “A uniform acceleration” means that the (proper) time derivative \dot{a}^α of the acceleration, projected into the hypersurface orthogonal to the four-velocity, vanishes. The magnitude of the acceleration is constant.

we find that the uniformly accelerated particles 1, 1' have the four-acceleration $-(R_o/\alpha^2) \partial/\partial R$, the magnitude $|a_o|$ being constant, and they are at rest at

$$R = \pm R_o = \mp \frac{a_o \alpha^2}{\sqrt{1 + a_o^2 \alpha^2}}.$$

The particle 1 (1') has, as measured at fixed T , a constant proper distance from the origin $\tilde{t} = \pi/2$, $\tilde{r} = 0$ ($\tilde{r} = \pi$). As with Rindler coordinates in Minkowski space, the static coordinates cover only a ‘‘half’’ of de Sitter space; in the other half the Killing vector $\partial/\partial T$ becomes space-like.

By the conformal transformation of the boosted Coulomb fields in Minkowski space, we constructed [5] test scalar and electromagnetic fields produced by charges moving along the worldlines 1, 1' in de Sitter space. The scalar field from two *identical* scalar charges s is given by

$$\Phi_{\text{sym}} = \frac{s}{4\pi} \frac{1}{\mathcal{Q}}, \quad (3)$$

$$\frac{\mathcal{Q}}{\alpha} = \left[\left(\sqrt{1 + a_o^2 \alpha^2} + a_o R \cos \vartheta \right)^2 - \left(1 - \frac{R^2}{\alpha^2} \right) \right]^{\frac{1}{2}}, \quad (4)$$

(Ref. [5], Eq. (5.4)), whereas the electromagnetic field due to *opposite* charges $+e$ and $-e$ is (Ref. [5], Eq. (5.7))

$$F_{\text{sym}} = -\frac{e}{4\pi} \frac{1}{\mathcal{Q}^3} \frac{a_o \alpha^4}{\sin^3 \tilde{t}} \left[\cos \tilde{t} \sin^2 \tilde{r} \sin \vartheta d\tilde{r} \wedge d\vartheta \right. \\ \left. + (a_o^{-1} \sqrt{a_o^2 + \alpha^{-2}} \sin \tilde{r} + \sin \tilde{t} \cos \vartheta) d\tilde{t} \wedge d\tilde{r} \right. \\ \left. - \sin \tilde{t} \cos \tilde{r} \sin \tilde{r} \sin \vartheta d\tilde{t} \wedge d\vartheta \right]. \quad (5)$$

$$F_{\text{Bas}} = -\frac{e}{4\pi} \frac{\alpha^3}{\mathcal{R}^3} \frac{1}{(\sin \tilde{t} + \cos \tilde{r})^3} \left[-(\sqrt{1 + a_o^2 \alpha^2} \cos \tilde{r} - a_o \alpha \sin \tilde{t}) \cos \vartheta d\tilde{t} \wedge d\tilde{r} \right. \\ \left. + (\sqrt{1 + a_o^2 \alpha^2} - a_o \alpha \cos \tilde{r} \sin \tilde{t}) \sin \tilde{r} \sin \vartheta d\tilde{t} \wedge d\vartheta + a_o \alpha \sin^2 \tilde{r} \cos \tilde{t} \sin \vartheta d\tilde{r} \wedge d\vartheta \right], \quad (8)$$

where

$$\frac{\mathcal{R}}{\alpha} = \frac{[(a_o \alpha \sin \tilde{t} - \sqrt{1 + a_o^2 \alpha^2} \cos \tilde{r})^2 + \sin^2 \tilde{r} \sin^2 \vartheta]^{\frac{1}{2}}}{\sin \tilde{t} + \cos \tilde{r}}.$$

In order to understand explicitly the relation of these fields to the classical Born solutions, consider Minkowski spacetime with spherical coordinates $\{t, r, \vartheta, \varphi\}$ with metric $g_M = -dt^2 + dr^2 + r^2 d\omega^2$. If we set

$$t = -\frac{\alpha \cos \tilde{t}}{\cos \tilde{r} + \sin \tilde{t}}, \quad r = \frac{\alpha \sin \tilde{r}}{\cos \tilde{r} + \sin \tilde{t}},$$

with ϑ, φ unchanged, we find that this Minkowski space is

We call these smooth (outside the sources) fields symmetric because they can be written as a symmetric combination of retarded and advanced effects from both charges.

Although Eqs. (3) and (5) represent fields due to uniformly accelerated charges in de Sitter space, their relation to the Born solutions is not transparent. In order to arrive at such a relation we have to consider sources located symmetrically with respect to the origin $\tilde{r} = 0$, similarly as the charges in the Born solution move along hyperbolae symmetrical with respect to the origin in Minkowski space. Hence, instead of the worldlines 1 and 1' we have to consider the worldlines 2 and 2' (Fig. 2) which, due to homogeneity and isotropy of de Sitter space, also represent uniformly accelerated particles. These worldlines and the resulting fields can be obtained from Eqs. (3)–(5) by a spatial rotation by $\pi/2$. We find the worldlines 2, 2' to be given by

$$\tan \tilde{t} = -\frac{\sqrt{1 + a_o^2 \alpha^2}}{\sinh(\lambda_{\text{ds}} \alpha^{-1} \sqrt{1 + a_o^2 \alpha^2})}, \quad (6)$$

$$\tan \tilde{r} = \pm \frac{\cosh(\lambda_{\text{ds}} \alpha^{-1} \sqrt{1 + a_o^2 \alpha^2})}{a_o \alpha}, \quad \vartheta, \varphi = 0.$$

The scalar and electromagnetic fields produced by sources moving along these worldlines are:

$$\Phi_{\text{Bas}} = \frac{s}{4\pi} \frac{\sin \tilde{t}}{\sin \tilde{t} + \cos \tilde{r}} \frac{1}{\mathcal{R}}, \quad (7)$$

conformally related to de Sitter space as follows (Fig. 2):

$$g_{\text{ds}} = \Omega^2 g_M, \quad \Omega = \frac{\cos \tilde{r} + \sin \tilde{t}}{\sin \tilde{t}} = \frac{2\alpha^2}{\alpha^2 - t^2 + r^2}. \quad (9)$$

In coordinates $\{t, r, \vartheta, \varphi\}$, which can also be used in de Sitter space³ the worldlines 2, 2', Eqs. (6), acquire

³ Coordinates $\{t, r, \vartheta, \varphi\}$ differ from the standard conformally flat coordinates $\{\tilde{t}, \tilde{r}, \vartheta, \varphi\}$ just by shift in \tilde{t} -direction by $\pi/2$ (cf. Figs. 1, 2). Coordinates $\{\tilde{t}, \tilde{r}, \vartheta, \varphi\}$ are related to the usual ‘‘steady-state’’ coordinates $\{\tilde{\eta}, \tilde{r}, \vartheta, \varphi\}$ of exponentially expanding $k = 0$ cosmologies by a simple time rescaling $\tilde{t} = -\alpha \exp(-\tilde{\eta}/\alpha)$.

the simple form: $\vartheta = 0$, $\varphi = 0$, and

$$t = b_o \sinh(\lambda_M/b_o), \quad r = \pm b_o \cosh(\lambda_M/b_o), \quad (10)$$

where λ_M is the proper time as measured by g_M , and

$$b_o/\alpha = \sqrt{1 + a_o^2 \alpha^2} - a_o \alpha.$$

The worldlines (10) are just two hyperbolae (Fig. 2), representing particles with uniform acceleration $1/b_o$ as measured in Minkowski space.

Transforming the fields (7) and (8) into conformally flat coordinates $\{t, r, \vartheta, \varphi\}$, we obtain

$$\Phi_{\text{BdS}} = \Omega^{-1} \frac{s}{4\pi} \frac{1}{\mathcal{R}}, \quad (11)$$

$$F_{\text{BdS}} = -\frac{e}{4\pi} \frac{1}{2b_o} \frac{\alpha^3}{\mathcal{R}^3} \left[-2t r^2 \sin \vartheta dr \wedge d\vartheta - (b_o^2 + t^2 - r^2) \cos \vartheta dt \wedge dr + r(b_o^2 + t^2 + r^2) \sin \vartheta dt \wedge d\vartheta \right], \quad (12)$$

the factor \mathcal{R} now being given by

$$\mathcal{R} = \frac{1}{2b_o} \sqrt{(b_o^2 + t^2 - r^2)^2 + 4b_o^2 r^2 \sin^2 \vartheta}. \quad (13)$$

Expressions (7), (8), and (11), (12) represent *the generalized Born scalar and electromagnetic fields from the sources moving with constant acceleration a_o along the worldlines (6), respectively (10), in de Sitter universe.*

To connect these fields with their counterparts in flat space, note that they are conformally related by transformation (9). Under the conformal transformation, the field Φ_{BdS} in (11) has to be multiplied by factor Ω , which gives $\Phi_{\text{BdS}} = (s/4\pi) \mathcal{R}^{-1}$, and F_{BdS} in (12) remains unchanged. The transformed fields then precisely coincide with the classical Born fields; see e.g. Refs. [1], [2], [7].

In order to see the limit for $\Lambda \rightarrow 0$, we parameterize the sequence of de Sitter spaces by Λ , identifying them in terms of coordinates $\{t, r, \vartheta, \varphi\}$. As $\Lambda = 3/\alpha^2 \rightarrow 0$, Eq. (9) implies $\Omega_\Lambda \rightarrow 2$, $g_{\text{dS}\Lambda} \rightarrow 4g_M$. After the trivial rescaling of t, r by factor 2, the standard Minkowski metric is obtained. The limit of the fields (11) and (12), in which b_o is kept constant (cf. $a_o = (1 - b_o^2 \alpha^{-2})/(2b_o)$), leads to the scalar and electromagnetic Born fields in flat space. Because of the rescaling of coordinates by factor 2, we get the physical acceleration $1/b_o = 2a_o$, and the scalar field rescaled by $1/2$.

What is the character of the generalized Born fields? Focusing on the electromagnetic case, we first decompose the field (8) into the orthonormal tetrad $\{e_\mu\}$ tied to coordinates $\{\tilde{t}, \tilde{r}, \vartheta, \varphi\}$; for example, $e_{\tilde{t}} = (\alpha^{-1} \sin \tilde{t}) \partial/\partial \tilde{t}$, etc., the dual tetrad $e^{\tilde{t}} = (\alpha/\sin \tilde{t}) d\tilde{t}$, etc. Splitting the field into the electric and magnetic parts,

$F_{\text{BdS}} = E \wedge e^{\tilde{t}} + B \cdot e^{\tilde{r}} \wedge e^{\vartheta} \wedge e^{\varphi}$, we get

$$E = \frac{e}{4\pi} \frac{\alpha \sin^2 \tilde{t}}{\mathcal{R}^3 (\sin \tilde{t} + \cos \tilde{r})^3} \times \left[-(\sqrt{1 + a_o^2 \alpha^2} \cos \tilde{r} - a_o \alpha \sin \tilde{t}) \cos \vartheta e_{\tilde{r}} + (\sqrt{1 + a_o^2 \alpha^2} - a_o \alpha \sin \tilde{t} \cos \tilde{r}) \sin \vartheta e_{\vartheta} \right], \quad (14)$$

$$B = -\frac{e}{4\pi} \frac{a_o \alpha^2 \sin^2 \tilde{t}}{\mathcal{R}^3 (\sin \tilde{t} + \cos \tilde{r})^3} \cos \tilde{t} \sin \tilde{r} \sin \vartheta e_{\varphi}.$$

The fields exhibit some features typical for the classical Born solution. The toroidal electric field, E_φ , vanishes; only B_φ is non-vanishing. At $\tilde{t} = \pi/2$, the moment of time symmetry, $B_\varphi = 0$. It vanishes also for $\vartheta = 0$ — there is no Poynting flux along the axis of symmetry.

The classical Born field decays rapidly ($E \sim r^{-4}$, $B \sim r^{-5}$) at spatial infinity, but it is “radiative” ($E, B \sim r^{-1}$) if we expand it along null geodesics $t - r = \text{constant}$, approaching thus null infinity. In de Sitter spacetime with standard slicing, the space is finite (S^3). However, we can approach infinity along spacelike hypersurfaces if, for example, we consider the “steady-state” half of de Sitter universe (cf. Fig. 1) with flat-space slices, i.e., if we take the “conformally flat” time $\tilde{t} = \text{constant}$ (see footnote 3). Introducing the orthogonal tetrad tied to conformally flat coordinates $\{\tilde{t}, \tilde{r}, \vartheta, \varphi\}$, the tetrad components of the fields decay as \tilde{r}^{-2} at $\tilde{t} = \text{constant}$, $\tilde{r} \rightarrow \infty$, so that the Poynting flux falls off as \tilde{r}^{-4} .

The fields decay very rapidly along *timelike* worldlines as \mathcal{I}^+ is approached. This is caused by the exponential expansion of slices $\tau = \text{constant}$ (cf. Eq. (1)). As $\tau \rightarrow \infty$ the electric field (14) becomes radial, $E_{\tilde{r}} \sim \exp(-2\tau/\alpha)$, and $B_\varphi \sim \exp(-2\tau/\alpha)$. The energy density, $u = \frac{1}{2}(E^2 + B^2)$, decays as (expansion factor) $^{-4}$ — as energy density in the radiation dominated standard cosmologies. The density of the conserved energy $u_{\text{conf}} = (\alpha/\sin \tilde{t})u \sim \exp(-3\tau/\alpha)$ (determined by a timelike conformal Killing vector $\partial/\partial \tilde{t}$) gets rarified at the same rate that the volume increases.

Will a slower decay occur if \mathcal{I}^+ is approached along null geodesics? To study the asymptotic behavior of a field along a null geodesic (see, e.g., Ref. [4]), we have to (i) find a geodesic and parameterize it by an affine parameter ζ , (ii) construct a tetrad parallelly propagated along the geodesic, and (iii) study the asymptotic expansion of the tetrad components of the field. The details of these calculations will be presented elsewhere. Here we just list some typical conclusions. Along null geodesics lying in the axis $\vartheta = 0$ (thus crossing the particles’ worldlines) the “radiation field”, i.e., the coefficient of the leading term in $1/\zeta$, vanishes, as could have been anticipated — particles do not radiate in the direction of their acceleration. The radiation field also vanishes along null geodesics reaching infinity along directions *opposite* to those of geodesics emanating from the particles (see Fig. 3). Along all other geodesics, the field *has* ra-

