

Gravitational waves from inspiralling binary black holes

This content has been downloaded from IOPscience. Please scroll down to see the full text.

2002 Class. Quantum Grav. 19 1267

(<http://iopscience.iop.org/0264-9381/19/7/305>)

View [the table of contents for this issue](#), or go to the [journal homepage](#) for more

Download details:

IP Address: 194.94.224.254

This content was downloaded on 08/07/2014 at 08:44

Please note that [terms and conditions apply](#).

Gravitational waves from inspiralling binary black holes

Alessandra Buonanno

Institut d'Astrophysique de Paris (GRECO, FRE 2435 du CNRS), 98bis Boulevard Arago, 75014 Paris, France

and

Theoretical Astrophysics and Relativity Group, California Institute of Technology, Pasadena, 91125 CA, USA

E-mail: buonanno@tapir.caltech.edu

Received 30 November 2001, in final form 28 January 2002

Published 11 March 2002

Online at stacks.iop.org/CQG/19/1267

Abstract

Binary black holes are the most promising candidate sources for the first generation of earth-based interferometric gravitational-wave detectors. We summarize and discuss the state-of-the-art analytical techniques developed during the last few years to better describe the late dynamical evolution of binary black holes of comparable masses.

PACS number: 0460N

1. Introduction

Binary systems made of compact objects (neutron stars or black holes) that spiral in towards coalescence because of gravitational-radiation damping are among the most promising candidate sources for interferometric gravitational-wave (GW) detectors, such as the Laser Interferometric Gravitational Wave Observatory (LIGO), VIRGO, GEO and TAMA [1]. The inspiral waveform will enter the detector frequency band during the last few minutes or seconds of evolution of the binary, and the GW community plans to track the signal phase and build up the signal-to-noise ratio by integrating the signal for the time during which it stays in the detector bandwidth. This is achieved by filtering the detector output with a template which is an (approximate) theoretical copy of the exact, observed signal.

Einstein theory predicts that the radiative transverse traceless (TT) gravitational field h_{ij}^{TT} , far away from the source, is related to the motion and the structure of the source, at lowest order in the post-Newtonian (PN) expansion, by the quadrupole formula

$$h_{ij}^{\text{TT}}(T, D) = \frac{2G}{c^4 D} \mathcal{P}_{ijkl}(N) \frac{d^2}{dT^2} Q_{km} \left(T - \frac{D}{c} \right) \quad (1)$$




Figure 1. Typical gravitational waveform emitted throughout the inspiral, plunge and ring-down phases.

where Q_{ij} ($i, j = 1, 2, 3$) is the tracefree quadrupole moment of the source, D is the distance from the source, $\mathbf{N} = \mathbf{X}/D$ is the unit vector from the source to the observer, $\mathcal{P}_{ijkm}(\mathbf{N})$ is the TT projection operator onto the plane orthogonal to \mathbf{N} , G is the Newton constant and c is the speed of light. From equation (1) (and its extensions at higher-PN orders) we see that the more precisely we know the two-body motion, the more accurately the PN template h_{ij}^{TT} will describe the gravitational waveform detected.

In figure 1 we show a typical gravitational waveform. The part of the waveform drawn with a continuous line is emitted during the inspiral phase when the two black holes are largely separated, e.g., $r \geq 10GM/c^2$ where we denote the radial separation by r and the total mass of the binary system by M . During the inspiral, the two black holes follow an adiabatic sequence of quasi-circular orbits. The equation of motion in the centre-of-mass frame can be written schematically as

$$\frac{d^2\mathbf{x}}{dt^2} = -\frac{GM\mathbf{x}}{r^3} [1 + \mathcal{O}(\epsilon) + \mathcal{O}(\epsilon^2) + \mathcal{O}(\epsilon^{5/2}) + \mathcal{O}(\epsilon^3) + \dots] \times [1 + \mathcal{O}(v) + \mathcal{O}(v^2) + \dots] \quad (2)$$

where \mathbf{x} denotes the separation vector between the two bodies and $r = |\mathbf{x}|$. Equation (2) is characterized by a double expansion: in the PN parameter $\epsilon \sim v^2/c^2 \sim GM/(c^2r)$ and in the parameter $v = m_1m_2/M^2$, where m_1 and m_2 are the masses of the two black holes. The parameter v ranges between 0 (test-mass limit) and 1/4 (equal-mass case).

It is well known that the PN expansion converges badly: as the two bodies draw closer and enter the nonlinear strong curvature phase, the motion becomes relativistic, e.g., $v/c \sim 0.3$, and it is more and more difficult to extract reliable information from the PN series. Specifically, when the distance between the inspiralling black holes shrinks to $r \lesssim 10GM/c^2$, the PN expansion can no longer be trusted [2]. The dashed line in figure 1 depicts the (less known) part of the waveform emitted during the final phase of evolution when nonlinearities and strong curvature effects become important and nonperturbative analytical and/or numerical techniques should be used to describe it. This final phase includes the transition from adiabatic inspiral to plunge, beyond which the two-body motion is driven (almost) only by the conservative part of the dynamics. For nonspinning binary black holes, the plunge starts at the innermost stable circular orbit (ISCO) of the binary black holes. Beyond the plunge

the two black holes merge, forming a Kerr black hole. As the system reaches the stationary Kerr state, the nonlinear dynamics of the merger resemble, more and more, the oscillations of the black-hole quasi-normal modes [3] and the gravitational signal will be a superposition of exponentially damped sinusoids (ring-down waveform). However, if the black holes carry a big spin, it is likely that the plunge stage will never be reached and the late dynamical phase will be much more complicated in this case.

It is likely that the first detection of gravitational waves with LIGO/VIRGO... interferometers will be from binary systems made of black holes of comparable masses, say, with a total mass $M \simeq 10M_{\odot}$ – $30M_{\odot}$. In fact, if we assume that the binary black holes are originated by ‘population synthesis’ [4], the estimated detection rate per year is $\lesssim 4 \times 10^{-3}$ – 0.6 at 100 Mpc [5], while if globular clusters are considered as ‘machines’ for making binary black holes [6], the detection rate per year is ~ 0.04 – 0.6 at 100 Mpc [5]. These numbers, especially the latter, are more optimistic than the estimated detection rate per year for neutron-star binaries, which is $\lesssim 3 \times 10^{-4}$ – 0.3 at 20 Mpc [5] or, for neutron-star/black-hole binaries, $\lesssim 4 \times 10^{-4}$ – 0.6 at 43 Mpc [5].

Although the number of cycles in the LIGO/VIRGO... frequency band of gravitational waves emitted by a comparable-mass black holes is not very high, on the order of 10–200, these particular sources demand a more careful analysis because the gravitational waves the detectors will be sensitive to are emitted during the final stages of inspiral where PN expansion fails. For example, in the nonspinning case, the GW frequency at the ISCO (evaluated using the Schwarzschild ISCO) is $f_{\text{GW}}^{\text{ISCO}} \simeq 220$ Hz for $M = 20M_{\odot}$ and $f_{\text{GW}}^{\text{ISCO}} \simeq 167$ Hz for $M = 30M_{\odot}$, well inside the LIGO/VIRGO... band. Moreover, comparable-mass black holes could carry big spins which could affect the waveforms, so if data analysis will be done with spinless templates, there is considerable chance to miss the gravitational-wave signal.

In the next section we shall summarize what has been done in the literature to cope with these problems and which issues remain to be solved.

2. Analytical methods to predict gravitational waveforms

For spinless black-hole binaries, we consider the so-called restricted waveform $h(t) = v^2 \cos(\varphi_{\text{GW}}(t))$, where $\varphi_{\text{GW}} = 2\varphi$ with φ being the orbital phase and v the invariantly defined velocity $v = (M\dot{\varphi})^{1/3} = (\pi M f_{\text{GW}})^{1/3}$. This waveform is obtained disregarding all the multipolar components appearing in the gravitational waveform except the quadrupolar one (see equation (1)). To determine in the adiabatic limit the evolution of the GW phase, $\varphi_{\text{GW}}(t)$, it is sufficient to use the energy-balance equation

$$\frac{d\mathcal{E}}{dt} = -\mathcal{F} \quad (3)$$

relating the orbital energy function \mathcal{E} (centre-of-mass energy that is conserved in absence of radiation reaction) to the gravitational-flux (or luminosity) function \mathcal{F} , which are known for quasi-circular orbits as a PN expansion in v . It is easily shown that equation (3) is equivalent to the following system of differential equations (see, e.g., [22]):

$$\frac{d\varphi_{\text{GW}}}{dt} = \frac{2v^3}{M} \quad \frac{dv}{dt} = -\frac{\mathcal{F}(v)}{M d\mathcal{E}(v)/dv} \quad (4)$$

whose solution provides the phasing $\varphi_{\text{GW}}(t)$ during the inspiral.

2.1. ‘Genuine’ post-Newtonian calculations

Let us summarize what we know about the two crucial ingredients \mathcal{E} and \mathcal{F} entering equation (4). The equations of motion of two compact bodies at 2.5PN approximation were

first derived in [7]. The 3PN equations of motion have been obtained by two separate groups: Damour, Jaranowski and Schäfer [8] used the Arnowitt–Deser–Misner (ADM) canonical approach, while Blanchet, Faye and de Andrade [10] worked with the PN iteration of the Einstein equations in harmonic gauge. Recently, Damour *et al* [9], working in the ADM formalism and applying dimensional regularization, uniquely determined the so-called static parameter which entered the 3PN equations of motion [8, 10] and till then was unknown. Thus, at the present time, the energy function \mathcal{E} is known up to 3PN order.

The gravitational flux emitted by compact binaries was first computed at 1PN order in [11]. Subsequently, it was determined at 2PN order using a formalism based on multipolar and post-Minkowskian approximations and independently by using a direct integration of the relaxed Einstein equations [12]. Nonlinear effects of tails at 2.5PN and 3.5PN orders were computed in [13]. More recently, the gravitational-flux function for quasi-circular orbits has been derived up to 3.5PN order [14]. However, at 3PN order [14], the gravitational-flux function depends on an arbitrary parameter, which is not fixed by the regularization scheme used by the authors.

Although the knowledge of higher-order PN corrections to \mathcal{E} and \mathcal{F} is a necessary ingredient to extract the phase of gravitational signals emitted by neutron star or black-hole/neutron-star binaries, it is not sufficient by itself for computing GWs of comparable-mass binaries. As underlined above, this is due to the fact that, for these sources, LIGO/VIRGO... will detect gravitational signals emitted when the motion is relativistic and (genuine) perturbative PN calculations can no longer be trusted.

2.2. Post-Newtonian resummation methods

Certainly, the best way of extracting the GW signal emitted by comparable-mass binaries during the last stages of inspiral would be to solve numerically the Einstein equations of a binary black-hole system. Unfortunately, despite the interesting progress made by the numerical relativity community during the recent years [15–20], an estimate of the waveform emitted by a black-hole binary has not yet been provided. Preliminary results for the plunge, merger and ring-down waveforms were only recently obtained [21] and they use initial conditions at the ISCO which differ from PN predictions. To overcome this gap and tackle the delicate issue of the late dynamical evolution, various *nonperturbative analytical* approaches have been proposed [22–25] to study the motion of two spinning and nonspinning bodies in general relativity.

The main features of the various PN resummation methods can be summarized as follows: (i) they provide an analytical (gauge-invariant) resummation of the orbital energy function \mathcal{E} and gravitational-flux function \mathcal{F} , (ii) they can describe the motion (and provide the gravitational waveform) beyond the adiabatic approximation and (iii) they can, in principle, be extendible to higher-PN orders. More importantly, they can be used to provide initial data for black holes just starting the plunge motion which can be used by the numerical relativity community to evolve the full Einstein equations during the merger phase. However, the resummation methods are also based on some assumptions that are hard to prove rigorously—for example in deriving the orbital energy and the gravitational-flux functions in the comparable-mass case, it is assumed that they are smooth deformations of the analogous quantities in the test-mass limit. Moreover, in absence of an exact solution or experimental data, we can test the robustness and reliability of those resummation approaches using only internal convergence tests.

As underlined at the beginning of section 2, in the absence of spins, the two crucial ingredients necessary to extract the GW phasing are the orbital energy function \mathcal{E} and the

gravitational-flux function \mathcal{F} . In section 2.2.1, we shall discuss a resummation approach which provides a better-behaved flux-type function [22], while in section 2.2.2, we shall summarize another resummation method which provides an improved energy-type function [23, 25]. By combining these two resummation methods, Buonanno and Damour [24] proposed a way of describing the black-hole motion beyond the adiabatic limit, including the transition from inspiral to plunge, whose main features are discussed in section 2.3.

2.2.1. Padé approximants. Starting from the PN expansions of $\mathcal{E}(v)$ and $\mathcal{F}(v)$ Damour, Iyer and Sathyaprakash [22] proposed a new class of waveforms based on the systematic use of Padé resummation, which is a standard mathematical technique used to accelerate the convergence of poorly converging or even divergent power series. For lack of space we shall discuss briefly only the Padé approximant to the flux function \mathcal{F} .

In the test-mass limit ($v \rightarrow 0$), the flux function has a simple pole at the light ring ($v^2 = 1/3$). Damour *et al* [22] argued that the origin of this pole is quite general and that we should expect a pole singularity also in the equal-mass case. Thus, after factoring out this pole (defined by what they call the better-behaved energy function $e(v)$, see [22]), they introduced the better-behaved flux function $f(v) \equiv (1 - v/v_{\text{pole}})\mathcal{F}(v)$. Using the 2.5PN expansion of $\mathcal{F}(v)$ and applying the (near diagonal) Padé resummation to $f(v)$, they derived the v^n -Padé approximant to the gravitational-flux function:

$$\mathcal{F}_{\text{Padé}'}^n(v; \nu) = \frac{1}{1 - v/v_{\text{pole}}^n(\nu)} f_{\text{Padé}'}^n(v; \nu). \quad (5)$$

To test the reliability of the result, Damour *et al* showed that in the test-mass limit the (near diagonal) Padé approximant to $\mathcal{F}(v; \nu = 0)$, given by equation (5), exhibits a very good convergence towards the exact result, which is numerically known when $\nu = 0$. (See figure 3 in [22].) Thus, arguing that the equal-mass case can be obtained as a smooth deformation of the test-mass limit, with deformation parameter ν , they propose $\mathcal{F}_{\text{Padé}'}^n(v; \nu)$ as the best estimation of the GW flux for comparable-mass binaries.

2.2.2. Effective-one-body reduction. The resummation technique discussed in this section, the so-called effective-one-body (EOB) approach [23], was originally inspired by a similar approach introduced by Brézin, Itzykson and Zinn-Justin [26] to study electromagnetically-interacting two-bodies. The basic idea, illustrated in figure 2, is to map the *real* conservative two-body dynamics up to 2PN order (see below for the extension at 3PN order) onto an *effective* one-body problem, where a test particle of mass $\mu = m_1 m_2 / M$, with m_1, m_2 being the black-hole masses and $M = m_1 + m_2$, moves in some effective background metric $g_{\mu\nu}^{\text{eff}}$. This mapping has been worked out within the Hamilton–Jacobi formalism by imposing that, whereas the action variables of the real and effective description coincide, i.e. $L_{\text{real}} = L_{\text{eff}}$, $\mathcal{I}_{\text{real}} = \mathcal{I}_{\text{eff}}$, where L denotes the total angular momentum and \mathcal{I} the radial action variable, the energy axis is allowed to change, $E_{\text{real}} = f(E_{\text{eff}})$ where f is a generic function. By applying the above rules defining the mapping, it was found that as long as radiation-reaction effects are not taken into account, the effective metric is just a deformation of the Schwarzschild metric, with deformation parameter $\nu = \mu/M$. The effective metric reads [23]

$$ds_{\text{eff}}^2 = -A(R)c^2 dt^2 + \frac{D(R)}{A(R)} dR^2 + R^2 d\Omega^2. \quad (6)$$

More importantly the reduction to the one-body dynamics provides the *improved* real Hamiltonian

$$\mathcal{H}_{\text{real}}^{\text{improved}} = Mc^2 \sqrt{1 + 2\nu \left(\frac{\mathcal{H}_{\text{eff}}^{\nu} - \mu c^2}{\mu c^2} \right)} \quad (7)$$




Figure 2. How the EOB approach matches the real two-body problem (on the left) and the effective one-body problem (on the right).

where

$$\mathcal{H}_{\text{eff}}(v, R, P_R, P_\varphi) = \mu c^2 \sqrt{A(R) \left(1 + \frac{A(R) P_R^2}{\mu^2 c^2 D(r)} + \frac{P_\varphi^2}{\mu^2 c^2 R^2} \right)}. \quad (8)$$

We refer to the real Hamiltonian (7) as the improved real Hamiltonian because, being a (partial) PN resummation of the original badly-convergent real Hamiltonian $\mathcal{H}_{\text{real}} = \mathcal{H}_{\text{New}} + \mathcal{H}_{1\text{PN}}/c^2 + \mathcal{H}_{2\text{PN}}/c^4 \dots$, it should better capture the nonperturbative effects of the final stage of the black-hole motion. Remarkably, the mapping between the real and the effective Hamiltonians in equation (7) coincides with the mapping obtained in [26] in the context of quantum electrodynamics, where these authors mapped the one-body relativistic Balmer formula onto the two-body energy formula.

The EOB approach was then extended at 3PN order in [25]. The authors found that, starting from the 3PN level, there are more equations to satisfy than the number of free parameters appearing in the energy map and in the effective metric. Hence, they suggested the following two possibilities. At the price of modifying the coefficients of the effective metric at 1PN and 2PN levels, and the energy map (7) as well, it is still possible at 3PN order to (uniquely) map the real two-body dynamics onto the dynamics of a test mass moving on a geodesic (see appendix A of [25] for details). However, this solution looks quite complicated and, more importantly, it does not look very natural to wait until one knows the 3PN Hamiltonian to derive the matching at 1PN and 2PN levels. The authors then suggested abandoning the hypothesis (used at 2PN order [23]) that the effective test mass moves along a geodesic and introduced in the Hamilton–Jacobi equation (arbitrary), higher-derivative terms which provide enough coefficients to obtain the matching. Because of these terms, the effective 3PN Hamiltonian is not uniquely fixed by the matching rules defined above; the general expression is given by




Figure 3. Comparison between the PN-expanded (left) and the EOB (right) binding energies for equal-mass binary.

equation (3.12) in [25]. It is interesting to note that in the non-geodesic case the relation between the effective and real Hamiltonians is still given by equation (7).

For a qualitative understanding of the way the EOB can accelerate the PN convergence, in figure 3 we compare the PN-expanded binding energy (left panel) to the EOB-resummed binding energy (right panel) versus the radial separation, for circular orbits, in the equal-mass case, for two choices of the angular momentum L and at different PN orders. (To obtain the plots, we used the 3PN-expanded and EOB-resummed Hamiltonians given by equations (3.12) and (4.27) of [25] with $\omega_s = 0$ [9] and $z_1 = 0 = z_2$. We also used the (generalized) canonical transformation between the ADM and the ‘effective’ coordinates given by equations (4.29)–(4.36) of [25].)

Note that the PN-expanded energy oscillates at the various PN orders while the EOB energy has a monotonic behaviour. The fractional difference between the PN-expanded and EOB binding energies is, e.g., for $L = 3.2\mu GM$, 2.3% (2PN order) and 0.24% (3PN order) at $R = 9GM/c^2$ and 17.6% (2PN order) and 2.8% (3PN order) at $R = 6GM/c^2$.

An interesting nonperturbative feature of binary black-hole systems is the presence of the ISCO defined as the solution of the equations $\partial\mathcal{H}/\partial r = 0 = \partial^2\mathcal{H}/\partial r^2$. In the test-mass limit (Schwarzschild metric) the ISCO exists and the nonrelativistic energy associated with it is $\mathcal{E}_{\text{isco}}^{\text{Schw}} = -0.01430Mc^2$. If we consider the PN-expanded real Hamiltonian (equation (4.27) of [25]) in the test-mass limit and look for the ISCO, we find that at 1PN order $\mathcal{E}_{\text{isco}}^{\text{PN-exp}} = -0.00778Mc^2$, at 2PN order the ISCO does not exist, while at 3PN order $\mathcal{E}_{\text{isco}}^{\text{PN-exp}} = -0.01129Mc^2$. On the other hand, because by definition the EOB converges in the test-mass limit to the Schwarzschild case, it automatically provides the correct ISCO. Thus, in the same way as the Padé approximants, the EOB resummation method provides, by construction, the right prediction in the test-mass limit. Although we cannot prove that in the comparable-mass case the EOB approach is converging to the right limit, we can certainly say that it shows reasonable stability in the predictions at 1PN, 2PN and 3PN orders.

In figure 4, we summarize the binding energy at the ISCO predicted by various post-Newtonian [23, 25, 27] and numerical relativity calculations [15–19]. The equal-mass ISCO binding energies predicted by the EOB approach at various PN orders are: $\mathcal{E}_{\text{isco}}^{\text{EOB}} = -0.01440Mc^2$ (1PN order), $\mathcal{E}_{\text{isco}}^{\text{EOB}} = -0.01498Mc^2$ (2PN order) and $\mathcal{E}_{\text{isco}}^{\text{EOB}} = -0.01670Mc^2$ (3PN order). Note that the fractional difference from 1PN to 2PN order is 4% and it increases to 10% from 2PN to 3PN order. Figure 4 shows that, except for the




Figure 4. Summary of the binding energy at the ISCO predicted by post-Newtonian and numerical relativity calculations for nonspinning binary black holes of total mass $15M_{\odot} + 15M_{\odot}$.

very recent result of [19], which is much closer to the PN estimates than the numerical relativity predictions, the PN and numerical relativity results differ very much. Yet the reason of these discrepancies has not been understood. A possible explanation could be related to the conformally flat spatial-metric assumption used in numerical relativity simulations, which already leads to a theory that differs from Einstein general relativity at 2PN order (see discussions in [9, 36]).

2.3. Beyond the adiabatic approximation: transition inspiral to plunge

Let us now introduce radiation-reaction effects. As discussed in section 2.2.1, the Padé approximants [22] give a good estimate of the energy-loss rate \mathcal{F} along circular orbits up to 2.5PN order. In [24], this resummation method was combined with the EOB approach and the authors deduced a system of ordinary differential equations which describe the late dynamical evolution of a binary–black-hole system. In spherical coordinates $(\varphi, R, P_{\varphi}, P_R)$, their relevant equations are [24]

$$\begin{aligned} \frac{dR}{dt} &= \frac{\partial \mathcal{H}_{\text{real}}^{\text{impr}}}{\partial P_R} & \frac{dP_R}{dt} + \frac{\partial \mathcal{H}_{\text{real}}^{\text{impr}}}{\partial R} &= 0 \\ \frac{d\varphi}{dt} &= \frac{\partial \mathcal{H}_{\text{real}}^{\text{impr}}}{\partial P_{\varphi}} & \frac{dP_{\varphi}}{dt} &= -\frac{\mathcal{F}_{\text{Padé}}(\dot{\varphi})}{\dot{\varphi}}. \end{aligned} \quad (9)$$

Different from equation (2), the above equations go beyond the adiabatic approximation (at least for the conservative part of the dynamics) and can be analytically or numerically solved to study the transition between the adiabatic inspiral and the plunge.

Let us discuss briefly the main features of this transition in the two extreme limits $\nu \ll 1$ and $\nu = 1/4$ (see [24] for more details). The case $\nu \ll 1$ refers to binary–black-hole systems in which a very small black hole spirals around a supermassive black hole. They are typical GW sources for the Laser Interferometer Space Antenna (LISA). In this case, it was found [23, 28] that the transition from adiabatic inspiral to plunge is sharply localized around the ISCO. Ori and Thorne [28] pointed out that LISA could possibly observe the transition from inspiral to plunge with a low signal-to-noise ratio.




Figure 5. Plunge and ring-down gravitational waveform obtained from the EOB approach [24] for an equal-mass binary.

For equal-mass binaries ($\nu = 1/4$), the radiation damping effects are important in an extended region on the order of $\Delta(Rc^2/GM) \sim 1$ above the naïve (Schwarzschild) ISCO $R = 6GM/c^2$. The transition from inspiral to plunge is rather blurred and the dephasing between the full and the adiabatic waveform becomes visible somewhat before the naïve ISCO. The plunge part of the exact waveform looks like a continuation of the inspiral part because the orbital motion remains quasi-circular throughout the plunge.

Figure 5 shows the plunge, merger and ring-down part of the waveform [24] for an equal-mass binary. The ring-down waveform contains only the mode that is damped more slowly, $l = 2, m = 2$ [3], at frequency $\omega_{qnm} \sim 1880(10M_\odot/M_{\text{BH}})$ Hz, where M_{BH} is the mass of the final black hole formed. The dimensionless rotation parameter is $a_{\text{BH}} = J_{\text{BH}}/(GM_{\text{BH}}^2) = 0.795$, where we denoted the angular momentum of the final Kerr black hole by J_{BH} . The energy emitted during the plunge is $\sim 0.7\%$ of M , with a comparable-energy loss $\sim 0.7\%$ of M during the ring-down phase [30]. This gives a total energy released of $\sim 1.4\%$ of M , which contrasts with the much larger value, 4–5%, of M recently estimated in [21].

Recently, Damour, Iyer and Sathyaprakash [29] investigated the consequences of the EOB waveform for LIGO/VIRGO... data analysis. They showed that the GW radiation coming from the plunge and merger can significantly enhance the signal-to-noise ratio for binaries of total mass $M \gtrsim 30M_\odot$. They found that the signal-to-noise ratio reaches the maximum value of ~ 8 for $M \simeq 80M_\odot$ at 100 Mpc. Previous estimations using maximally-spinning binaries found that the merger dominates on inspiral. For example, Flanagan and Hughes [31] predicted that the energy released during the plunge is $\sim 10\%M$ while during the ring-down phase $\sim 3\%M$, and the signal-to-noise ratio reaches the maximum value of ~ 40 for $M \simeq 200M_\odot$ at 100 Mpc.

3. Open issues

3.1. Spinning binary black holes

The theoretical prediction of GWs from comparable-mass binaries is not only affected by the failure of the PN-expansion, but also by spin effects. Various studies [32, 33] estimated that




Figure 6. Expected Fourier-domain phase $\phi(f)$ of the GW signal versus frequency for various approaches modelling the two-body dynamics.

if the binary’s holes carry a spin, the evolution in time of the GW phase will be significantly affected by it—for example, the spin can introduce modulations and irregularities in the gravitational waveforms. These features can become quite ‘dramatic’ as long as the two spins are big and not aligned or anti-aligned with the orbital angular momentum.

To give a general idea of the effects we are alluding to and currently investigating [40], in figure 6 we sketch the Fourier-domain phase $\phi(f)$ of the GW signal versus frequency for various approaches, including possible modulations due to spin effects. The various methods give the same prediction for the Fourier-domain phase up to the frequency where the PN series fails—for example for $M = 20M_{\odot}$, $f_{\text{PN-failure}} \sim 50$ Hz, while for $M = 30M_{\odot}$, $f_{\text{PN-failure}} \sim 33$ Hz [2], which are well inside the LIGO/VIRGO . . . band.

More recently, Levin [34] claimed that because of spin effects the two-body dynamics could be affected by chaos, or more generally the dynamics, and as a consequence the gravitational waveform, could depend strongly on the initial conditions. However, Schnittman and Rasio re-analysed this issue in [35]. By calculating the divergence of nearby trajectories for a broad sample of initial conditions, they concluded that the divergent time is much greater than the inspiral time. So, even if chaos were present, it should not affect the detection of inspiral waveforms.

To tackle these delicate issues it would be desirable to extend the resummation methods discussed above to the case of spinning binaries. In this respect, we note that the Padé resummation method has been recently extended to the gravitational-flux function including spin–orbit and spin–spin effects [37]. Moreover, on the line of the EOB approach, Damour [36] has recently mapped the conservative dynamics of two spinning black holes into the one of a test particle moving in an external ν -deformed Kerr metric. It will be very important to complete Damour’s analysis by including radiation-reaction effects and describe the late nonadiabatic dynamics of spinning black-hole binaries.

3.2. Possible strategy for not missing the GW signal from comparable-mass binary black holes

In sections 2.2.1 and 2.2.2 we discussed the *best* theoretical gravitational waveforms currently available. These waveforms should certainly be used in detecting and extracting physical parameters (masses and spins) for GWs emitted by neutron-star binaries and neutron-star/black-hole binaries. On the other hand, the direct application of these templates to

detect GWs from comparable-mass black-hole binaries is less straightforward. Indeed, the approaches analysed in sections 2.2.1 and 2.2.2 are inevitably affected by approximations and assumptions, introduced to overtake the fact that the two-body problem is known only up to a certain PN order. It would be important for detection purposes to quantify such uncertainties and take them into account when building the best theoretical GW templates. If not, the risk is to miss the GW signal. A possibility [38] is to *deform* and *expand* (e.g., by introducing new parameters) the template family generated by the resummation method with better PN convergent properties, in such a way that the new (higher-dimensional) template space could (i) include, or at least be not very far from, all other approaches characterized by worse PN convergent properties and (ii) describe signals (hopefully the real GW signal!) whose functional form cannot be described by any of the original template families. To reach this goal the chirp fast transform technique, recently proposed in [39], deserves attention. Moreover, for comparable-mass binaries, the inclusion of spin effects and the enlargement of the template space should hopefully, not make the total number of templates huge because the number of cycles expected in the LIGO/VIRGO... band for nonspinning binaries is already rather small ~ 20 – 100 . (The situation would be different for neutron stars or neutron-star/black-hole binaries for which, in the non-spinning case, we expect a number of cycles on the order of $\sim 10^4$ – 10^6 .)

The new template space [40] could be used for an on-line search. When an acceptable signal-to-noise ratio is found, say $\gtrsim 8$, then the output signal could be re-analysed by templates provided, e.g., by the PN Taylor-expanded Padé approximant and EOB methods, to determine which approach better describes the real signal and extract the binary's parameters as the masses and the spins.

Acknowledgments

The author wishes to thank T Damour, Y Chen, D Chernoff, B S Sathayaprakash, K S Thorne and M Vallisneri for fruitful collaborations and/or very useful discussions.

Note added. While this manuscript was being refereed, an interesting paper by Blanchet [41] appeared on the web. He compares the minimum energy for equal-mass–binary circular orbits at 3PN order with the numerical relativity results [19] and finds good agreement.

References

- [1] Abramovici A *et al* 1992 *Science* **256** 325
Caron B *et al* 1997 *Class. Quantum Grav.* **14** 1461
Lück H *et al* 1997 *Class. Quantum Grav.* **14** 1471
Ando M *et al* 2001 *Preprint astro-ph/0105473*
- [2] Brady P R, Creighton J D E and Thorne K S 1998 *Phys. Rev. D* **58** 061501
- [3] Chandrasekhar S and Detweiler S 1975 *Proc. R. Soc. A* **344** 441
- [4] Lipunov V M, Postnov K A and Prokhorov M E 1997 *New Astron.* **2** 43
- [5] Thorne K S 2000 The scientific case for mature LIGO interferometers *LIGO Document* N. P000024-00-R
webpage www.ligo.caltech.edu/docs/P/P000024-00.pdf
- [6] Portegies Zwart S F and McMillan S L 1999 *Preprint astro-ph/9910061*
- [7] Damour T and Deruelle N 1981 *Phys. Lett. A* **87** 81
Damour T 1982 *C. R. Acad. Sci. II* **294** 1355
- [8] Jaranowski P and Schäfer G 1998 *Phys. Rev. D* **57** 7274
Jaranowski P and Schäfer G 1999 *Phys. Rev. D* **60** 124003
Damour T, Jaranowski P and Schäfer G 2000 *Phys. Rev. D* **62** 044024
Damour T, Jaranowski P and Schäfer G 2000 *Phys. Rev. D* **62** 021501
Damour T, Jaranowski P and Schäfer G 2001 *Phys. Rev. D* **63** 044021

- [9] Damour T, Jaranowski P and Schäfer G 2001 *Phys. Lett. B* **513** 147
- [10] Blanchet L and Faye G 2000 *Phys. Lett. A* **271** 58
(Blanchet L and Faye G 2000 *Preprint* gr-qc/0006100)
Blanchet L and Faye G 2000 *Phys. Rev. D* **63** 062005
de Andrade V C, Blanchet L and Faye G 2001 *Class. Quantum Grav.* **18** 753
- [11] Wagoner R V and Will C M 1976 *Astrophys. J.* **210** 764
- [12] Blanchet L, Damour T, Iyer B R, Will C M and Wiseman A G 1995 *Phys. Rev. Lett.* **74** 3515
Blanchet L, Damour T and Iyer B R 1995 *Phys. Rev. D* **51** 536
Will C M and Wiseman A G 1996 *Phys. Rev. D* **54** 4813
- [13] Blanchet L 1996 *Phys. Rev. D* **54** 1417
Blanchet L 1998 *Class. Quantum Grav.* **15** 113
- [14] Blanchet L, Iyer B R and Joguet B 2001 *Preprint* gr-qc/0105098
- [15] Cook G B 1994 *Phys. Rev. D* **50** 5025
- [16] Baumgarte T W 2000 *Phys. Rev. D* **62** 024018
- [17] Baumgarte T W, Cook G B, Scheel M A, Shapiro S L and Teukolsky S A 1998 *Phys. Rev. D* **57** 7299
- [18] Pfeiffer H P, Teukolsky S A and Cook G B 2000 *Phys. Rev. D* **62** 104018
- [19] Gourgoulhon E, Grandclément P and Bonazzola S 2001 *Preprint* gr-qc/0106015
(Grandclément P, Gourgoulhon E and Bonazzola S 2001 *Preprint* gr-qc/0106016)
- [20] Alcubierre M, Brügmann B, Pollney D, Seidel E and Takahashi R 2001 *Phys. Rev. D* **64** 061501
Kidder L E, Scheel M A and Teukolsky S A 2001 *Phys. Rev. D* **64** 064017
- [21] Baker J, Brügmann B, Campanelli M, Lousto C O and Takahashi R 2001 *Preprint* gr-qc/0102037
- [22] Damour T, Iyer B R and Sathyaprakash B S 1998 *Phys. Rev. D* **57** 885
- [23] Buonanno A and Damour T 1999 *Phys. Rev. D* **59** 084006
- [24] Buonanno A and Damour T 2000 *Phys. Rev. D* **62** 064015
- [25] Damour T, Jaranowski P and Schäfer G 2000 *Phys. Rev. D* **62** 084011
- [26] Brézin E, Itzykson C and Zinn-Justin J 1970 *Phys. Rev. D* **1** 2349
- [27] Kidder L E, Will C M and Wiseman A G 1992 *Class. Quantum Grav.* **9** L127
Kidder L E, Will C M and Wiseman A G 1993 *Phys. Rev. D* **47** 3281
- [28] Ori A and Thorne K S 2000 *Phys. Rev. D* **62** 124022
- [29] Damour T, Iyer B R and Sathyaprakash B S 2001 *Phys. Rev. D* **63** 044023
- [30] Buonanno A and Damour T 2000 *9th Marcel Grossmann Meeting (Rome, July 2000)*
(Buonanno A and Damour T 2000 *Preprint* gr-qc/0011052)
- [31] Flanagan E E and Hughes S A 1998 *Phys. Rev. D* **57** 4535
- [32] Kidder L E, Will C M and Wiseman A G 1993 *Phys. Rev. D* **47** 4183
Kidder L E 1995 *Phys. Rev. D* **52** 821
- [33] Apostolatos T A, Cutler C, Sussman G J and Thorne K S 1994 *Phys. Rev. D* **15** 6274
Apostolatos T A 1996 *Phys. Rev. D* **54** 2438
- [34] Levin J 2000 *Phys. Rev. Lett.* **84** 3515
(Levin J 2000 *Preprint* gr-qc/0010100)
- [35] Schnittman J D and Rasio F 2001 *Phys. Rev. Lett.* **87** 121101
- [36] Damour T 2001 *Phys. Rev. D* **64** 124013
- [37] Porter E and Sathyaprakash B S in preparation
- [38] Buonanno A, Chen Y and Vallisneri M in preparation
- [39] Jenet F and Prince T 2000 *Phys. Rev. D* **62** 122001
- [40] Buonanno A, Chen Y, Chernoff D and Vallisneri M in preparation
- [41] Blanchet L 2001 *Preprint* gr-qc/0112056