

Effective loop quantum cosmology as a higher-derivative scalar-tensor theory

Hongguang Liu,^{1,2,3,*} Karim Noui,^{2,3,†} Edward Wilson-Ewing,^{4,‡} and David Langlois^{3,§}

¹*Centre de Physique Théorique (UMR CNRS 7332) ,*

Universités d'Aix-Marseille et de Toulon, 13288 Marseille, France

²*Laboratoire de Mathématiques et Physique Théorique (UMR CNRS 7350),*

Université François Rabelais, Parc de Grandmont, 37200 Tours, France

³*Laboratoire Astroparticule et Cosmologie,*

Université Denis Diderot Paris 7, 75013 Paris, France

⁴*Max Planck Institute for Gravitational Physics (Albert Einstein Institute),*

Am Mühlenberg 1, 14476 Golm, Germany, EU

(Dated: April 3, 2017)

Recently, Chamseddine and Mukhanov introduced a higher-derivative scalar-tensor theory which leads to a modified Friedmann equation allowing for bouncing solutions. As we note in the present work, this Friedmann equation turns out to reproduce exactly the loop quantum cosmology effective dynamics for a flat isotropic and homogeneous space-time. We generalize this result to obtain a class of scalar-tensor theories, belonging to the family of mimetic gravity, which all reproduce the loop quantum cosmology effective dynamics for flat, closed and open isotropic and homogeneous space-times.

I. INTRODUCTION

Understanding the very early universe remains a fascinating open question in cosmology. In the context of general relativity, an expanding universe containing “standard” matter fields (which satisfy the null energy condition) is generically associated with an initial singularity, where the space-time curvature becomes infinite. In this sense, classical general relativity fails to explain the very origin of our universe. When the value of the curvature approaches the Planck scale, quantum gravity effects are expected to become physically important and could prevent the formation of space-time singularities. This is exactly what happens in the context of loop quantum cosmology (LQC) [1–4] where quantum gravity effects are repulsive, in opposition to attractive classical gravity, and lead to a bouncing universe.

However, quantizing gravity might not be necessary to resolve the initial cosmological singularity and one could envisage modifications of gravity at high curvature so that the singularity in general relativity is replaced by a bounce in a modified gravity theory (see [5, 6] for recent reviews). Of course, these two approaches to avoid the singularity could be two sides of the same coin if the classical equations derived from modified gravity can be interpreted as an effective description of the quantum behaviour.

Scalar-tensor theories provide a very large class of models for modified gravity theories. Among these, higher-order scalar-tensor theories, whose Lagrangians contain not only first order but also second order derivatives of the scalar field, have attracted a lot of attention lately. Allowing for higher-order time derivatives in the Lagrangian is potentially dangerous as this could lead to higher-order equations of motion which may require extra initial conditions and thus introduce an additional degree of freedom, known as the Ostrogradsky ghost, because it leads to an Ostrogradsky

*Electronic address: hongguang.LIU@etu.univ-amu.fr

†Electronic address: karim.noui@lmpt.univ-tours.fr

‡Electronic address: wilson-ewing@aei.mpg.de

§Electronic address: langlois@apc.univ-paris7.fr

instability [7, 8]. It is however possible to find higher-order scalar-tensor theories that contain a single scalar degree of freedom (in addition to the usual tensorial modes associated with gravity) by imposing some restrictions on the initial Lagrangian. Initially, it was believed that a theory of this type was necessarily characterized by second order Euler-Lagrange equations, thus pointing to Horndeski theories [9] (see also [10]). In fact, requiring second order equations of motion turns out to be restrictive and a much larger class of theories, dubbed Degenerate Higher-Order Scalar-Tensor (DHOST) theories, has been recently identified, showing that the absence of an extra unstable scalar mode is compatible with higher order Euler-Lagrange equations [11–18]. These theories could provide an interesting arena to construct models for the early universe, as well as late-time cosmology.

Among DHOST theories, one can distinguish a special family of scalar-tensor theories that share properties similar to those of mimetic gravity. Mimetic gravity is a higher order scalar-tensor theory which admits, in addition to the usual invariance under diffeomorphisms, a conformal invariance (which can be generalized to a conformal-disformal invariance). Mimetic gravity was introduced in [19] as a model for dark matter (see also [20]). More recently, this model has been shown to admit (as a number of other scalar-tensor theories) non-singular bouncing cosmologies [21–25].

In the present work, we concentrate on the specific mimetic gravity Lagrangian proposed in [21]. We note that the corresponding classical equations of motion for a cosmological background are exactly the same as the so-called effective equations of loop quantum cosmology. This result suggests that it may be possible to describe loop quantum gravity at an effective level in some appropriate regimes as a higher-derivative scalar-tensor theory. The main purpose of this paper is to highlight this relation between mimetic gravity and loop quantum cosmology. While this relation has only been established at the cosmological level, it nonetheless provides a proposal for an effective description of loop quantum gravity in terms of higher-order scalar-tensor theories. Such an effective description is potentially very interesting, especially as it may give important insights into the relation between the quantization of gravity à la loop and the usual perturbative quantization techniques.

The paper is organized as follows. In the following section, we give a short presentation of (degenerate) higher-order scalar-tensor theories and we present some basic properties of mimetic gravity, which can be seen as a particular example of these theories. Then, in Sec. III, we provide a brief review of loop quantum cosmology and explain how the effective equations are derived. We then show how the loop quantum cosmology effective dynamics can be derived from an action principle $S[a, N]$ with a Lagrangian (invariant under time reparametrizations) that depends on the scale factor a and on the lapse function N , for all isotropic cosmologies. (This calculation is already known for the spatially flat case [26, 27].) In Sec. IV we show that there exists a family of DHOST mimetic actions $S[g_{\mu\nu}, \phi]$ which all reduce to $S[a, N]$ for homogeneous and isotropic space-time. These actions generalize the model proposed recently by Chamseddine and Mukhanov in [21] and can be viewed as a proposal for an effective description of loop quantum gravity. We conclude in Sec. V with a discussion.

II. HIGHER-ORDER SCALAR-TENSOR THEORIES

In this section, we briefly review the main aspects of degenerate higher-order scalar-tensor (DHOST) theories. Their Lagrangian depends on a metric $g_{\mu\nu}$ and on a scalar field ϕ , including its first and second derivatives, $\nabla_\mu \phi \equiv \phi_\mu$ and $\nabla_\mu \nabla_\nu \phi \equiv \phi_{\mu\nu}$:

$$S[g_{\mu\nu}, \phi] = \int d^4x \sqrt{-g} \mathcal{L}(\phi, \phi_\mu, \phi_{\mu\nu}; g_{\mu\nu}). \quad (2.1)$$

In general, such theories propagate an extra degree of freedom in addition to the usual scalar mode and the two tensor modes of the metric (assuming a linear dependence on the Riemann tensor¹), see (2.5) below for an explicit example of such an action. This additional degree of freedom leads to instabilities (at least at the quantum level) and is known as an Ostrogradsky ghost [7, 8].

However, it is possible to find higher-order scalar-tensor theories that do not contain any Ostrogradsky ghost by imposing appropriate degeneracy conditions on the Lagrangian, thus defining DHOST theories. DHOST theories whose Lagrangian is at most cubic in $\phi_{\mu\nu}$ have already been classified [18]. In principle, this classification could be generalized to higher powers of $\phi_{\mu\nu}$.

Below, we first recall the main properties of DHOST theories and then concentrate on mimetic theories, which form a special family within DHOST theories.

A. Evading the Ostrogradsky Instability

Starting with a Lagrangian with second derivatives of ϕ is unusual in physics. Generically, the corresponding equation of motion for ϕ is fourth order in time derivatives, which means that more than two initial conditions (per space point) are required to fully specify the evolution. This signals the presence of an extra degree of freedom in the theory.

However, there exist special Lagrangians with higher-order derivative terms for which the Euler-Lagrange equations remain second order. This is precisely the property verified by Horndeski theories in the context of scalar-tensor theories. It is even possible to find Lagrangians leading to third or fourth order Euler-Lagrange equations but without the dangerous extra scalar mode. Examples of scalar-tensor theories of this type are the so-called beyond Horndeski theories, later encompassed in the DHOST theories. All these models are degenerate, a property which can also be seen in other contexts [28–31].

By construction, DHOST theories satisfy some degeneracy conditions so that they contain at most one scalar degree of freedom. To implement this degeneracy, it is useful to work with a Hamiltonian formulation, based on the usual (3+1) ADM-decomposition of the metric on a space-time of the form $\Sigma \times \mathbb{R}$

$$g_{\mu\nu} = \begin{pmatrix} -N^2 + q_{ab}N^aN^b & q_{ab}N^b \\ q_{ab}N^a & q_{ab} \end{pmatrix}, \quad (2.2)$$

where q_{ab} is the induced metric on the space slice Σ , N is the lapse function and N^a the shift vector. In this framework, the action (2.1) explicitly depends on second time derivatives of the scalar field and takes the general form² (up to boundary terms that we neglect)

$$S[g_{\mu\nu}, \phi] = \int d^4x N \sqrt{q} \mathcal{L}(q_{ab}, K_{ab}, N, N^a; \phi, A_*, \dot{A}_*), \quad (2.3)$$

with

$$K_{ab} \equiv \frac{1}{2N} (\dot{q}_{ab} - D_a N_b - D_b N_a), \quad A_* \equiv \frac{\dot{\phi} - N^a \partial_a \phi}{N}, \quad (2.4)$$

¹ If the Lagrangian is not linear in the Riemann tensor, then the theory can admit up to 8 degrees of freedom, most of them being unstable.

² We do not consider theories which involve, after a (3+1)-decomposition, second time derivatives of the metric components which are not total derivatives. Such theories are expected to propagate Ostrogradsky ghosts that cannot be removed.

where D_a is the covariant derivative compatible with q_{ab} . For simplicity, we use the same notation for the Lagrangian densities \mathcal{L} in the covariant (2.1) and the non-covariant (2.3) versions of the action, even though they are not strictly speaking the same function.

To perform the Hamiltonian analysis [14], it is convenient to use the auxiliary variable A_* as an independent variable, so that all second time derivatives of ϕ are absorbed in A_* . This procedure thus introduces a new pair of variables, A_* and its conjugate momentum, which a priori describes an extra scalar degree of freedom. However, it is still possible that the theory propagates no more than one scalar degree of freedom if there exist constraints (in addition to the usual four constraints associated with space-time diffeomorphism invariance) so that the effective number of physical degrees of freedom is reduced. The existence of a primary constraint is equivalent to the requirement that the Hessian matrix of (2.3) (whose coefficients are the second derivatives of the action with respect to velocities of the fields) is degenerate. This property of degeneracy of the Hessian matrix has been used systematically to construct DHOST Lagrangians, initially with a quadratic dependence on $\phi_{\mu\nu}$ [13] and, more recently, with a cubic dependence [18]. Note that the primary constraint is usually of the second-class type and imposing its time conservation leads to a secondary constraint, which is also second-class. Both constraints thus eliminate the dangerous extra degree of freedom [14]. The special case where the primary constraint is first-class, signalling an additional local symmetry of the action, is seen in the mimetic models, which will be discussed in the next subsection.

All the DHOST theories that have been identified can be written in the form

$$S[g, \phi] = \int d^4x \sqrt{-g} \left[f_2(X, \phi) R + C_{(2)}^{\mu\nu\rho\sigma} \phi_{\mu\nu} \phi_{\rho\sigma} + f_3(X, \phi) G_{\mu\nu} \phi^{\mu\nu} + C_{(3)}^{\mu\nu\rho\sigma\alpha\beta} \phi_{\mu\nu} \phi_{\rho\sigma} \phi_{\alpha\beta} \right], \quad (2.5)$$

where the functions f_2 and f_3 depend only on the scalars ϕ and $X \equiv \phi_\mu \phi^\mu$; R and $G_{\mu\nu}$ denote, respectively, the usual Ricci scalar and Einstein tensor associated with the metric $g_{\mu\nu}$. The tensors $C_{(2)}$ and $C_{(3)}$ are the most general tensors constructed from the metric $g_{\mu\nu}$ and the first derivative of the scalar field ϕ_μ . It is easy to see that the quadratic terms can be rewritten as

$$C_{(2)}^{\mu\nu\rho\sigma} \phi_{\mu\nu} \phi_{\rho\sigma} = \sum_{A=1}^5 a_A(X, \phi) L_A^{(2)}, \quad (2.6)$$

with the elementary quadratic Lagrangians given by

$$\begin{aligned} L_1^{(2)} &= \phi_{\mu\nu} \phi^{\mu\nu}, & L_2^{(2)} &= (\square\phi)^2, & L_3^{(2)} &= (\square\phi) \phi^\mu \phi_{\mu\nu} \phi^\nu, \\ L_4^{(2)} &= \phi^\mu \phi_{\mu\rho} \phi^{\rho\nu} \phi_\nu, & L_5^{(2)} &= (\phi^\mu \phi_{\mu\nu} \phi^\nu)^2. \end{aligned} \quad (2.7)$$

In a similar fashion, the cubic terms can be written as

$$C_{(3)}^{\mu\nu\rho\sigma\alpha\beta} \phi_{\mu\nu} \phi_{\rho\sigma} \phi_{\alpha\beta} = \sum_{A=1}^{10} b_A(X, \phi) L_A^{(3)}, \quad (2.8)$$

with the elementary cubic Lagrangians being

$$\begin{aligned} L_1^{(3)} &= (\square\phi)^3, & L_2^{(3)} &= (\square\phi) \phi_{\mu\nu} \phi^{\mu\nu}, & L_3^{(3)} &= \phi_{\mu\nu} \phi^{\nu\rho} \phi_\rho^\mu, \\ L_4^{(3)} &= (\square\phi)^2 \phi_\mu \phi^{\mu\nu} \phi_\nu, & L_5^{(3)} &= \square\phi \phi_\mu \phi^{\mu\nu} \phi_{\nu\rho} \phi^\rho, & L_6^{(3)} &= \phi_{\mu\nu} \phi^{\mu\nu} \phi_\rho \phi^{\rho\sigma} \phi_\sigma, \\ L_7^{(3)} &= \phi_\mu \phi^{\mu\nu} \phi_{\nu\rho} \phi^{\rho\sigma} \phi_\sigma, & L_8^{(3)} &= \phi_\mu \phi^{\mu\nu} \phi_{\nu\rho} \phi^\rho \phi_\sigma \phi^{\sigma\lambda} \phi_\lambda, \\ L_9^{(3)} &= \square\phi (\phi_\mu \phi^{\mu\nu} \phi_\nu)^2, & L_{10}^{(3)} &= (\phi_\mu \phi^{\mu\nu} \phi_\nu)^3. \end{aligned} \quad (2.9)$$

In general, as explained above, theories with an action of the form (2.1) contain two tensor modes and two scalar modes, one of which leads to an Ostrogradsky instability. DHOST theories correspond to specific restrictions of the functions a_A and b_A , so that the Hessian matrix is degenerate. One finds that these theories contain at most one propagating scalar mode. DHOST theories include Horndeski and beyond-Horndeski theories but many other higher-order scalar-tensor theories as well.

B. Mimetic theories

Originally, mimetic gravity was introduced by Chamseddine and Mukhanov as a scalar-tensor theory defined by the usual Einstein-Hilbert action [19]

$$S_{CM}[\tilde{g}_{\mu\nu}, \phi] = S_{EH}[g_{\mu\nu}] \equiv \frac{1}{16\pi G} \int d^4x \sqrt{-g} R[g_{\mu\nu}] \quad (2.10)$$

where the metric $g_{\mu\nu}$ is related to $\tilde{g}_{\mu\nu}$ and ϕ by the non-invertible conformal transformation

$$g_{\mu\nu} \equiv -\tilde{X} \tilde{g}_{\mu\nu} \quad \text{with} \quad \tilde{X} \equiv \tilde{g}^{\mu\nu} \phi_\mu \phi_\nu. \quad (2.11)$$

It is immediate to see that the metric $g_{\mu\nu}$ is invariant under a local conformal transformation of the metric $\tilde{g}_{\mu\nu}$ while ϕ is left unchanged

$$\tilde{g}_{\mu\nu} \mapsto \Omega(x) \tilde{g}_{\mu\nu}, \quad \phi \mapsto \phi. \quad (2.12)$$

Here Ω is an arbitrary function on the space-time. Hence, the action (2.10) is invariant under this same local transformation provided that the coupling to matter (if there is any) is defined with respect to $g_{\mu\nu}$.

There exist two different equivalent reformulations of the mimetic action (2.10) as it was emphasized in [32]. The observation that

$$X \equiv g^{\mu\nu} \phi_\mu \phi_\nu = -\frac{1}{\tilde{X}} \tilde{g}^{\mu\nu} \phi_\mu \phi_\nu = -1, \quad (2.13)$$

enables us to replace the action (2.10) by the following one

$$S_{CM}^{(1)}[g_{\mu\nu}, \phi; \lambda] \equiv S_{EH}[g_{\mu\nu}] + \int d^4x \sqrt{-g} \lambda (X + 1), \quad (2.14)$$

where the equation of motion for λ reproduces exactly the condition (2.13). It follows that the actions (2.10) and (2.14) are classically equivalent.

In the second reformulation of mimetic gravity, one considers the action written in terms of the metric $\tilde{g}_{\mu\nu}$ and ϕ . Using the transformation law of the Ricci tensor under a conformal transformation of the metric, the action (2.10) can be rewritten as the following higher-derivative scalar-tensor theory

$$S_{CM}^{(2)}[\tilde{g}_{\mu\nu}, \phi] \equiv \frac{1}{16\pi G} \int d^4x \sqrt{-\tilde{g}} \left(\tilde{X} R[\tilde{g}_{\mu\nu}] + \frac{6}{\tilde{X}} \tilde{\phi}_\mu{}^\nu \tilde{\phi}^{\mu\rho} \phi_\nu \phi_\rho \right), \quad (2.15)$$

where $\tilde{\phi}_{\mu\nu} = \tilde{\nabla}_\mu \phi_\nu$ with $\tilde{\nabla}$ being the covariant derivative compatible with $\tilde{g}_{\mu\nu}$, and indices are lowered and raised with $\tilde{g}_{\mu\nu}$ and its inverse. The action (2.15) is clearly of the form (2.6), where the only non-trivial coefficients are

$$f_2 = \frac{\tilde{X}}{16\pi G} \quad \text{and} \quad a_4 = \frac{3}{8\pi G \tilde{X}}. \quad (2.16)$$

As expected, this theory is degenerate and can be shown to belong to the class Ia (according to the classification given in [17]) which corresponds to Lagrangians which are obtained from a conformal-disformal transformation of the quadratic Horndeski action. This is in total agreement with the fact that the mimetic gravity action has been obtained from a (non-invertible) conformal transformation of the Einstein-Hilbert action (which can be viewed as a particular case of Horndeski theory).

We close this short review with a remark concerning generalizations of mimetic gravity. First of all, one can generalize the action (2.10) assuming that $g_{\mu\nu}$ is now a general non-invertible conformal-disformal transformation of $\tilde{g}_{\mu\nu}$ [33], i.e.,

$$g_{\mu\nu} \equiv A(\tilde{X}, \phi)\tilde{g}_{\mu\nu} + B(\tilde{X}, \phi)\phi_\mu\phi_\nu \quad \text{with} \quad \frac{\partial}{\partial\tilde{X}}(A + \tilde{X}B) = 0. \quad (2.17)$$

In that case, it is easy to see that the local conformal invariance of $\tilde{g}_{\mu\nu}$ has been generalized to invariance under the local symmetry

$$\delta\tilde{g}_{\mu\nu} = \alpha(x)\tilde{g}_{\mu\nu} + \beta(x)\phi_\mu\phi_\nu \quad \text{with} \quad (A - \tilde{X}A_{\tilde{X}})\alpha(x) = \tilde{X}^2A_{\tilde{X}}\beta(x), \quad (2.18)$$

where $\alpha(x)$ and $\beta(x)$ are functions on the space-time and $A_{\tilde{X}} \equiv \partial_{\tilde{X}}A$. Hence, we obtain in this way an action $S[\tilde{g}_{\mu\nu}, \phi]$ that is invariant under the symmetry (2.18), and therefore the theory is degenerate. In fact, given any higher-order scalar-tensor action $S[g_{\mu\nu}, \phi]$, the action defined by

$$\tilde{S}[\tilde{g}_{\mu\nu}, \phi] \equiv S[g_{\mu\nu}, \phi] \quad (2.19)$$

where $g_{\mu\nu}$ is defined by (2.17) is necessarily degenerate. This family of actions provides a large generalization of mimetic gravity theories.

III. LOOP QUANTUM COSMOLOGY AND ITS EFFECTIVE DYNAMICS

As discussed in the introduction, the recent work [21] by Chamseddine and Mukhanov, based on a particular model of mimetic gravity, leads to a modified Friedmann equation that coincides with the effective dynamics of loop quantum cosmology (LQC). This connection was not mentioned in [21]. In order to clarify this link, we first briefly review LQC for spatially flat FLRW space-times and then perform a Hamiltonian analysis for a large class of mimetic gravity theories restricted to (homogeneous and isotropic) cosmological geometries. This allows us to determine the conditions on the mimetic gravity action under which the effective dynamics of LQC are recovered. Interestingly, we find that there exists a large class of such theories, among which the Chamseddine-Mukhanov model is a particular example. Furthermore, we extend this analysis to the case of spatially curved FLRW space-times and find, once again, that there exists a large class of higher-derivative scalar-tensor theories which give the LQC effective dynamics for all FLRW space-times.

A. Loop Quantum Cosmology: A Review

LQC is a proposal for quantizing cosmological space-times using the variables and the non-perturbative techniques of LQG. More specifically, as in the full theory, LQC is based on the Ashtekar-Barbero connection variables, and the elementary variables to be promoted to fundamental operators in the quantum theory are holonomies along edges and fluxes across surfaces. For dedicated reviews of LQC, see, e.g., [1–4].

1. *Hamiltonian Framework with Ashtekar-Barbero Variables*

The Ashtekar-Barbero variables are related to the metric and extrinsic curvature as follows. We first introduce the densitized triads $E_i^a = \sqrt{q}e_i^a$ (where q is the determinant of the spatial metric and the triads e_i^a are related to the inverse of the spatial metric by $q^{ab} = e_i^a e_j^b \delta^{ij}$). The conjugate variable to E_i^a is the $\mathfrak{su}(2)$ -valued Ashtekar-Barbero connection

$$A_a^i = \Gamma_a^i + \gamma K_a^i, \quad (3.1)$$

where $K_a^i = K_{ab}e_i^b$ encodes the extrinsic curvature, Γ_a^i is the spin-connection such that

$$D_a e_i^b \equiv \partial_a e_i^b - \Gamma_{ac}^b e_i^c + \epsilon_{ij}{}^k \Gamma_a^j e_k^b = 0, \quad (3.2)$$

with Γ_{ac}^b being the usual Christoffel symbols on the spatial slice, and γ is the real-valued Barbero-Immirzi parameter. The symplectic structure of gravity in the Ashtekar-Barbero variables is given by the Poisson bracket

$$\{A_a^i(x), E_j^b(y)\} = 8\pi G \gamma \delta_a^b \delta_j^i \delta^{(3)}(x-y). \quad (3.3)$$

For simplicity, the matter field is often assumed to be a scalar ψ , with a Lagrangian

$$L_\psi = \int d^4x \sqrt{-g} \left[-\frac{1}{2}(\partial\psi)^2 - V(\psi) \right]. \quad (3.4)$$

In the Hamiltonian framework, ψ comes with a conjugate momentum π_ψ such that

$$\{\psi, \pi_\psi\} = 1 \quad \text{and} \quad \pi_\psi \equiv \frac{\partial L_\psi}{\partial \dot{\psi}} = \frac{\sqrt{q}}{N} \dot{\psi}. \quad (3.5)$$

The lapse function N in the metric is not a dynamical variable and, in the Hamiltonian formulation, plays the role of a Lagrange multiplier that enforces that the scalar constraint \mathcal{H} vanish, with \mathcal{H} given by [34–36]

$$\mathcal{H} = -\frac{E_i^a E_j^b}{16\pi G \gamma^2 \sqrt{q}} \epsilon^{ij}{}^k \left(F_{ab}{}^k - (1 + \gamma^2) \Omega_{ab}{}^k \right) + \frac{\pi_\psi^2}{2\sqrt{q}} + \sqrt{q} V(\psi), \quad (3.6)$$

where $F_{ab}{}^k = 2\partial_{[a} A_{b]}^k + \epsilon_{ij}{}^k A_a^i A_b^j$ is the field strength of the Ashtekar-Barbero connection, while the tensor $\Omega_{ab}{}^k = 2\partial_{[a} \Gamma_{b]}^k + \epsilon_{ij}{}^k \Gamma_a^i \Gamma_b^j$ measures the spatial curvature.

In the particular case of a spatially flat FLRW space-time, with metric

$$ds^2 = -N^2 dt^2 + a(t)^2 d\vec{x}^2, \quad (3.7)$$

the densitized triads E_i^a are given by

$$E_i^a = p \left(\frac{\partial}{\partial x^i} \right)^a \quad \text{with} \quad p = a^2, \quad (3.8)$$

and the Ashtekar-Barbero connection can be written as

$$A_a^i = c (dx^i)_a \quad \text{with} \quad c = \frac{\gamma \dot{a}}{N}, \quad (3.9)$$

since $\Gamma_a^i = 0$ because of the space-time symmetries. Therefore, the gravitational sector of the phase space is two-dimensional, and the symplectic structure, inherited from (3.3), reduces to³

$$\{c, p\} = \frac{8\pi G\gamma}{3}. \quad (3.10)$$

Moreover, the scalar constraint (3.6) becomes

$$\mathcal{H} = -\frac{3}{8\pi G\gamma^2} p^{1/2} c^2 + \frac{\pi_\psi^2}{2p^{3/2}} + p^{3/2} V(\psi). \quad (3.11)$$

Note that $\Omega_{ab}{}^k = 0$ for the spatially flat FLRW space-time. The dynamical evolution of any observable \mathcal{O} is then given by the smeared Hamiltonian constraint according to

$$\frac{d\mathcal{O}}{dt} = \{\mathcal{O}, \mathcal{C}_H\} \quad \text{with} \quad \mathcal{C}_H = \int d^3\vec{x} N\mathcal{H}. \quad (3.12)$$

Note that the spatial diffeomorphism and Gauss constraints cannot contribute to the smeared Hamiltonian constraint since they identically vanish for the choice of variables (3.8) and (3.9). Also, the spatial integral of $N\mathcal{H}$ is trivial in FLRW space-times since every term is independent of position due to homogeneity. One can use the relation (3.12) with $\mathcal{O} = p$ to recover the usual Friedmann equation.

2. Quantum Theory

The quantization of such a theory (assuming now for simplicity that $V(\psi) = 0$) following the standard Wheeler-de Witt procedure will give a quantum cosmology where the classical big-bang singularity is not resolved in any meaningful sense. Indeed, sharply-peaked wave packets closely follow the classical (singular) solutions, and the expectation value of, e.g., the energy density of ψ can become arbitrarily large [1].

The situation is markedly different in LQC for the reason that the fundamental operators of the theory are holonomies and areas, not operators corresponding to the connection A_a^i itself. For this reason, in LQC it is not possible to directly promote the symmetry-reduced scalar constraint (3.11) to an operator in the quantum theory since there is no operator corresponding to \hat{c} . Instead, it is necessary to go back one step and construct an operator corresponding to (3.6).

This can be done in two parts. First, since the p contained in the E_i^a corresponds to an area, it can directly be promoted to be an operator. Second, the $\mathfrak{su}(2)$ -valued field strength can be expressed in terms of holonomies in the same fashion as in lattice gauge theories,

$$F_{ab} \simeq \frac{h_{\square ab} - \mathbb{I}}{Ar_{\square}}, \quad (3.13)$$

where $h_{\square ab}$ is the holonomy of the connection A_a around a loop in the a - b plane, \mathbb{I} is the identity and Ar_{\square} is the area of that loop. In a lattice gauge theory, one would be interested in the limit of the right-handside of (3.13) when $Ar_{\square} \rightarrow 0$, in which case the relation (3.13) becomes exact.

³ As a technical remark, note that for the symplectic structure to be well-defined in such a space-time, it is necessary to restrict integrals to some finite region, called the fiducial cell. This restriction on the integrals acts as an infrared regulator which should be removed once the dynamics are determined by taking the limit of the fiducial cell going to the entire spatial manifold. For simplicity, here we choose the fiducial cell such that its volume with respect to the metric $ds^2 = d\vec{x}^2$ is 1.

However, this is not natural in LQC, since the spectrum of the area operator in LQG is discrete and has a minimum non-zero eigenvalue

$$\Delta = 4\sqrt{3}\pi\gamma\ell_{\text{Pl}}^2 \quad (3.14)$$

where ℓ_{Pl} is the Planck length. Therefore, what is done in LQC is to express F_{ab} in terms of the holonomy of A_a around a loop with this minimal area Δ .

More specifically, given the symmetries of the FLRW space-time, the loop \square_{ab} is assumed to be a square loop in the a - b plane. The holonomy of A_a along edges parallel transported by the vectors $(\partial/\partial x^i)^a$ is easily evaluated. Since the Ashtekar-Barbero connection $A_a = A_a^i \tau_i$ is $\mathfrak{su}(2)$ -valued⁴, it is necessary to choose a representation in which to calculate the holonomy. This is usually chosen to be the $j = 1/2$ representation. This is not only the simplest non-trivial representation, but also corresponds to the smallest excitation Δ of area possible in LQG, which is precisely the area that has been chosen for the loop \square_{ab} to have from physical grounds, as argued in the previous paragraph.

In the $j = 1/2$ representation, the τ_i can be chosen to be the Pauli matrices (up to a factor of $i/2$ in order to have the correct normalization as shown in footnote 4). Hence, the holonomy of A_a along a path parallel to $(\partial/\partial x^i)^a$ and of length ℓ with respect to the fiducial metric $d\hat{s}^2 = dx^2$, is (no sum over i)

$$h_i(\ell) = \exp\left(\int_0^\ell dx^i A_a \left(\frac{\partial}{\partial x^i}\right)^a\right) = \cos\frac{\ell c}{2}\mathbb{I} + 2\sin\frac{\ell c}{2}\tau_i. \quad (3.15)$$

Note that the fiducial metric allows to identify internal (Lie algebra) indices i, j, k, \dots with space indices a, b, c, \dots so that, from now on, we will use the same notation i, j, k, \dots to label indifferently internal and space directions. An important point here is that the length ℓ is measured with respect to the fiducial metric, and the physical length of the edge along which h_i is evaluated is given by $a\ell$ where a is the scale factor. Therefore, requiring that the physical length of h_i be $\sqrt{\Delta}$ corresponds to setting $\sqrt{p}\ell = \sqrt{\Delta} \Rightarrow \ell = \sqrt{\Delta/p}$ where we used $a = \sqrt{p}$. Then, the holonomy $h_{\square_{ij}}$ around a square loop in the x^i - x^j plane with a physical area equal to Δ (3.14) is

$$h_{\square_{ij}} = h_j(\bar{\mu})^{-1} h_i(\bar{\mu})^{-1} h_j(\bar{\mu}) h_i(\bar{\mu}), \quad (3.16)$$

where $\bar{\mu} = \sqrt{\Delta/p}$, and $Ar_{\square} = \bar{\mu}^2$.

Now, the holonomy (3.16) can be defined in the quantum theory, and therefore can be used as the operator corresponding to \hat{F}_{ab} . Then, given the field strength operator, it is now easy to define operators corresponding to the scalar constraint, and this completes the quantum theory. For the precise details concerning the Hilbert space and the Hamiltonian constraint operator (which are not necessary here for our purposes), see, e.g., [1, 3].

The resulting quantum theory resolves the big-bang singularity in a precise sense: first, there is an upper bound on the operator corresponding to the matter energy density, and second, the

⁴ The elements τ_i form a basis of the $\mathfrak{su}(2)$ Lie algebra satisfying

$$[\tau_i, \tau_j] = 2\epsilon_{ij}{}^k \tau_k \quad \text{and} \quad \text{Tr}(\tau_i \tau_j) = -2\delta_{ij}.$$

We used the notation $\epsilon_{ij}{}^k$ for the totally antisymmetric symbol with $\epsilon_{123} = +1$, indices are raised and lowered with δ_{ij} , and Tr denotes the trace in the 2-dimensional fundamental representation (known as the Killing form). In the spin-1/2 representation, the elements τ_i are represented by the 2-dimensional matrices

$$\tau_1 = \begin{pmatrix} 0 & -i \\ -i & 0 \end{pmatrix}, \quad \tau_2 = \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix}, \quad \tau_3 = \begin{pmatrix} -i & 0 \\ 0 & i \end{pmatrix}.$$

states corresponding to singular space-times (i.e., $p = a^2 = 0$) decouple from non-singular states under the action of the Hamiltonian constraint operator and thus an initial state which is non-singular will always remain non-singular. These important differences from the Wheeler-de Witt theory arise from expressing the field strength operator in terms of the holonomy of the connection around a loop of area Δ rather than directly promoting c to be an operator.

Furthermore, for the case that the scalar field ψ is massless, it provides a good relational clock and it is possible to speak of the relational evolution of the wave function $\Psi(p, \psi)$ with respect to ψ . In particular, it is possible to construct an ‘initial’ state $\Psi(p, \psi_o)$ at some ‘initial’ relational time ψ_o and to evolve it using the Hamiltonian constraint operator. One especially interesting possibility is to construct a wave packet sharply peaked around a classical configuration at a low curvature scale when general relativity can be trusted, and then evolve the wave packet towards the high-curvature regime. An important result in this case is that the wave packet remains sharply peaked throughout its entire evolution, assuming (i) it is initially sharply peaked and (ii) that the expectation value of p always remains large compared to ℓ_{Pl}^2 . For such states, the full quantum dynamics are extremely well approximated by an effective theory [37–39].

3. Effective Dynamics

The effective dynamics of LQC are obtained by expressing the classical field strength F_{ab} in terms of the holonomy around a square loop of area Δ as we did in the previous section, and then treating the resulting \mathcal{H} classically. It is easy to verify that the LQC effective scalar constraint is

$$\mathcal{H}^{\text{eff}} = -\frac{3p^{3/2}}{8\pi G\Delta\gamma^2} \sin^2 \bar{\mu}c + \frac{\pi_{\psi}^2}{2p^{3/2}} + p^{3/2}V(\psi). \quad (3.17)$$

In this way, the effective theory captures the physics corresponding to the discrete nature of geometry in LQC (specifically, the existence of the area gap Δ), but ignores the effect of quantum fluctuations since \mathcal{H}^{eff} is treated classically⁵.

Using (3.12) with (3.17), it is easy to derive the time derivative of p :

$$\dot{p} = 2N \frac{p}{\gamma\sqrt{\Delta}} \sin \bar{\mu}c \cos \bar{\mu}c. \quad (3.18)$$

Moreover, the constraint (3.17) implies

$$\sin^2 \bar{\mu}c = \frac{\rho}{\rho_c}, \quad (3.19)$$

where ρ is the matter energy density,

$$\rho = \frac{\pi_{\psi}^2}{2p^3} + V(\psi), \quad (3.20)$$

and ρ_c is a constant defined by

$$\rho_c = \frac{3}{8\pi G\gamma^2\Delta} = \frac{\sqrt{3}}{32\pi^2\gamma^3}\rho_{\text{Pl}}, \quad (3.21)$$

⁵ This is a good approximation for states that are initially sharply peaked (i.e., all expectation values satisfy $\langle \hat{O}^2 \rangle \approx \langle \hat{O} \rangle^2$ which means that quantum fluctuations are negligible) so long as the spatial volume of the space-time is much larger than the Planck volume since for these states quantum fluctuations do not grow significantly and hence always remain negligible [39].

with ρ_{Pl} being the Planck density. Combining the above relations yields the effective Friedmann equation

$$H^2 = \left(\frac{\dot{p}}{2Np} \right)^2 = \frac{8\pi G}{3} \rho \left(1 - \frac{\rho}{\rho_c} \right), \quad (3.22)$$

where $H = \dot{a}/(Na)$ is the Hubble parameter. It can also be checked that the continuity equation in effective LQC is the same as in classical general relativity.

As can easily be seen from the LQC effective Friedmann equation, the big-bang singularity of general relativity is replaced by a bounce that occurs when the energy density of ψ reaches the critical energy density $\rho_c \propto \rho_{\text{Pl}}$ (3.21). This bounce clearly originates from the quantum geometry of LQG, and in the limit of $\Delta \rightarrow 0$, the classical Friedmann equation is recovered. Numerical simulations of the dynamics generated by the LQC Hamiltonian constraint operator for sharply-peaked wave functions have explicitly shown that the effective equations do indeed provide an excellent approximation to the full quantum dynamics, even around and at the bounce point [37]. Thus, it is clear that the bounce occurs due to the discrete geometry of LQG, irrespective of quantum fluctuations.

Note that the bounce occurs when the energy density is of the order of the Planck scale but the volume of the spatial slice at the bounce time can be much larger than the Planck volume. Consequently, as long as the bounce occurs at a spatial volume much larger than ℓ_{Pl}^3 , the effective theory can be trusted at all times for sharply-peaked states. Note that this condition is automatically satisfied for non-compact FLRW space-times since their spatial volume is always infinite (so long as the scale factor remains non-vanishing, which is always true in LQC).

In conclusion, the LQC effective dynamics is a powerful tool which significantly simplifies calculations of quantum gravity effects in semi-classical cosmological states. Clearly, it would be very helpful if it were possible to develop an effective theory that would hold more generally, for instance for all states in LQG that have nice semi-classical properties and whose geometric observables of interest concern (spatial) regions that are much larger than ℓ_{Pl}^3 . For this reason, we explore scalar-tensor theories that are able to reproduce the LQC effective dynamics for cosmological geometries.

B. Loop Quantum Cosmology from Mimetic Gravity

The goal now is to find a family of modified gravity theories that, when restricted to the spatially flat FLRW space-time, reproduce precisely the LQC effective Friedmann equation (3.22). In fact, one such modified gravity theory with precisely this property has already been found [26, 27] (see also [40] for an $f(R)$ modified gravity theory whose dynamics are a good approximation to the LQC Friedmann equation). Here, we will generalize these earlier results to a whole class of scalar-tensor theories, and in Sec. III C we extend these results to the case of non-vanishing spatial curvature.

To begin, we look for an action $S[a, N, \psi]$ where the dynamical variables are the scale factor $a(t)$, the lapse function $N(t)$ and a field $\psi(t)$ that represents the matter content of the universe. The action is of course invariant under time reparametrizations. Afterwards, we will construct a class of covariant actions of modified gravity which reduce to $S[a, N, \psi]$ when the metric is fixed by the flat FLRW metric (3.7). We assume that the field ψ is a massless scalar field minimally coupled to gravity. Hence, the modified action of gravity we are looking for takes the form

$$\int d^4x \sqrt{|g|} \left(\frac{1}{16\pi G} R - \frac{1}{2} g^{\mu\nu} \psi_\mu \psi_\nu + \dots \right), \quad (3.23)$$

where the remaining (so far unknown) part does not involve the matter content represented here

by ψ . As a consequence, on an FLRW space-time, the previous action reduces to

$$S[a, N, \psi] = \int dt \left(-\frac{3a\dot{a}^2}{8\pi GN} + a^3 \frac{\dot{\psi}^2}{2N} + Na^3 \mathcal{L} \left(a, \frac{\dot{a}}{N} \right) \right) \quad (3.24)$$

where the unknown function \mathcal{L} has to be fixed in such a way that $S[a, N, \psi]$ reproduces the LQC effective dynamics. The fact that \mathcal{L} depends on \dot{a}/N (rather than on \dot{a} and N separately) is a consequence of requiring invariance under time reparametrization. Furthermore, the Lagrangian does not involve non-trivial higher derivatives (which cannot be eliminated from the action with integrations by parts) of the scale factor, otherwise the associated classical equations of motion would (necessarily) be higher order, hence they would not reproduce the LQC effective dynamics.

As can be seen from (3.17), up to an overall prefactor the effective Hamiltonian constraint of loop quantum cosmology is expressed only in terms of the combination c/\sqrt{p} , which classically corresponds to the Hubble rate $H = \dot{a}/(Na)$ (up to the prefactor $\gamma\sqrt{\Delta}$), as can be seen from the definitions (3.8) and (3.9). This suggests restricting the function \mathcal{L} to be to the form

$$\mathcal{L} \left(a, \frac{\dot{a}}{N} \right) = \mathcal{F}(H) . \quad (3.25)$$

A Hamiltonian analysis of the action (3.24) with $\mathcal{L} = \mathcal{F}(H)$ clarifies the link with LQC. Due to the invariance under time reparametrization, the lapse N is still a Lagrange multiplier and the only non-trivial pairs of canonically conjugate pairs of variables are

$$\{a, \pi_a\} = 1 = \{\psi, \pi_\psi\} . \quad (3.26)$$

The Lagrangian is clearly not degenerate and the momenta are given in terms of the velocities by

$$\pi_a = a^2 \left[-\frac{3H}{4\pi G} + a^2 \mathcal{F}'(H) \right] , \quad \pi_\psi = \frac{a^3}{N} \dot{\psi} , \quad (3.27)$$

where \mathcal{F}' is the derivative of the function \mathcal{F} . The shape of the LQC effective Hamiltonian suggests the ansatz

$$\pi_a = \alpha a^n \arcsin \left(\beta \frac{\dot{a}}{Na} \right) , \quad (3.28)$$

where n , α and β are constants to be fixed. The condition that the momentum π_a should be approximately given by the classical result $\pi_a = -3a\dot{a}/4\pi GN$ at low curvatures (or small H) sets $n = 2$ and further requires that $\alpha\beta = -3/4\pi G$. Then, for this ansatz (3.28) for π_a , \mathcal{F} must be

$$\mathcal{F}(H) = \alpha H \arcsin(\beta H) + \frac{\alpha}{\beta} \sqrt{1 - \beta^2 H^2} + \frac{3H^2}{8\pi G} - \frac{\alpha}{\beta} , \quad (3.29)$$

where the integration constant has been fixed so that $\mathcal{F}(0) = 0$, which means that one recovers the standard general relativity action in the low curvature regime.

Given this explicit form for the Lagrangian, one finds for the Hamiltonian density

$$\mathcal{H} = a^3 \left(\frac{\pi_\psi^2}{2a^6} - \frac{8\pi G}{3} \alpha^2 \sin^2 \left(\frac{\pi_a}{2\alpha a^2} \right) \right) . \quad (3.30)$$

Using the same procedure as in the previous subsection, one easily finds that the modified Friedman equation is given by

$$H^2 = \frac{8\pi G}{3} \rho \left(1 - \frac{\rho}{\rho_c} \right) \quad \text{with} \quad \rho = \frac{\pi_\psi^2}{2a^6} \quad \text{and} \quad \rho_c = \frac{8\pi G}{3} \alpha^2 . \quad (3.31)$$

This coincides with the LQC effective dynamics provided

$$\alpha = \frac{3}{8\pi G\gamma\sqrt{\Delta}}. \quad (3.32)$$

Hence, α is determined by Newton's constant and the Barbero-Immirzi parameter. We recover the result of Chamseddine and Mukhanov from a Hamiltonian point of view. Let us emphasize that exactly the same function has been found in a rather different context much earlier in [26, 27] from a Lagrangian point of view.

Before showing the large class of scalar-tensor theories whose Hamiltonian constraint reduces to (3.30) for spatially flat FLRW space-times, we will show that this result can be extended to allow for non-vanishing spatial curvature.

C. Spatial Curvature

It is possible to generalize the previous procedure to the case of a spatially curved FLRW space-time. In classical general relativity, one gets an additional contribution from the Einstein-Hilbert term coming due to the 3-dimensional curvature 3R evaluated in a non-flat homogeneous and isotropic space-time. Hence, we start with the action

$$S_k[a, N, \psi] = \int dt \left(-\frac{3a\dot{a}^2}{8\pi GN} + a^3 \frac{\dot{\psi}^2}{N} + \frac{3Nka}{8\pi G} + Na^3 \mathcal{L}_k \left(a, \frac{\dot{a}}{N} \right) \right), \quad (3.33)$$

where k denotes the usual spatial curvature parameter. As in previous subsection, the extra term \mathcal{L}_k (such that $\mathcal{L}_0 = \mathcal{L}$) contains the (yet unknown) additional terms that are necessary to obtain the LQC effective dynamics in place of the classical general relativity ones.

1. Curved Cosmology in Mimetic Gravity

We now assume that the curvature dependence can be taken into account by simply adding to \mathcal{L} obtained in the flat case a new term that depends on the scale factor but not on its derivatives:

$$\mathcal{L}_k \left(a, \frac{\dot{a}}{N} \right) = \mathcal{F}(H) - \frac{3}{8\pi G} V_k(a). \quad (3.34)$$

The potential-like term V_k must vanish for $k = 0$, in order to recover the results of the flat case. The overall normalization is chosen for later convenience.

Since the two new terms in the action (3.33) depend only on a (and not on \dot{a}), it is straightforward to generalize the Hamiltonian analysis of the flat case. The new Hamiltonian constraint is given by

$$\mathcal{H} = a^3 \left(\rho - \rho_c \sin^2 \left(\frac{\pi a}{2\alpha a^2} \right) - \frac{3k}{8\pi G a^2} + \frac{3V_k(a)}{8\pi G} \right), \quad \rho = \frac{\pi^2 \dot{\psi}^2}{2a^6}, \quad (3.35)$$

and then the modified Friedmann equation becomes

$$H^2 = \left(\frac{8\pi G}{3} \rho - \frac{k}{a^2} + V_k(a) \right) \left(1 - \frac{1}{\rho_c} \left[\rho - \frac{3k}{8\pi G a^2} + \frac{3V_k(a)}{8\pi G} \right] \right). \quad (3.36)$$

One can now compare this modified Friedmann equation, based on the very simple ansatz (3.34), with the LQC effective Friedmann equation obtained for spatially curved FLRW space-times.

2. Curved Cosmology in Effective LQC and Quantization Ambiguities

Before making this comparison, it is important to review a quantization ambiguity which arises when performing the loop quantization of a homogeneous space-time with non-vanishing spatial curvature: this quantization ambiguity concerns the precise quantity that is to be expressed in terms of Planck-length holonomies. Due to this quantization ambiguity, there exist three quantization prescriptions that give slightly different effective theories, which can each be compared to (3.36).

To be more specific concerning this quantization ambiguity, for closed FLRW space-times there exists a direct generalization of the procedure followed for the spatially flat space-time reviewed above in Sec. III A 2, i.e., to express the field strength in terms of the holonomy of the Ashtekar-Barbero connection around a closed loop [41, 42]. This is known as the ‘F’ loop quantization. The ‘F’ loop quantization is not possible for the open FLRW space-time, nor for spatially curved Bianchi space-times (the problem is that, when expressing the field strength in this fashion, the resulting function is not almost-periodic in the connection, and so cannot be promoted to be an operator in the quantum theory, see [43–45] for details). For the Bianchi space-times, an alternative way forward is to express the connection itself (rather than the field strength) in terms of a Planck-length holonomy, as proposed in [44, 45]; this alternative quantization is known as the ‘A’ loop quantization. However, for the open FLRW space-time, neither the ‘F’ nor the ‘A’ loop quantizations are viable. Instead, it has been proposed that the ‘K’ quantization, where one considers ‘holonomies’ of the extrinsic curvature rather than of the Ashtekar-Barbero connection, may provide a reasonable approximation to a proper loop quantization [43]. Interestingly, the ‘F’, ‘A’, and ‘K’ loop quantizations are all possible in the closed FLRW model, and it is possible to compare the resulting effective theories that result from each of these quantization prescriptions [46, 47], with the surprising result that the ‘K’ quantization in fact appears to provide a better approximation to the ‘F’ loop quantization than the ‘A’ loop quantization does. (All three quantizations are also possible for the spatially flat FLRW space-time and also Bianchi type I models, but in both cases all three quantizations turn out to be exactly equivalent.) Therefore, for the closed FLRW space-time it is possible to compare the effective theories resulting from the ‘F’, ‘A’ and ‘K’ quantization prescriptions to (3.36), while for the open space-time only the effective theory resulting from the ‘K’ quantization is known.

Let us start with the ‘F’ loop quantization. In this case, the LQC effective Friedmann equation is [41, 46]

$$H^2 = \left(\frac{8\pi G}{3} \rho - \frac{k}{a^2} - \frac{k}{\gamma^2 a^2} + \frac{1}{\Delta \gamma^2} \sin^2 \frac{\sqrt{k\Delta}}{a} \right) \times \left(1 - \frac{1}{\rho_c} \left[\rho - \frac{3k}{8\pi G a^2} - \frac{3k}{8\pi G \gamma^2 a^2} + \frac{3}{8\pi G \Delta \gamma^2} \sin^2 \frac{\sqrt{k\Delta}}{a} \right] \right), \quad (3.37)$$

which is precisely of the form of (3.36) with

$$V_k(a) = -\frac{k}{\gamma^2 a^2} + \frac{k}{\Delta \gamma^2} \sin^2 \frac{\sqrt{k\Delta}}{a}. \quad (3.38)$$

Note that it is not guaranteed that the effective Friedmann equation has the general form of (3.36) for some $V_k(a)$, and therefore this result is encouraging because it indicates that the same modified gravity theory can be used to describe the LQC effective dynamics of both spatially flat and closed FLRW space-times (assuming the ‘F’ quantization for the closed FLRW space-time).

Now, while it is not known how to perform a proper loop quantization for the open FLRW space-time, it appears reasonable to assume that the effective dynamics for such a loop quantization would

also have the form (3.36) where $V_k(a)$ is (3.38) with the only difference that now k is negative. If this is indeed the case, then the resulting LQC effective Friedmann equation for the open FLRW space-time would be

$$H^2 = \left(\frac{8\pi G}{3} \rho + \frac{|k|}{a^2} + \frac{|k|}{\gamma^2 a^2} - \frac{1}{\Delta \gamma^2} \sinh^2 \frac{\sqrt{|k|\Delta}}{a} \right) \times \left(1 - \frac{1}{\rho_c} \left[\rho + \frac{3|k|}{8\pi G a^2} + \frac{3|k|}{8\pi G \gamma^2 a^2} - \frac{3}{8\pi G \Delta \gamma^2} \sinh^2 \frac{\sqrt{|k|\Delta}}{a} \right] \right). \quad (3.39)$$

Moving on to the ‘A’ loop quantization, the effective equation for the closed FLRW space-time turns out to have a form which is different than (3.36) [46]. This provides an explicit example that shows that, in general, generic modifications to the Friedmann equations cannot be written in the form (3.36). It is only in some special cases that it will be possible to describe quantum gravity effects via a mimetic scalar-tensor theory, as is the case for the ‘F’ loop quantization.

Finally, for the ‘K’ loop quantization the effective Friedmann equation for the closed and open FLRW space-times is [43, 47]

$$H^2 = \left(\frac{8\pi G}{3} \rho + \frac{k}{a^2} \right) \times \left(1 - \frac{1}{\rho_c} \left[\rho + \frac{3k}{8\pi G a^2} \right] \right), \quad (3.40)$$

where $k > 0$ for a closed universe and $k < 0$ for an open universe. Interestingly, in this case we find that the effective Friedmann equation is again of the form (3.36), this time with $V_k(a) = 0$. There are three important points here. First, for the ‘K’ quantization, which can be performed for both open and closed FLRW space-times, the effective theory can be understood to come from a scalar-tensor theory, and the same theory describes equally well space-times with positive or negative spatial curvature. Second, in agreement with earlier results [46, 47], we find that the ‘K’ quantization is more similar to the ‘F’ quantization than the ‘A’ quantization is, in that its resulting effective Friedmann equations can be described by a scalar-tensor theory. And third, the ‘K’ quantization does not require a $V_k(a)$ term in the action (3.33). Since a non-vanishing $V_k(a)$ term breaks Lorentz invariance (as we shall discuss in more detail in the following section), it is quite interesting that the effective theory of the ‘K’ quantization in fact corresponds to a modified gravity theory which is Lorentz invariant.

To summarize, there is a quantization ambiguity that arises in LQC when considering spatially curved homogeneous space-times. There exist ‘F’, ‘A’ and ‘K’ quantizations for the closed FLRW space-time, while only the ‘K’ quantization is known for the open topology. For closed FLRW space-time, the effective equations resulting from the ‘F’ quantization prescription (but not the ‘A’ quantization) can be understood to come from a mimetic gravity theory with the action (3.33), for a particular choice of $V_k(a)$. This mimetic gravity theory in fact provides a candidate effective theory for a proper loop quantization of the open FLRW space-time. Finally, the effective equations for the ‘K’ quantization of FLRW space-times (for all k) can also be understood to follow from a mimetic gravity action of the form (3.33), in this case with $V_k(a) = 0$.

In short, these results show that the LQC effective equations (with the exception of the ‘A’ quantization) for homogeneous and isotropic cosmologies are identical to the Friedmann equations of a particular mimetic gravity theory.

IV. EFFECTIVE LOOP QUANTUM GRAVITY AND MIMETIC GRAVITY

In this section, we address the question of finding a covariant action of modified gravity that reduces to

$$S[a, N, \psi] = \int dt N a^3 \left(\frac{\dot{\psi}^2}{2N^2} - \frac{\rho_c}{2} \left[\beta H \arcsin(\beta H) + \sqrt{1 - \beta^2 H^2} - 1 \right] \right), \quad \beta^2 = \frac{3}{2\pi G \rho_c}, \quad (4.1)$$

in a spatially flat homogeneous and isotropic space-time (the non-flat case will be discussed below), as found in the previous section. Of course, such a condition is not very restrictive and one can expect that many different covariant actions could yield the same cosmological action.

Because of the non-linearity of the Lagrangian in the Hubble parameter H , it is not possible to find an $f(R)$ theory which exactly reproduces (4.1) in the cosmological sector, although an approximate construction has been found in [40] (where one considers $f(R)$ theories à la Palatini). The reason is that the Ricci scalar, given by

$$R = 6 \left(\frac{1}{a} \frac{1}{N} \frac{d}{dt} \left(\frac{\dot{a}}{N} \right) + \left(\frac{\dot{a}}{Na} \right)^2 \right), \quad (4.2)$$

involves second derivatives of the scale factor and this prevents us from recovering a Lagrangian $f(R)$ that depends only on a and \dot{a} in the cosmological sector.

Another possibility would be to consider an action involving nonlinear combinations of the Riemann tensor. It was shown in [27] that a Lagrangian containing contractions of the Ricci tensor of the form $R_{\mu_1}{}^{\mu_2} R_{\mu_2}{}^{\mu_3} \dots R_{\mu_n}{}^{\mu_1}$ indeed reduces to (4.1) for spatially flat FLRW space-times. However, the explicit expression of this Lagrangian is very cumbersome and it does not appear to be suited for calculations away from the homogeneous and isotropic sector. Furthermore, as the Lagrangian involves higher powers of the Ricci tensor, the theory will propagate more degrees of freedom than the usual two tensor modes, and these additional degrees of freedom will lead to Ostrogradsky instabilities.

Following [21], we will try to explore scalar-tensor actions, in particular mimetic theories. Following the results recalled in the previous sections, we look for a covariant action of the form

$$S[g_{\mu\nu}, \phi] = \int d^4x \sqrt{-g} \left(\frac{fR}{16\pi G} + \mathcal{L}_\phi(\phi, \phi_\mu, \phi_{\mu\nu}) + \lambda(X + 1) - \frac{1}{2} g^{\mu\nu} \psi_\mu \psi_\nu \right) \quad (4.3)$$

where f is a function of ϕ only and \mathcal{L}_ϕ is a scalar function which depends on ϕ and its first and second derivatives ϕ_μ and $\phi_{\mu\nu}$ only.

As we said above, the solution of our problem is far from being unique and one can find an infinite class of solutions with few restrictions on the Lagrangian \mathcal{L}_ϕ . Here, we want to propose the simplest class of solutions which generalize the Chamseddine-Mukhanov model, and we restrict ourselves to higher-derivative Lagrangians for the scalar field of the form

$$\mathcal{L}_\phi = \mathcal{L}_\phi \left(\phi, L_1^{(2)}, L_2^{(2)}, L_3^{(2)}, L_4^{(2)}, L_5^{(2)} \right) \quad (4.4)$$

where the $L_A^{(2)}$ are the five elementary quadratic Lagrangians introduced in (2.7). Of course, one could also consider further generalizations including the cubic elementary Lagrangians.

Now, we find the conditions that f and \mathcal{L}_ϕ must satisfy for the action (4.3) to reduce to (3.24)

when the metric $g_{\mu\nu}$ is (3.7) and the fields ϕ and ψ depend on time only. In this case,

$$L_1^{(2)} = (\mathcal{D}_t^2 \phi)^2 + 3 \left(\mathcal{D}_t \phi \frac{\mathcal{D}_t a}{a} \right)^2, \quad L_2^{(2)} = \left[\frac{1}{a^3} \mathcal{D}_t (a^3 \mathcal{D}_t \phi) \right]^2, \quad (4.5)$$

$$L_3^{(2)} = -\frac{1}{a^3} (\mathcal{D}_t \phi)^2 \mathcal{D}_t^2 \phi \mathcal{D}_t (a^3 \mathcal{D}_t \phi), \quad (4.6)$$

$$L_4^{(2)} = -(\mathcal{D}_t \phi \mathcal{D}_t^2 \phi)^2, \quad L_5^{(2)} = \left[(\mathcal{D}_t \phi)^2 \mathcal{D}_t^2 \phi \right]^2, \quad (4.7)$$

where we have introduced the notation

$$\mathcal{D}_t \varphi \equiv \frac{\dot{\varphi}}{N} \quad (4.8)$$

for any time-dependent function $\varphi(t)$.

The advantage to consider mimetic theories is that the time derivative of the scalar field is automatically normalized since we have, in the cosmological background,

$$X = - \left(\frac{\dot{\phi}}{N} \right)^2 = -1. \quad (4.9)$$

As consequence, in the cosmological context, the elementary quadratic Lagrangians simplify drastically:

$$L_1^{(2)} = 3 \left(\frac{\dot{a}}{Na} \right)^2, \quad L_2^{(2)} = 9 \left(\frac{\dot{a}}{Na} \right)^2, \quad L_3^{(2)} = L_4^{(2)} = L_5^{(2)} = 0. \quad (4.10)$$

Due to the form of (3.24), the functions f and \mathcal{L}_ϕ are necessarily independent of ϕ . All these ingredients allow us to conclude immediately that the action (4.3) reproduces the LQC effective dynamics when the Lagrangian \mathcal{L}_ϕ takes the form

$$\mathcal{L}_\phi(L_1^{(2)}, L_2^{(2)}, L_3^{(2)}, L_4^{(2)}, L_5^{(2)}) = \left(\sum_{i=1}^5 \alpha_i L_i^{(2)} \right) + U(L_1^{(2)}, L_2^{(2)}, L_3^{(2)}, L_4^{(2)}, L_5^{(2)}) \quad (4.11)$$

with the conditions

$$\alpha_1 + 3\alpha_2 = \frac{f}{8\pi G}, \quad U(3H^2, 9H^2, 0, 0, 0) = \frac{\rho_c}{2} \left(1 - \sqrt{1 - \beta^2 H^2} - \beta H \arcsin(\beta H) \right), \quad (4.12)$$

for any value of H . The mimetic gravity model proposed by Chamseddine and Mukhanov clearly belongs to this class and consists in taking $f = 24\pi G \alpha_2 = 1$, with all other $\alpha_A = 0$, and U is assumed to be a function of $L_2^{(2)} = (\square\phi)^2$ only. However, this result shows that there exist a lot of different possibilities to get the LQC effective dynamics.

Let us close this section with a discussion on the case of a non-flat cosmology. The only difference with what has been done so far is that the scalar-tensor action, when evaluated for a spatially curved FLRW space-time, must include the additional $V_k(a)$ term that arises in (3.34) (at least for the ‘F’ loop quantization, but not for the ‘K’ quantization, as explained in III C 2). Such a term can be obtained by adding to the effective covariant action (4.3) with \mathcal{L}_ϕ given by (4.11) a new potential term

$$- \int d^4x \sqrt{|g|} \mathcal{V}(g_{\mu\nu}, \phi) \quad (4.13)$$

which reproduces exactly $-V_k(a)$ when evaluated on a curved FLRW space-time. As in the previous situation, we obviously do not expect the solution for \mathcal{V} to be unique. However, we can exhibit some properties \mathcal{V} must have. The fact that $V_k(a)$ is a highly non-linear function (even non-polynomial) of the scale factor a which does not depend on its time derivatives implies that \mathcal{V} cannot be constructed from the full space-time components of the Riemann tensor only (with no contractions with derivatives of ϕ) nor from the elementary scalar-tensor Lagrangians $L_A^{(2)}$. The reason is that, when evaluated on a curved FLRW space-time, such terms necessarily produce derivatives of the scale factor which cannot be removed by integration by parts. Cubic (or even higher-order) scalar-tensor terms as $L_A^{(3)}$ produce the same problem. Only the components of the 3-dimensional Riemann tensor are functions of a only when evaluated on a curved FLRW solution. This can be easily illustrated with the 3-dimensional Ricci scalar 3R which reduces to

$${}^3R = -\frac{6k}{a^2}. \quad (4.14)$$

From this point of view, the potential $\mathcal{V}(g_{\mu\nu}, \phi)$ can be constructed from the 3-dimensional Riemann tensor, in which case it breaks the Lorentz invariance. Furthermore, as V_k can be expanded as the following series

$$V_k(a) = \sum_{n>0} v_n \left(\frac{6k}{a^2}\right)^n \equiv \tilde{V}\left(-\frac{6k}{a^2}\right), \quad (4.15)$$

one can choose for the potential \mathcal{V}_k a function which depends only on 3R according to

$$\mathcal{V}(g_{\mu\nu}, \phi) = \tilde{V}({}^3R). \quad (4.16)$$

Interestingly, similar potential are considered in Hořava-Lifshitz-type models of gravity to construct renormalizable theories of gravity. Note that, using the Stückelberg method, it is nonetheless possible to render \mathcal{V} fully covariant using the scalar field ϕ as a covariant clock. The price to pay is that the covariant version of \mathcal{V} would involve terms like $R_{\mu\nu\rho\sigma}\phi^\mu\phi^\nu\phi^\rho\phi^\sigma$ in the action which would most probably introduce many unhealthy degrees of freedom in the theory.

V. PERSPECTIVES

In this paper, we have constructed a family of higher-derivative scalar-tensor theories that possess the property to reproduce exactly the effective dynamics of loop quantum cosmology for flat, closed and open homogeneous and isotropic space-times, leading to bouncing solutions. This family thus generalizes the particular model considered by Chamseddine and Mukhanov for the spatially flat case [21].

An important question is whether this family, identified only at the level of the homogeneous and isotropic dynamics, contains a theory that could fully represent an effective description of the full loop quantum gravity (LQG). Indeed, a study limited to homogeneous cosmologies is too restrictive to be conclusive and one should go beyond homogenous and isotropic solutions to test further the interest of the theories we have identified. In particular, it would be interesting to investigate whether some of them can adequately describe anisotropic space-times, cosmological perturbations, black holes, and more.

Concerning anisotropic space-times, it should be stressed that the equations of motion for the Bianchi I space-time derived in the specific mimetic theory of [21] do not coincide with the LQC effective equations given in [48, 49], even if their qualitative dynamics are quite similar (see, e.g.,

[50]). It would thus be worth studying whether another theory among the family identified in the present work is able to reproduce these Bianchi I equations. We leave this question for future work.

It would also be interesting to extend our study to include linear cosmological perturbations around the FLRW background. Indeed, in the presence of perturbative inhomogeneities a number of important qualitative similarities have already been noticed between some modified gravity theories and the LQC effective dynamics [51]. One might hope that there is in fact an exact correspondence for at least one mimetic theory of gravity.

Furthermore, it would be interesting to study black hole solutions for the scalar-tensor theories considered here. For the specific mimetic theory proposed in [21], the spherically symmetric black hole solution has been studied in [52]. We now have at our disposal a large class of mimetic theories to consider. In particular, it might be possible to find solutions related to those constructed in the context of LQG such as the Planck stars [53, 54].

Beyond the natural extensions discussed above, it should also be mentioned that an alternative description of the LQC effective dynamics offers a complementary framework that could provide new insights. One such example is the question of a possible signature change in LQC: due to a modification in the Dirac algebra of the (effective) constraints of LQC, it has been suggested that the signature of the metric may change from Lorentzian to Euclidean around the bounce point, see [55] for details. While this question is difficult to address in LQC due to the gauge-fixings that are necessary before quantization, the metric clearly remains Lorentzian at all times in the mimetic theories considered here and this could suggest that there is no signature change in LQC.

Finally, this result suggests some new links between LQG, Hořava-Lifshitz gravity, and non-commutative geometry. As seen in Sec. III C 2, for some (but not all) versions of LQC it is necessary to add a Lorentz-violating term in order to recover the correct effective Friedmann equations in the presence of scalar curvature, along the lines of what is done in Hořava-Lifshitz gravity. As Hořava-Lifshitz gravity is known to be perturbatively renormalizable, it is possible that this effective theory may be renormalizable as well. In addition, the mimetic condition (2.13) has been argued to arise naturally in non-commutative geometry when requiring the quantization of the three-dimensional volume [56]. It is intriguing that it is precisely modified gravity theories that satisfy this condition which give the LQC effective dynamics for isotropic cosmologies. An exploration in greater depth of the links between these different approaches to the problem of quantum gravity may provide important new insights.

Note: During the completion of this paper, we were informed that the independent work [57] also points out the relation between the mimetic gravity theory of [21] and the effective equations of LQC.

Acknowledgments

We would like to warmly thank Norbert Bodendorfer for valuable discussions and for telling us of his work [57] before its publication.

-
- [1] A. Ashtekar and P. Singh, “Loop Quantum Cosmology: A Status Report,” *Class. Quant. Grav.* **28** (2011) 213001, [arXiv:1108.0893](#).
 - [2] M. Bojowald, “Loop quantum cosmology,” *Living Rev. Rel.* **11** (2008) 4.
 - [3] K. Banerjee, G. Calcagni, and M. Martín-Benito, “Introduction to loop quantum cosmology,” *SIGMA* **8** (2012) 016, [arXiv:1109.6801](#).
 - [4] E. Wilson-Ewing, “Testing loop quantum cosmology,” [arXiv:1612.04551](#).

- [5] D. Battfeld and P. Peter, “A Critical Review of Classical Bouncing Cosmologies,” *Phys. Rept.* **571** (2015) 1–66, [arXiv:1406.2790](#).
- [6] R. Brandenberger and P. Peter, “Bouncing Cosmologies: Progress and Problems,” [arXiv:1603.05834](#).
- [7] M. Ostrogradsky, “Mémoires sur les équations différentielles, relatives au problème des isopérimètres,” *Mem. Acad. St. Petersbourg* **VI** 4 (1850) 385–517.
- [8] R. P. Woodard, “Ostrogradsky’s theorem on Hamiltonian instability,” *Scholarpedia* **10** (2015) 32243, [arXiv:1506.02210](#).
- [9] G. W. Horndeski, “Second-order scalar-tensor field equations in a four-dimensional space,” *Int. J. Theor. Phys.* **10** (1974) 363–384.
- [10] C. Deffayet, X. Gao, D. A. Steer, and G. Zahariade, “From k-essence to generalised Galileons,” *Phys. Rev.* **D84** (2011) 064039, [arXiv:1103.3260](#).
- [11] J. Gleyzes, D. Langlois, F. Piazza, and F. Vernizzi, “Healthy theories beyond Horndeski,” *Phys. Rev. Lett.* **114** (2015) 211101, [arXiv:1404.6495](#).
- [12] J. Gleyzes, D. Langlois, F. Piazza, and F. Vernizzi, “Exploring gravitational theories beyond Horndeski,” *JCAP* **1502** (2015) 018, [arXiv:1408.1952](#).
- [13] D. Langlois and K. Noui, “Degenerate higher derivative theories beyond Horndeski: evading the Ostrogradski instability,” *JCAP* **1602** (2016) 034, [arXiv:1510.06930](#).
- [14] D. Langlois and K. Noui, “Hamiltonian analysis of higher derivative scalar-tensor theories,” *JCAP* **1607** (2016) 016, [arXiv:1512.06820](#).
- [15] M. Crisostomi, M. Hull, K. Koyama, and G. Tasinato, “Horndeski: beyond, or not beyond?,” *JCAP* **1603** (2016) 038, [arXiv:1601.04658](#).
- [16] M. Crisostomi, K. Koyama, and G. Tasinato, “Extended Scalar-Tensor Theories of Gravity,” *JCAP* **1604** (2016) 044, [arXiv:1602.03119](#).
- [17] J. Ben Achour, D. Langlois, and K. Noui, “Degenerate higher order scalar-tensor theories beyond Horndeski and disformal transformations,” *Phys. Rev.* **D93** (2016) 124005, [arXiv:1602.08398](#).
- [18] J. Ben Achour, M. Crisostomi, K. Koyama, D. Langlois, K. Noui, and G. Tasinato, “Degenerate higher order scalar-tensor theories beyond Horndeski up to cubic order,” *JHEP* **12** (2016) 100, [arXiv:1608.08135](#).
- [19] A. H. Chamseddine and V. Mukhanov, “Mimetic Dark Matter,” *JHEP* **11** (2013) 135, [arXiv:1308.5410](#).
- [20] A. H. Chamseddine, V. Mukhanov, and A. Vikman, “Cosmology with Mimetic Matter,” *JCAP* **1406** (2014) 017, [arXiv:1403.3961](#).
- [21] A. H. Chamseddine and V. Mukhanov, “Resolving Cosmological Singularities,” *JCAP* **1703** (2017) 009, [arXiv:1612.05860](#).
- [22] A. Ijjas, J. Ripley, and P. J. Steinhardt, “NEC violation in mimetic cosmology revisited,” *Phys. Lett.* **B760** (2016) 132–138, [arXiv:1604.08586](#).
- [23] A. Ijjas and P. J. Steinhardt, “Classically stable nonsingular cosmological bounces,” *Phys. Rev. Lett.* **117** (2016) 121304, [arXiv:1606.08880](#).
- [24] A. Ijjas and P. J. Steinhardt, “Fully stable cosmological solutions with a non-singular classical bounce,” *Phys. Lett.* **B764** (2017) 289–294, [arXiv:1609.01253](#).
- [25] H. Firouzjahi, M. A. Gorji, and A. H. Mansoori, “Instabilities in Mimetic Matter Perturbations,” [arXiv:1703.02923](#).
- [26] G. Date and S. Sengupta, “Effective Actions from Loop Quantum Cosmology: Correspondence with Higher Curvature Gravity,” *Class. Quant. Grav.* **26** (2009) 105002, [arXiv:0811.4023](#).
- [27] R. C. Helling, “Higher curvature counter terms cause the bounce in loop cosmology,” [arXiv:0912.3011](#).
- [28] H. Motohashi, K. Noui, T. Suyama, M. Yamaguchi, and D. Langlois, “Healthy degenerate theories with higher derivatives,” *JCAP* **1607** (2016) 033, [arXiv:1603.09355](#).
- [29] R. Klein and D. Roest, “Exorcising the Ostrogradsky ghost in coupled systems,” *JHEP* **07** (2016) 130, [arXiv:1604.01719](#).
- [30] C. de Rham and A. Matas, “Ostrogradsky in Theories with Multiple Fields,” *JCAP* **1606** (2016), no. 06, 041, [arXiv:1604.08638](#).
- [31] M. Crisostomi, R. Klein, and D. Roest, “Higher Derivative Field Theories: Degeneracy Conditions and Classes,” [arXiv:1703.01623](#).

- [32] F. Arroja, N. Bartolo, P. Karmakar, and S. Matarrese, “The two faces of mimetic Horndeski gravity: disformal transformations and Lagrange multiplier,” *JCAP* **1509** (2015) 051, [arXiv:1506.08575](#).
- [33] N. Deruelle and J. Rua, “Disformal Transformations, Veiled General Relativity and Mimetic Gravity,” *JCAP* **1409** (2014) 002, [arXiv:1407.0825](#).
- [34] J. F. Barbero G., “Real Ashtekar variables for Lorentzian signature space times,” *Phys. Rev.* **D51** (1995) 5507–5510, [arXiv:gr-qc/9410014](#).
- [35] S. Holst, “Barbero’s Hamiltonian derived from a generalized Hilbert-Palatini action,” *Phys. Rev.* **D53** (1996) 5966–5969, [arXiv:gr-qc/9511026](#).
- [36] T. Thiemann, *Modern canonical quantum general relativity*, Cambridge University Press, Cambridge, 2008. [arXiv:gr-qc/0110034](#).
- [37] A. Ashtekar, T. Pawłowski, and P. Singh, “Quantum Nature of the Big Bang: Improved dynamics,” *Phys. Rev.* **D74** (2006) 084003, [arXiv:gr-qc/0607039](#).
- [38] V. Taveras, “Corrections to the Friedmann Equations from LQG for a Universe with a Free Scalar Field,” *Phys. Rev.* **D78** (2008) 064072, [arXiv:0807.3325](#).
- [39] C. Rovelli and E. Wilson-Ewing, “Why are the effective equations of loop quantum cosmology so accurate?,” *Phys. Rev.* **D90** (2014) 023538, [arXiv:1310.8654](#).
- [40] G. J. Olmo and P. Singh, “Effective Action for Loop Quantum Cosmology a la Palatini,” *JCAP* **0901** (2009) 030, [arXiv:0806.2783](#).
- [41] A. Ashtekar, T. Pawłowski, P. Singh, and K. Vandersloot, “Loop quantum cosmology of $k=1$ FRW models,” *Phys. Rev.* **D75** (2007) 024035, [arXiv:gr-qc/0612104](#).
- [42] L. Szulc, W. Kamiński, and J. Lewandowski, “Closed FRW model in Loop Quantum Cosmology,” *Class. Quant. Grav.* **24** (2007) 2621–2636, [arXiv:gr-qc/0612101](#).
- [43] K. Vandersloot, “Loop quantum cosmology and the $k = -1$ RW model,” *Phys. Rev.* **D75** (2007) 023523, [arXiv:gr-qc/0612070](#).
- [44] A. Ashtekar and E. Wilson-Ewing, “Loop quantum cosmology of Bianchi type II models,” *Phys. Rev.* **D80** (2009) 123532, [arXiv:0910.1278](#).
- [45] E. Wilson-Ewing, “Loop quantum cosmology of Bianchi type IX models,” *Phys. Rev.* **D82** (2010) 043508, [arXiv:1005.5565](#).
- [46] A. Corichi and A. Karami, “Loop quantum cosmology of $k=1$ FRW: A tale of two bounces,” *Phys. Rev.* **D84** (2011) 044003, [arXiv:1105.3724](#).
- [47] P. Singh and E. Wilson-Ewing, “Quantization ambiguities and bounds on geometric scalars in anisotropic loop quantum cosmology,” *Class. Quant. Grav.* **31** (2014) 035010, [arXiv:1310.6728](#).
- [48] D.-W. Chiou and K. Vandersloot, “The Behavior of non-linear anisotropies in bouncing Bianchi I models of loop quantum cosmology,” *Phys. Rev.* **D76** (2007) 084015, [arXiv:0707.2548](#).
- [49] A. Ashtekar and E. Wilson-Ewing, “Loop quantum cosmology of Bianchi I models,” *Phys. Rev.* **D79** (2009) 083535, [arXiv:0903.3397](#).
- [50] B. Gupt and P. Singh, “Quantum gravitational Kasner transitions in Bianchi-I spacetime,” *Phys. Rev.* **D86** (2012) 024034, [arXiv:1205.6763](#).
- [51] Y.-F. Cai and E. Wilson-Ewing, “Non-singular bounce scenarios in loop quantum cosmology and the effective field description,” *JCAP* **1403** (2014) 026, [arXiv:1402.3009](#).
- [52] A. H. Chamseddine and V. Mukhanov, “Nonsingular Black Hole,” *Eur. Phys. J.* **C77** (2017), no. 3, 183, [arXiv:1612.05861](#).
- [53] C. Rovelli and F. Vidotto, “Planck stars,” *Int. J. Mod. Phys.* **D23** (2014), no. 12, 1442026, [arXiv:1401.6562](#).
- [54] T. De Lorenzo, C. Pacilio, C. Rovelli, and S. Speziale, “On the Effective Metric of a Planck Star,” *Gen. Rel. Grav.* **47** (2015), no. 4, 41, [arXiv:1412.6015](#).
- [55] T. Cailleteau, J. Mielczarek, A. Barrau, and J. Grain, “Anomaly-free scalar perturbations with holonomy corrections in loop quantum cosmology,” *Class. Quant. Grav.* **29** (2012) 095010, [arXiv:1111.3535](#).
- [56] A. H. Chamseddine, A. Connes, and V. Mukhanov, “Quanta of Geometry: Noncommutative Aspects,” *Phys. Rev. Lett.* **114** (2015), no. 9, 091302, [arXiv:1409.2471](#).
- [57] N. Bodendorfer, A. Schäfer, and J. Schliemann, “On the canonical structure of general relativity with a limiting curvature and its relation to loop quantum gravity,” [arXiv preprint](#).